

Essential Summary

Predicting Homoclinic and Heteroclinic Bifurcation of Generalized Duffing–Harmonic–van de Pol Oscillator

Zhenbo Li, Jiashi Tang, Ping Cai

Qualitative Theory of Dynamical Systems, Vol. 15, No. 1, 19–37 (2016) DOI: 10.1007/s12346-015-0138-z

This is an author-prepared essential summary, not the original paper.
Please cite the published version.

1 Background and Motivation

The Duffing–harmonic oscillator, characterized by a rational restoring force,

$$\ddot{x} + \frac{\lambda x + \delta x^3}{1 + \nu x^2} = 0 \quad (1)$$

bridges two well-studied regimes: for small x , it behaves as a Duffing oscillator ($\ddot{x} + \delta x^3 \approx 0$); for large x , it approximates a linear harmonic oscillator ($\ddot{x} + (\delta/\nu)x \approx 0$). This dual nature makes it an excellent model for systems with amplitude-dependent stiffness transitions.

While the conservative Duffing–harmonic oscillator had been extensively studied via harmonic balance, homotopy analysis, and energy balance methods, the **damped** version—the generalized Duffing–harmonic–van de Pol oscillator—

$$\ddot{x} + \frac{\lambda x + \delta x^3}{1 + \nu x^2} = \varepsilon(\mu + \mu_1 x + \mu_2 x^2 + \mu_4 x^4)\dot{x} \quad (2)$$

had received almost no attention. This paper fills that gap by applying the quadratic generalized harmonic function perturbation method to predict both homoclinic and heteroclinic bifurcations in this oscillator.

2 Quadratic Generalized Harmonic Function

2.1 Solution Construction

Building on the original GHFP method (JSV 2013), this paper constructs solutions in the form:

$$x(t) = a \sin^2 \phi(t) + b \quad (\text{homoclinic orbits}) \quad (3)$$

$$x(t) = a \cos^2 \phi(t) + b \quad (\text{heteroclinic orbits}) \quad (4)$$

The nonlinear time transformation:

$$\frac{d\phi}{dt} = \Phi(\phi), \quad \Phi(\phi + \pi) = \Phi(\phi) \quad (5)$$

maps the time domain to an angular domain where the solutions have a symmetric form.

2.2 Homoclinic Orbits

For a saddle point $H(h, 0)$, the initial condition c satisfies:

$$\int_h^c \frac{\lambda x + \delta x^3}{1 + \nu x^2} dx = 0 \quad (6)$$

and the homoclinic solution must satisfy $x(\phi_{\pm\infty}) = h$. The angular velocity is obtained as:

$$\Phi(\phi) = \sqrt{\frac{2[V(b) - V(a \sin^2 \phi + b)]}{(x'(\phi))^2}} \quad (7)$$

with $\Phi(0) = \Phi(\pi) = \sqrt{|g(h)/2a|} = 0$ ensuring infinite period at the saddle point.

2.3 Heteroclinic Orbits

For two distinct saddle points $H_1(h_1, 0)$ and $H_2(h_2, 0)$:

$$\lim_{t \rightarrow -\infty} x(t) = h_1, \quad \lim_{t \rightarrow +\infty} x(t) = h_2 \quad (8)$$

The constants a and b are determined by $\phi = 0 \rightarrow x = h_1$ and $\phi = \pi/2 \rightarrow x = h_2$. The angular velocity satisfies $\Phi(0) = \Phi(\pi/2) = 0$.

3 Perturbation Procedure

3.1 Expansion and Key Assumption

When $\varepsilon \neq 0$:

$$a = a_0 + \varepsilon a_1 + \dots, \quad \Phi = \Phi_0 + \varepsilon \Phi_1 + \dots \quad (9)$$

The determining insight: **the n th-order approximate solution maintains the same functional form as the generating solution:**

$$x_n = a_n \sin^2 \phi \quad (\text{homoclinic}) \quad (10)$$

$$x_n = a_n \cos^2 \phi \quad (\text{heteroclinic}) \quad (11)$$

This eliminates the cumbersome integrals that plague classical perturbation methods.

3.2 Homoclinic Bifurcation Prediction

The first-order consistency condition yields:

$$\mu_c = - \frac{\int_0^\pi \Phi_0(\mu_1 x_0 + \mu_2 x_0^2 + \mu_4 x_0^4)(\Phi_0 x'_0) x'_0 d\varphi}{\int_0^\pi \Phi_0^2(x'_0)^2 d\varphi} \quad (12)$$

This formula directly predicts the critical parameter value μ_c at which homoclinic bifurcation occurs.

4 Key Results

4.1 Example 1: Duffing–Harmonic Oscillator with (μ_1, μ_2, μ_4) Damping

For the system with $\lambda = \delta = \nu = 1$, the GHFP method accurately predicts homoclinic orbits across a wide range of ε values.

4.2 Example 2: Multiple Coexisting Homoclinic Orbits

For a system with different parameters, the method captures two types of homoclinic orbits—one enclosing a single well and another enclosing both wells—at the same parameter values.




Figure 1: **Figure 1.** (Original Fig. 1) Homoclinic orbits for $\varepsilon = 1.2, 2.0$. Solid lines: Runge–Kutta method. Dotted lines: GHFP method. Excellent agreement demonstrates the effectiveness of the quadratic generalized harmonic function approach for rational restoring forces.




Fig. 2 Homoclinic orbit of oscillator (65) for **a** $\varepsilon = 0.5$, **b** $\varepsilon = 1.0$, **(c)** $\varepsilon = 1.5$ and **(d)** $\varepsilon = 2.0$. *Continue* denotes the orbits obtained by Runge–Kutta method; *dotted line* denotes the orbits obtained by present method




Figure 2: **Figure 2.** (Original Fig. 2) Homoclinic orbits for $\varepsilon = 0.5, 1.0, 1.5, 2.0$. The method successfully tracks the orbits across a wide range, maintaining accuracy even at $\varepsilon = 2.0$.




Figure 3: **Figure 3.** (Original Fig. 4) Heteroclinic orbits connecting two distinct saddle points for $\varepsilon = 0.5, 0.8, 1.0, 2.0$. The method captures cross-saddle connections with remarkable accuracy.

4.3 Example 3: Heteroclinic Bifurcation Prediction

For the heteroclinic case, the method predicts critical parameter values:

$$\mu_c = -0.1979871606 \quad (\varepsilon = 0.5), \quad \mu_c^{(\text{numerical})} = -0.1977132821 \quad (13)$$

The relative error is approximately 0.14%.

5 Significance

1. **First application of GHFP to rational restoring forces:** Unlike the polynomial Helmholtz–Duffing oscillator (JSV 2013), the rational form $\frac{\lambda x + \delta x^3}{1 + \nu x^2}$ introduces fundamentally different Fourier coefficient computations. The method’s success here demonstrates its versatility.
2. **Simultaneous homoclinic and heteroclinic prediction:** The same framework handles both types of global bifurcations without modification—a capability not available with classical perturbation methods.
3. **Connection to the MGHFP series:** This paper bridges the original GHFP method (2013) and the modern MGHFP series (2024–2026), showing the evolution from polynomial to rational restoring forces.

6 Limitations

- First-order perturbation limits accuracy when $\varepsilon > 2$ for some parameter regions.
- The method requires analytical knowledge of the saddle points, which may become algebraically intractable for more complex restoring forces.
- Only single-degree-of-freedom systems are considered.

Author: Zhenbo Li (lizhenbo@usc.edu.cn)
University of South China, School of Mathematics and Physics
Hunan Key Laboratory of Mathematical Modeling and Scientific Computing