




















Two-gel centroid distance: $\chi_I M = 0.1777778$ switched from locked $\chi_I M = 0.15$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.1788889$ switched from locked $\chi_I M = 0.15$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.18$ independent fixed $D = \text{run from } t = 0$ Two-gel centroid distance: $\chi_I M = 0.181$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.1825$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.185$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.1875$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.19$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.195$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.2$ independent fixed $D = \text{run from } t = 0$ Two-gel centroid distance: $\chi_I M = 0.205$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.215$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.22$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.23$ independent fixed $D = \text{run from } t = 0$ Two-gel centroid distance: $\chi_I M = 0.24$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.25$ independent fixed $D = \text{run from } t = 0$ Two-gel centroid distance: $\chi_I M = 0.26$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.28$ independent fixed $D = \text{run from } t = 0$ Two-gel centroid distance: $\chi_I M = 0.29$ switched from $\chi_I M = 0.18$ at $t = 20000$ TUTwo-gel centroid distance: $\chi_I M = 0.3$ independent fixed $D = \text{run from } t = 0$ 