
(a) Component ablation
PPK-FIP (full)
simplified F only (no averaging)
full F only (no averaging)
no fingerprint prior (joint, BMA)
no prior, BIC selection
two-step, simplified (ORIG)
two-step, full (ORIG-full)
fixed λ=1, s_p=0.05 (no meta-selection)
fixed λ=1, s_p=0.2 (no meta-selection)
fixed λ=3, s_p=0.05 (no meta-selection)
fixed λ=3, s_p=0.2 (no meta-selection)
no pseudo-processivity
no deactivation
no processivity, no deactivation
2.0
2.2
2.4
2.6
pooled LOTO RMSE (molar %)
(b) RMSE ratio to PPK-FIP per case
PPK-FIP (full) 1.00 1.00 1.00 1.00 1.00 1.00
simplified F only (no averaging) 0.99 0.91 1.00 1.32 1.03 1.11
full F only (no averaging) 0.96 1.00 1.00 0.97 0.98 1.00
no fingerprint prior (joint, BMA) 1.24 1.11 1.00 0.96 0.98 1.00
no prior, BIC selection 1.53 1.16 1.00 0.95 0.96 1.00
two-step, simplified (ORIG) 0.97 1.07 1.73 1.55 1.10 1.16
two-step, full (ORIG-full) 0.88 0.94 1.73 1.06 1.06 1.07
fixed λ=1, s_p=0.05 (no meta-selection) 0.96 1.04 1.13 1.03 1.07 1.03
fixed λ=1, s_p=0.2 (no meta-selection) 0.96 1.07 1.12 1.03 1.07 1.03
fixed λ=3, s_p=0.05 (no meta-selection) 1.00 1.00 1.00 1.00 1.00 1.00
fixed λ=3, s_p=0.2 (no meta-selection) 1.00 1.00 1.00 1.00 1.00 1.00
no pseudo-processivity 0.99 1.11 1.63 1.11 1.08 1.00
no deactivation 0.98 1.00 1.00 0.97 1.00 1.00
no processivity, no deactivation 0.96 1.11 1.63 1.06 1.08 1.00
Åkerman 70 °C
Åkerman 100 °C
Bornadel
Tao a_w 0.25
Tao a_w 0.35+0.45
Mao batches 1-3
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
log ratio (red = worse)