

# Incorporating stochastic sensitivity as an uncertainty measure for data assimilation and parcel trajectories in geophysical flows

Liam Blake

September 27, 2022

*Thesis submitted for the degree of  
Master of Philosophy  
in  
Applied Mathematics  
at The University of Adelaide  
Faculty of Sciences, Engineering and Technology  
School of Mathematical Sciences*



THE UNIVERSITY  
of ADELAIDE



# Contents



# List of Tables



# List of Figures





# Signed Statement

I certify that this work contains no material which has been accepted for the award of any other degree or diploma in my name in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text. In addition, I certify that no part of this work will, in the future, be used in a submission in my name for any other degree or diploma in any university or other tertiary institution without the prior approval of the University of Adelaide and where applicable, any partner institution responsible for the joint award of this degree.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library Search and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

Signed: ..... Date: .....



# Acknowledgements



# Dedication



# Abstract

Chapter ?? introduces





# Chapter 1

## Introduction



# Chapter 2

## Background

### 2.1 Notation

For a

### 2.2 Lagrangian Dynamics

We are typically interested in a spatial domain  $\Omega \subseteq \mathbb{R}^n$  and over some finite time interval  $[0, T]$  for finite  $T$ . Lagrangian trajectories are solutions of the first-order differential equation

$$\frac{dx_t}{dt} = u(x_t, t), \quad x_0 = x \in \Omega, \quad (2.1)$$

where  $u : \Omega \times [0, T] \rightarrow \mathbb{R}^n$  describes the velocity at a point in space and time.

#### 2.2.1 Flow map

In the mathematical treatment of Lagrangian dynamics, and in particular Lagrangian coherent structures (?), trajectories solving (??) are summarised by the flow map. The flow map is an operator mapping

The flow map can be defined more formally as follows.

**Definition 2.2.1 (Flow map)**

Suppose  $t_1, t_2 \in [0, T]$ . The **flow map**  $F_{t_1}^{t_2} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  from time  $t_1$  to  $t_2$  associated with (??) is the solution to

$$\frac{\partial F_{t_1}^\tau(x)}{\partial \tau} = u(F_{t_1}^\tau(x), \tau), \quad F_{t_1}^{t_1}(x) = x,$$

solved up to time  $\tau = t_2$ . Equivalently,

$$F_{t_1}^{t_2}(x) = x + \int_{t_1}^{t_2} u(F_{t_1}^\tau(x), \tau) d\tau.$$

The flow map satisfies the following properties

The gradient of the flow map satisfies a useful property; the equation of variations.

**Theorem 2.2.1**

Let  $F_{t_0}^t$  be the flow map corresponding to (??). Then, the spatial gradient  $\nabla F_{t_0}^t(x)$  satisfies the equation of variations

$$\frac{\partial \nabla F_{t_0}^t(x)}{\partial t} = \nabla u(F_{t_0}^t(x), t) \nabla F_{t_0}^t(x). \quad (2.2)$$

**Proof.**

□

## 2.2.2 Lagrangian coherent structures

TODO: Briefly discuss LCSs. Do not need to go into too much detail here. Mention whatever is relevant

## 2.3 Stochastic Differential Equations

$$dy_t = u(y_t, t) dt + \sigma(y_t, t) dW_t. \quad (2.3)$$

A solution to the stochastic differential equation can be defined rigorously ?.

**Definition 2.3.1**

A stochastic process  $\{y_t\}_{t \in [0, T]}$  taking values in  $R^n$  is said to be a **strong solution** of (??) with initial condition  $y_0 = \xi$  if the following holds:

1. For each  $t$ ,

2.

$$\int_0^T \left( \|u(y_t, t)\| + \|\sigma(y_t, t)\|^2 \right) dt < \infty \text{ a.s.}$$

3. For each  $t \in [0, T]$ ,

## 2.4 Stochastic Sensitivity

In most practical situations, the Eulerian velocity data driving ocean and atmospheric models relies upon measurements of estimates obtained on a low resolution spatial discretisation. There are limited tools within the LCS context that explicitly characterise the impact of these uncertainties. As such, there is recent interest in addressing this deficiency (??) . ? introduces stochastic sensitivity as a new tool for directly quantifying the impact of Eulerian uncertainty on Lagrangian trajectories. The evolution of Lagrangian trajectories is modelled as solution to a Itô stochastic ordinary differential equation.

TODO: Deterministic

The SDE model is

$$dy_t = u(y_t, t) dt + \varepsilon \sigma(y_t, t) dW_t, \quad (2.4)$$

where  $0 < \varepsilon \ll 1$  is a parameter quantifying the scale of the noise,  $\sigma : \mathbb{R}^2 \times [0, T] \rightarrow \mathbb{R}^{2 \times 2}$  is the  $2 \times 2$  diffusion matrix, and  $W_t$  is the canonical two-dimensional Wiener process. Since  $\sigma$  can vary by both space and time, the noise is multiplicative.

To quantify uncertainty in a way that is independent of the noise scale  $\varepsilon$ , ? defined the random variable  $z_\varepsilon(x, t)$  on  $\mathbb{R}^2 \times [0, T]$  as

$$z_\varepsilon(x, t) := \frac{y_t - F_0^t(x)}{\varepsilon}.$$

The main aim is to compute statistics of  $z_\varepsilon$  at the final time  $T$ , so that of  $z_\varepsilon(x, T)$ . ? then considers the signed projection of  $z_\varepsilon(x, T)$  onto a ray emanating from the deterministic position  $F_0^T(x)$  in a given direction, defining

$$P_\varepsilon(x, \theta) := \hat{n}^\top z_\varepsilon(x, T),$$

where  $\theta \in [-\pi/2, \pi/2)$  and

$$\hat{n}(\theta) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}.$$

The statistics of  $z_\varepsilon(x, T)$  and  $P_\varepsilon(x, \theta)$  are considered in the limit as  $\varepsilon \downarrow 0$ , which

TODO: something

The first result established by ? is that the expected location is deterministic, in the following sense.

### Theorem 2.4.1 (?)

For all  $x \in \mathbb{R}^2$ ,

$$\lim_{\varepsilon \downarrow 0} \mathbb{E}[z_\varepsilon(x, T)] = 0.$$

probably need  
some non-Sanjeeva  
citations here

The variance of  $P_\varepsilon(x, \theta)$  is used to assign a computable scalar measure of uncertainty to the trajectory.

**Definition 2.4.1 (?)**

a) The **anisotropic uncertainty** is a scalar field  $A : \mathbb{R}^2 \times [-\pi/2, \pi/2) \rightarrow [0, \infty)$  defined by

$$A(x, \theta) := \sqrt{\lim_{\varepsilon \downarrow 0} \mathbb{V} [P_\varepsilon(x, \theta)]}.$$

b) The **stochastic sensitivity** is a scalar field  $S : \mathbb{R}^2 \rightarrow [0, \infty)$  defined by

$$S^2(x) := \lim_{\varepsilon \downarrow 0} \sup_{\theta} \mathbb{V} [P_\varepsilon(x, \theta)].$$

**Theorem 2.4.2 (?)**

For  $x \in \mathbb{R}^2$  and  $\theta \in [-\pi/2, \pi/2)$ ,

$$A(x, \theta) = \left( \int_0^T \left\| \Lambda(F_0^t(x), t) J \hat{n}(\theta) \right\| dt \right)^{1/2}$$

# Chapter 3

## Extending stochastic sensitivity

There are several shortcomings of ? that warrant further extension, namely:

1. The tools are restricted to two-dimensional models, and the constructions using projections have no obvious extension to  $n$ -dimensions. Extending stochastic sensitivity to  $n$ -dimensions will enable application to a much broader class of models beyond the fluid flow context, including high-dimensional climate and ??? models.
2. ? only computes the expectation and variance of the projections  $P_\varepsilon(x, \theta)$ , which does not give us the distribution under the limit as  $\varepsilon$  approaches 0.
- 3.





# Bibliography

- Balasuriya, S. (2020*a*), ‘Stochastic approaches to Lagrangian coherent structures’, *The Role of Metrics in the Theory of Partial Differential Equations* **85**, 95–105.
- Balasuriya, S. (2020*b*), ‘Stochastic Sensitivity: A Computable Lagrangian Uncertainty Measure for Unsteady Flows’, *SIAM Review* **62**, 781–816.
- Balasuriya, S. and Gottwald, G. A. (2018), ‘Estimating stable and unstable sets and their role as transport barriers in stochastic flows’, *Physical Review E* **98**(1), 013106.
- Balasuriya, S., Ouellette, N. T. and Rypina, I. I. (2018), ‘Generalized Lagrangian coherent structures’, *Physica D: Nonlinear Phenomena* **372**, 31–51.
- Kallianpur, G. and Sundar, P. (2014), *Stochastic Analysis and Diffusion Processes*, number 24 in ‘Oxford Graduate Texts in Mathematics’, first edition edn, Oxford University Press, Oxford, United Kingdom.