

# Computable characterisations of uncertainty in geophysical models

Liam Blake

March 2, 2023

*Thesis submitted for the degree of  
Master of Philosophy  
in  
Applied Mathematics  
at The University of Adelaide  
Faculty of Sciences, Engineering and Technology  
School of Computer and Mathematical Sciences*



THE UNIVERSITY  
*of* ADELAIDE



# Contents

<b>Signed Statement</b>	<b>ix</b>
<b>Acknowledgements</b>	<b>xi</b>
<b>Dedication</b>	<b>xiii</b>
<b>Abstract</b>	<b>xv</b>
<b>1 Introduction</b>	<b>1</b>
<b>2 Background</b>	<b>3</b>
2.1 Notation . . . . .	3
2.2 Results from dynamical systems . . . . .	3
2.3 Stochastic differential equations . . . . .	4
2.3.1 Analytical tools for Itô calculus . . . . .	5
2.4 Aspects of stochastic parameterisation . . . . .	6
2.4.1 Additive versus multiplicative noise . . . . .	6
2.5 Stochastic sensitivity . . . . .	6
2.5.1 Current applications & shortcomings . . . . .	8
<b>3 Extending stochastic sensitivity</b>	<b>11</b>
<b>4 The Gaussian Limit</b>	<b>13</b>
<b>Bibliography</b>	<b>15</b>



# List of Tables



# List of Figures





# Signed Statement

I certify that this work contains no material which has been accepted for the award of any other degree or diploma in my name in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text. In addition, I certify that no part of this work will, in the future, be used in a submission in my name for any other degree or diploma in any university or other tertiary institution without the prior approval of the University of Adelaide and where applicable, any partner institution responsible for the joint award of this degree.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library Search and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

Signed: ..... Date: .....



# Acknowledgements



# Dedication



# Abstract

Chapter ?? introduces





# Chapter 1

## Introduction



# Chapter 2

## Background

### 2.1 Notation

The norm symbol  $\|\cdot\|$  without any additional qualifiers denotes the standard Euclidean norm for a vector, and the spectral (operator) norm induced by the Euclidean norm, i.e. for an  $n \times n$  matrix  $A$

$$\|A\| = \sup \left\{ \frac{\|Av\|}{\|v\|} \mid v \in \mathbb{R}^n, \|v\| \neq 0 \right\}.$$

For example, we say that a random variable  $A$  is equal to another random variable  $B$  almost surely (a.s.) if  $P(A = B) = 1$ .

### 2.2 Results from dynamical systems

$$\frac{dw_t}{dt} = u(w_t, t), \quad w_0 = x \in \Omega, \quad (2.1)$$

where  $u : \Omega \times [0, T] \rightarrow \mathbb{R}^n$  describes the velocity at a point in space and time.

In the mathematical treatment of Lagrangian dynamics, and in particular Lagrangian coherent structures [5], trajectories solving (2.1) are summarised by the flow map. The flow map is an operator mapping

The flow map can be defined formally as follows.

**Definition 2.2.1 (Flow map)** Suppose  $t_1, t_2 \in [0, T]$ . The **flow map**  $F_{t_1}^{t_2} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  from time  $t_1$  to  $t_2$  associated with (2.1) is the unique solution to

$$\frac{\partial F_{t_1}^\tau(x)}{\partial \tau} = u(F_{t_1}^\tau(x), \tau), \quad F_{t_1}^{t_1}(x) = x, \quad (2.2)$$

solved up to time  $\tau = t_2$ . Equivalently,

$$F_{t_1}^{t_2}(x) = x + \int_{t_1}^{t_2} u(F_{t_1}^\tau(x), \tau) d\tau.$$

The flow map satisfies the following properties

The gradient of the flow map satisfies a useful property; the equation of variations.

**Theorem 2.2.1** *Let  $F_{t_1}^t$  be the flow map corresponding to (2.1). Then, the spatial gradient  $\nabla F_{t_1}^t(x)$  satisfies the equation of variations*

$$\frac{\partial \nabla F_{t_1}^t(x)}{\partial t} = \nabla u(F_{t_1}^t(x), t) \nabla F_{t_1}^t(x). \quad (2.3)$$

**Proof.** Taking the gradient on both sides of (2.2) and using the chain rule gives

$$\nabla \left( \frac{\partial F_{t_1}^{t_2}(x)}{\partial t} \right) = \nabla u(F_{t_1}^t(x), t) \nabla F_{t_1}^t(x).$$

SMOOTHNESS □

An important inequality

**Theorem 2.2.2 (Grönwall's inequality)** *Let  $\alpha, \beta, u : [a, b] \rightarrow \mathbb{R}$  be functions such that  $\beta$  and  $u$  are continuous and that the negative part of  $\alpha$  is integrable on every closed and bounded subset of  $[a, b]$ . Then, if  $\beta$  is non-negative and for all  $t \in [a, b]$ ,*

$$u(t) \leq \alpha(t) + \int_a^t \beta(\tau) u(\tau) d\tau$$

then

$$u(t) \leq \alpha(t) + \int_a^t \alpha(\tau) \beta(\tau) \exp \left( \int_\tau^t \beta(s) ds \right) d\tau.$$

Additionally, if  $\alpha$  is non-decreasing, then

$$u(t) \leq \alpha(t) \exp \left( \int_a^t \beta(\tau) d\tau \right)$$

## 2.3 Stochastic differential equations

$$dy_t = u(y_t, t) dt + \sigma(y_t, t) dW_t. \quad (2.4)$$

A solution to the stochastic differential equation can be defined rigorously [10].

**Definition 2.3.1** A stochastic process  $\{y_t\}_{t \in [0, T]}$  taking values in  $\mathbb{R}^n$  is said to be a **strong solution** of (2.4) with initial condition  $y_0 = \xi$  if the following holds:

1. For each  $t$ ,

2.

$$\int_0^T \left( \|u(y_t, t)\| + \|\sigma(y_t, t)\|^2 \right) dt < \infty \text{ a.s.}$$

3. For each  $t \in [0, T]$ ,

$$y_t = \xi + \int_0^t u(y_\tau, \tau) d\tau + \int_0^t \sigma(y_\tau, \tau) dW_\tau \quad \text{a.s.}$$

**Theorem 2.3.1** Then, (2.4) has a unique strong solution.

**Proof.** See Theorem 6.2.1 of [10], for instance. □

### 2.3.1 Analytical tools for Itô calculus

There are several tools available for the analytic treatment of Itô integrals and solutions to stochastic differential equations, which we make use of throughout. The first is Itô's Lemma (or the Itô Formula), which is a change-of-variables formula in stochastic calculus and can be thought of as a generalisation of the chain rule from deterministic calculus. We state and use the multidimensional form of the Lemma for solutions to Itô stochastic differential equations, although more general forms exist (e.g. see Theorem 5.4.1 of [7]).

**Theorem 2.3.2 (Itô's Lemma)** Let  $X_t$  be the strong solution to the stochastic differential equation

$$dX_t = a(X_t, t) dt + b(X_t, t) dW_t,$$

where  $a : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^n$ ,  $b : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^{n \times p}$  and  $W_t$  is the canonical  $p$ -dimensional Wiener process. If  $f : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^m$  is twice continuously-differentiable, then the stochastic process  $Y_t := f(X_t, t)$  is a strong solution to the stochastic differential equation

$$dY_t = \left( \frac{\partial f}{\partial t}(X_t, t) + \nabla f(X_t, t) a(X_t, t) + \frac{1}{2} \text{tr} \left[ b(X_t, t)^\top \nabla \nabla f(X_t, t) b(X_t, t) \right] \right) dt + \nabla f(X_t, t) b(X_t, t) dW_t.$$

**Proof.** □

**Theorem 2.3.3 (Burkholder-Davis-Gundy Inequality)** *Let  $M_t$  be an Itô-integrable stochastic process taking values in  $\mathbb{R}^n$ . Then, for any  $p > 0$  there exists constants  $c_p, C_p > 0$  independent of the stochastic process  $M_t$  such that*

$$c_p \mathbb{E} \left[ \left( \int_0^t \|M_\tau\|^2 d\tau \right)^p \right] \leq \mathbb{E} \left[ \sup_{\tau \in [0, t]} \left\| \int_0^\tau M_s dW_s \right\|^{2p} \right] \leq C_p \mathbb{E} \left[ \left( \int_0^t \|M_\tau\|^2 d\tau \right)^p \right].$$

**Proof.**

□

## 2.4 Aspects of stochastic parameterisation

When using deterministic systems to model real-world phenomena, there are many ways in which uncertainty can arise. For instance,

Stochastic parameterisation is a These unresolved subgrid effects are accounted for by introducing stochastic noise into the otherwise deterministic model.

Berner et al. [6]

Leutbecher et al. [11]

### 2.4.1 Additive versus multiplicative noise

When  $\sigma = \sigma(t)$  depends only on  $t$ , then noise is considered *additive*. If there is spatial dependence in  $\sigma$ , i.e.  $\sigma = \sigma(x, t)$ , then the noise considered *multiplicative*.

For instance, Sura et al. [12] shows that the non-Gaussian statistics observed in atmospheric regimes can arise from linear models with multiplicative noise.

## 2.5 Stochastic sensitivity

In most practical situations, the Eulerian velocity data driving ocean and atmospheric models relies upon measurements of estimates obtained on a low resolution spatial discretisation.

There are limited tools within the LCS context that explicitly characterise the impact of these uncertainties As such, there is recent interest in addressing this deficiency [? 2]. Balasuriya [3] introduces stochastic sensitivity as a new tool for directly quantifying the impact of Eulerian uncertainty on Lagrangian trajectories. The evolution of Lagrangian trajectories is modelled as solution to a Itô stochastic ordinary differential equation.

TODO: Deterministic

The SDE model is

$$dy_t = u(y_t, t) dt + \epsilon \sigma(y_t, t) dW_t, \quad (2.5)$$

where  $0 < \epsilon \ll 1$  is a parameter quantifying the scale of the noise,  $\sigma : \mathbb{R}^2 \times [0, T] \rightarrow \mathbb{R}^{2 \times 2}$  is the  $2 \times 2$  diffusion matrix, and  $W_t$  is the canonical two-dimensional Wiener process. Since  $\sigma$  can vary by both space and time, the noise is multiplicative.

To quantify uncertainty in a way that is independent of the noise scale  $\epsilon$ , [3] defined the random variable  $z_\epsilon(x, t)$  on  $\mathbb{R}^2 \times [0, T]$  as

$$z_\epsilon(x, t) := \frac{y_t - F_0^t(x)}{\epsilon}.$$

The main aim is to compute statistics of  $z_\epsilon$  at the final time  $T$ , so that of  $z_\epsilon(x, T)$ . Balasuriya [3] then considers the signed projection of  $z_\epsilon(x, T)$  onto a ray emanating from the deterministic position  $F_0^T(x)$  in a given direction, defining

$$P_\epsilon(x, \theta) := \hat{n}^\top z_\epsilon(x, T),$$

where  $\theta \in [-\pi/2, \pi/2)$  and

$$\hat{n}(\theta) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}.$$

The statistics of  $z_\epsilon(x, T)$  and  $P_\epsilon(x, \theta)$  are considered in the limit as  $\epsilon \downarrow 0$ , which

TODO: something

The first result established by [3] is that the expected location is deterministic, in the following sense.

**Theorem 2.5.1** ([3]) *For all  $x \in \mathbb{R}^2$ ,*

$$\lim_{\epsilon \downarrow 0} \mathbb{E}[z_\epsilon(x, T)] = 0.$$

The variance of  $P_\epsilon(x, \theta)$  is used to assign a computable scalar measure of uncertainty to the trajectory.

**Definition 2.5.1** ([3]) *a) The **anisotropic uncertainty** is a scalar field  $A : \mathbb{R}^2 \times [-\pi/2, \pi/2) \rightarrow [0, \infty)$  defined by*

$$A(x, \theta) := \sqrt{\lim_{\epsilon \downarrow 0} \mathbb{V}[P_\epsilon(x, \theta)]}.$$

*b) The **stochastic sensitivity** is a scalar field  $S : \mathbb{R}^2 \rightarrow [0, \infty)$  defined by*

$$S^2(x) := \lim_{\epsilon \downarrow 0} \sup_{\theta} \mathbb{V}[P_\epsilon(x, \theta)].$$

By employing techniques from both deterministic and stochastic calculus (i.e. Grönwall's inequality, the Burkholder-Davis-Gundy inequality, Itô's Lemma), Balasuriya further established expressions for both the anisotropic uncertainty and the stochastic sensitivity that are computable given only the flow map and velocity data.

**Theorem 2.5.2 ([3])** *For  $x \in \mathbb{R}^2$ , set  $w := F_0^t(x)$ . Then, for any  $\theta \in [-\pi/2, \pi/2)$ ,*

$$A(x, \theta) = \left( \int_0^T \|\Lambda(x, t, T) J \hat{n}(\theta)\| dt \right)^{1/2},$$

where

$$\Lambda(x, t, T) := e^{\int_t^T [\nabla \cdot u](F_0^\xi(x), \xi) d\xi} \sigma(F_0^t(x), t)^\top J \nabla_w F_T^t(w),$$

with the gradients  $\nabla_w$  of the flow map taken with respect to the mapped position  $w$ , and

$$J := \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Additionally, stochastic sensitivity is computed as

$$S^2(x) = P(x) + N(x),$$

with

$$\begin{aligned} L(x) &:= \frac{1}{2} \sum_{i=1}^2 \int_0^T \left[ \Lambda_{i2}(x, t, T)^2 - \Lambda_{i1}(x, t, T)^2 \right] dt \\ M(x) &:= \sum_{i=1}^2 \int_0^T \Lambda_{i1}(x, t, T) \Lambda_{i2}(x, t, T) dt \\ N(x) &:= \sqrt{L^2(x) + M^2(x)} \\ P(x) &:= \left| \frac{1}{2} \sum_{i=1}^2 \sum_{j=1}^2 \int_0^T \Lambda_{ij}(x, t, T)^2 dt \right|, \end{aligned}$$

where  $\Lambda_{ij}$  is the  $(i, j)$ -element of  $\Lambda$ .

### 2.5.1 Current applications & shortcomings

Since stochastic sensitivity is only a recent development, it has only been applied in a limited number of places so far. Here, we briefly review the literature in which the original formulation stochastic sensitivity by [3] has been applied.



- Balasuriya [4] uses stochastic sensitivity to compute an error bound for the finite-time Lyapunov computation.
- Fang et al. [8]
- Badza et al. [1] investigate the impact of velocity uncertainty on Lagrangian coherent structures (e.g. see the reviews by Balasuriya et al. [5] and Hadjighasem et al. [9]) extracted as robust sets with stochastic sensitivity.

Although

1. The tools are restricted to two-dimensional models, and the constructions using projections have no obvious extension to  $n$ -dimensions. Extending stochastic sensitivity to  $n$ -dimensions will enable application to a much broader class of models beyond the fluid flow context, including high-dimensional climate and ??? models.
2. Balasuriya [3] only computes the expectation and variance of the projections  $P_\epsilon(x, \theta)$ , which does not give us the distribution under the limit as  $\epsilon$  approaches 0.
3. The computational formula for the anisotropic uncertainty and stochastic sensitivity require knowledge of the divergence  $\nabla \cdot u$  of the velocity field.



## Chapter 3

# Extending stochastic sensitivity

There are several shortcomings of [3] that warrant further extension, namely:



## Chapter 4

# The Gaussian Limit



# Bibliography

- [1] A. Badza, T. W. Mattner, and S. Balasuriya. How sensitive are Lagrangian coherent structures to uncertainties in data? *Physica D: Nonlinear Phenomena*, 444:133580, Feb. 2023. ISSN 0167-2789. doi: 10.1016/j.physd.2022.133580.
- [2] S. Balasuriya. Stochastic approaches to Lagrangian coherent structures. *The Role of Metrics in the Theory of Partial Differential Equations*, 85:95–105, Jan. 2020. doi: 10.2969/aspm/08510095.
- [3] S. Balasuriya. Stochastic Sensitivity: A Computable Lagrangian Uncertainty Measure for Unsteady Flows. *SIAM Review*, 62:781–816, Nov. 2020. doi: 10.1137/18M1222922.
- [4] S. Balasuriya. Uncertainty in finite-time Lyapunov exponent computations. *Journal of Computational Dynamics*, 7(2):313–337, 2020. doi: 10.3934/jcd.2020013.
- [5] S. Balasuriya, N. T. Ouellette, and I. I. Rypina. Generalized Lagrangian coherent structures. *Physica D: Nonlinear Phenomena*, 372:31–51, June 2018. ISSN 0167-2789. doi: 10.1016/j.physd.2018.01.011.
- [6] J. Berner, U. Achatz, L. Batté, L. Bengtsson, A. de la Cámara, H. M. Christensen, M. Colangeli, D. R. B. Coleman, D. Crommelin, S. I. Dolaptchiev, C. L. E. Franzke, P. Friederichs, P. Imkeller, H. Järvinen, S. Juricke, V. Kitsios, F. Lott, V. Lucarini, S. Mahajan, T. N. Palmer, C. Penland, M. Sakradzija, J.-S. von Storch, A. Weisheimer, M. Weniger, P. D. Williams, and J.-I. Yano. Stochastic Parameterization: Toward a New View of Weather and Climate Models. *Bulletin of the American Meteorological Society*, 98(3):565–588, Mar. 2017. ISSN 0003-0007, 1520-0477. doi: 10.1175/BAMS-D-15-00268.1.
- [7] P. Brémaud. *Probability Theory and Stochastic Processes*. Universitext. Springer, first edition, 2020. ISBN 978-3-030-40182-5.
- [8] L. Fang, S. Balasuriya, and N. T. Ouellette. Disentangling resolution, precision, and inherent stochasticity in nonlinear systems. *Physical Review Research*, 2(2):023343, June 2020. doi: 10.1103/PhysRevResearch.2.023343.

- [9] A. Hadjighasem, M. Farazmand, D. Blazeovski, G. Froyland, and G. Haller. A critical comparison of Lagrangian methods for coherent structure detection. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27(5):053104, May 2017. ISSN 1054-1500. doi: 10.1063/1.4982720.
- [10] G. Kallianpur and P. Sundar. *Stochastic Analysis and Diffusion Processes*. Number 24 in Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, United Kingdom, first edition edition, 2014. ISBN 978-0-19-965706-3 978-0-19-965707-0.
- [11] M. Leutbecher, S.-J. Lock, P. Ollinaho, S. T. K. Lang, G. Balsamo, P. Bechtold, M. Bonavita, H. M. Christensen, M. Diamantakis, E. Dutra, S. English, M. Fisher, R. M. Forbes, J. Goddard, T. Haiden, R. J. Hogan, S. Juricke, H. Lawrence, D. MacLeod, L. Magnusson, S. Malardel, S. Massart, I. Sandu, P. K. Smolarkiewicz, A. Subramanian, F. Vitart, N. Wedi, and A. Weisheimer. Stochastic representations of model uncertainties at ECMWF: State of the art and future vision. *Quarterly Journal of the Royal Meteorological Society*, 143(707):2315–2339, 2017. ISSN 1477-870X. doi: 10.1002/qj.3094.
- [12] P. Sura, M. Newman, C. Penland, and P. Sardeshmukh. Multiplicative Noise and Non-Gaussianity: A Paradigm for Atmospheric Regimes? *Journal of the Atmospheric Sciences*, 62(5):1391–1409, May 2005. ISSN 0022-4928, 1520-0469. doi: 10.1175/JAS3408.1.