

233B - Final

5.6.12 Let X be a prevariety over an algebraically closed field k , and let $P \in X$ be a (closed) point of X . Let $D = \text{Spec } k[x]/(x^2)$ be the “double point”. Show that the tangent space $T_{X,P}$ to X at P can be canonically identified with the set of morphisms $D \rightarrow X$ that map the unique point of D to P .

Proof. □

5.6.13 Let X be an affine variety, let Y be a closed subscheme of X defined by the ideal $I \subset A(X)$, and let \tilde{X} be the blow-up of X at I . Show that:

- (i) $\tilde{X} = \text{Proj}(\bigoplus_{d \geq 0} I^{(d)})$, where $I^{(0)} := A(X)$.
- (ii) The projection map $\tilde{X} \rightarrow X$ is the morphism induced by the ring homomorphism $I^{(0)} \rightarrow \bigoplus_{d \geq 0} I^{(d)}$.
- (iii) The exceptional divisor of the blow-up, i.e. the fiber $Y \times_X \tilde{X}$ of the blow-up $\tilde{X} \rightarrow X$ over Y , is isomorphic to $\text{Proj}(\bigoplus_{d \geq 0} I^{(d)}/I^{(d+1)})$.

Proof. □

6.7.3 Let $X \subset \mathbb{P}^n$ scheme with Hilbert polynomial χ . Define the arithmetic genus of X to be $g(X) = (-1)^{\dim X} \cdot (\chi(0) - 1)$.

- (i) Show that $g(\mathbb{P}^n) = 0$.
- (ii) If X is a hypersurface of degree d in \mathbb{P}^n , show that $g(X) = \binom{d-1}{n}$. In particular, if $C \subset \mathbb{P}^2$ is a plane curve of degree d , then $g(C) = \frac{1}{2}(d-1)(d-1)$.
- (iii) Compute the arithmetic genus of the union of the three coordinate axes

$$Z(x_1x_2, x_1x_3, x_2x_3) \subset \mathbb{P}^3.$$

6.7.8 Let $C_1 = \{f_1 = 0\}$ and $C_2 = \{f_2 = 0\}$ be affine curves in \mathbb{A}_k^2 , and let $P \in C_1 \cap C_2$ be a point. Show that the intersection multiplicity of C_1 and C_2 at P (i.e. the length of the component at P of the intersection scheme $C_1 \cap C_2$) is equal to the dimension of the vector space $\mathcal{O}_{\mathbb{A}^2, P}/(f_1, f_2)$ over k .

7.8.8 What is the line bundle on $\mathbb{P}^n \times \mathbb{P}^m$ leading to the Segre embedding $\mathbb{P}^n \times \mathbb{P}^m \rightarrow \mathbb{P}^N$ by the correspondence of ... What is the line bundle leading to the degree- d Veronese embedding $\mathbb{P}^n \rightarrow \mathbb{P}^N$?

7.8.10 Let X be a smooth projective curve, and let $P \in X$ be a point. Show that there is a rational function on X that is regular everywhere except at P .