

# Finding and Counting Substructures in Graphs and Hypergraphs

Liam Hardiman

December 10, 2020

# 1 A Finding Problem

## Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .
- The **degree** of a vertex  $v$ , denoted  $d_G(v)$ , is the number of edges of  $G$  that  $v$  appears in.

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .
- The **degree** of a vertex  $v$ , denoted  $d_G(v)$ , is the number of edges of  $G$  that  $v$  appears in.
- picture here

# History and Motivation

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

# History and Motivation

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

## Proof sketch:

asdf





# History and Motivation

- What about odd subgraphs?

# History and Motivation

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

# History and Motivation

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

## Conjecture (A. Scott - 2001)

*There exists some constant  $c > 0$  such that every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq cn$  and  $G[W]$  has all degrees odd.*