

# Finding and Counting Substructures in Graphs and Hypergraphs

Liam Hardiman

December 10, 2020

- 1 A Finding Problem
- 2 A Counting Problem
- 3 Another Counting Problem

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .
- The **degree** of a vertex  $v$ , denoted  $d_G(v)$ , is the number of edges of  $G$  that  $v$  appears in.

# Quick Definitions

- A **graph**  $G = (V, E)$  consists of a (finite) set of **vertices**  $V$  and a set  $E$  of unordered pairs of vertices called **edges**.
- If  $X \subseteq V$ , then the **subgraph induced by**  $X$ ,  $G[X]$  is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .
- The **degree** of a vertex  $v$ , denoted  $d_G(v)$ , is the number of edges of  $G$  that  $v$  appears in.
- picture here

# History and Motivation

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

# History and Motivation

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

## Proof sketch:

asdf





# History and Motivation

- What about odd subgraphs?

# History and Motivation

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

# History and Motivation

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

## Conjecture (A. Scott - 2001)

*There exists some constant  $c > 0$  such that every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq cn$  and  $G[W]$  has all degrees odd.*

# History and Motivation

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

## Conjecture (A. Scott - 2001)

*There exists some constant  $c > 0$  such that every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq cn$  and  $G[W]$  has all degrees odd.*

# History and Motivation

## Definition

The **Erdős-Renyi random graph**,  $\mathcal{G}(n, p)$ , is the random variable that outputs a graph on  $n$  vertices, any two of which are independently connected with probability  $p$ .

# History and Motivation

## Definition

The **Erdős-Renyi random graph**,  $\mathcal{G}(n, p)$ , is the random variable that outputs a graph on  $n$  vertices, any two of which are independently connected with probability  $p$ .

## Theorem (A. Scott - 1992)

*Let  $G \sim \mathcal{G}(n, 1/2)$ . Then with high probability (that is, with probability  $1 - o(1)$ ),  $G$  has an induced subgraph on at least  $0.7729n$  vertices with all degrees odd.*

# Other Moduli and Remainders

## Conjecture (A. Scott - 2001)

*For any positive integer  $q$  at least 2, there exists a constant  $c_q$  so that every graph on  $n$  vertices without isolated vertices has an induced subgraph on at least  $c_q n$  vertices with all degrees  $1 \pmod q$ .*

# Other Moduli and Remainders

## Conjecture (A. Scott - 2001)

*For any positive integer  $q$  at least 2, there exists a constant  $c_q$  so that every graph on  $n$  vertices without isolated vertices has an induced subgraph on at least  $c_q n$  vertices with all degrees  $1 \pmod{q}$ .*

## Theorem (A. Ferber, H., M. Krivelevich - 2020+)

*Let  $q \geq 2$  and let  $r$  be an integer. Then there exists a constant  $c_q$  such that, with high probability, the random graph  $G \sim \mathcal{G}(n, 1/2)$  has an induced subgraph on at least  $c_q n$  vertices with all degrees  $r \pmod{q}$ .*



# Key Idea Behind Proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.

# Key Idea Behind Proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.
- In this case, the  $i$ -th entry of  $M\mathbf{1}$  is the degree of vertex  $i$  in  $G$ , where  $\mathbf{1}$  is the length  $n$  all-1 vector.

# Key Idea Behind Proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.
- In this case, the  $i$ -th entry of  $M\mathbf{1}$  is the degree of vertex  $i$  in  $G$ , where  $\mathbf{1}$  is the length  $n$  all-1 vector.

## Observation

*$G$  contains a subgraph with all degrees  $r \pmod{q}$  if and only if its adjacency matrix contains a principal submatrix  $B$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .*

# A Useful Computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables.

# A Useful Computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables.

$$\Pr[x_1 + \dots + x_t \equiv r] = \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)]$$

# A Useful Computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables.

$$\begin{aligned}\Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right]\end{aligned}$$

# A Useful Computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables.

$$\begin{aligned}\Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \prod_{j=1}^t \mathbb{E} e^{2\pi i \ell x_j / q}\end{aligned}$$

# A Useful Computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables.

$$\begin{aligned}\Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \prod_{j=1}^t \mathbb{E} e^{2\pi i \ell x_j / q} \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \left( \frac{1 + e^{2\pi i \ell / q}}{2} \right)^t.\end{aligned}$$



# A Useful Computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

# A Useful Computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

# A Useful Computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

- Calculus trick: when  $|x| \leq \pi/2$ ,  $|\sin x| \geq (2/\pi)|x|$ .

# A Useful Computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \dots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

- Calculus trick: when  $|x| \leq \pi/2$ ,  $|\sin x| \geq (2/\pi)|x|$ .
- From this, it follows that  $|\cos(\pi\ell/q)| \leq e^{-2/q^2}$  for all  $\ell = 1, \dots, q-1$ .

# A Useful Computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \dots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

- Calculus trick: when  $|x| \leq \pi/2$ ,  $|\sin x| \geq (2/\pi)|x|$ .
- From this, it follows that  $|\cos(\pi\ell/q)| \leq e^{-2/q^2}$  for all  $\ell = 1, \dots, q-1$ .

$$\left| \Pr[x_1 + \dots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{q-1}{q} e^{-2t/q^2}.$$

# A Useful Computation

- The distribution of a random Bernoulli sum modulo  $q$  is asymptotically uniform!

# A Useful Computation

- The distribution of a random Bernoulli sum modulo  $q$  is asymptotically uniform!
- By independence we have.

## Lemma

*Let  $M$  be an  $s \times t$  matrix whose entries are iid  $\text{Bern}(1/2)$  random variables. Then for any  $v \in \mathbb{Z}_q^s$ ,*

$$\Pr[M\mathbf{1} \equiv v] = \frac{1}{q^s} \left( 1 + O\left(e^{-2t/q^2}\right) \right)^s.$$

# Symmetric Case

- But the adjacency matrix is symmetric!



# Symmetric Case

- But the adjacency matrix is symmetric!
- A similar, but more technical argument handles the symmetric case.

# Symmetric Case

- But the adjacency matrix is symmetric!
- A similar, but more technical argument handles the symmetric case.

## Lemma

*Let  $M$  be an  $m \times m$  symmetric matrix whose diagonal is zero and whose entries above the diagonal are iid  $\text{Bern}(1/2)$  random variables. Then for any  $v \in \mathbb{Z}_q^m$ ,*

$$\Pr[M\mathbf{1} \equiv v] = \begin{cases} \frac{1}{q^m} (1 + O(e^{-m/q^2})) & \text{if } q \text{ is odd} \\ \frac{2}{q^m} (1 + O(e^{-m/q^2})) & \text{if } q \text{ is even and } \sum v_i \text{ is even} \\ 0 & \text{if } q \text{ is even and } \sum v_i \text{ is odd.} \end{cases}$$

# Parity?

- We have

$$\sum_i (M\mathbf{1})_i = \sum_{i,j} M_{ij} = 2 \sum_{i < j} M_{ij},$$

which is always even modulo  $q$  if  $q$  is even and can be any residue modulo odd  $q$ .

# Parity?

- We have

$$\sum_i (M\mathbf{1})_i = \sum_{i,j} M_{ij} = 2 \sum_{i < j} M_{ij},$$

which is always even modulo  $q$  if  $q$  is even and can be any residue modulo odd  $q$ .

- The symmetric lemma then says that the distribution of  $M\mathbf{1} \pmod{q}$  is asymptotically uniform over all “feasible” values.

# Proof (Sketch) of Theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

# Proof (Sketch) of Theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

- Let  $k = c_q n$  for some  $c_q > 0$  and let  $X_k$  be the number of  $k \times k$  principal submatrices  $B$  of the adjacency matrix of  $G$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .

# Proof (Sketch) of Theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

- Let  $k = c_q n$  for some  $c_q > 0$  and let  $X_k$  be the number of  $k \times k$  principal submatrices  $B$  of the adjacency matrix of  $G$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .
- Show that  $c_q$  can be chosen so that  $\text{Var}[X_k] = o(E[X_k]^2)$ . Then  $G$  has an induced subgraph of size  $c_q n$  with high probability.

# Proof (Sketch) of Theorem

Maybe have some of the calculations and that overlap diagram.



State theorem for arbitrary distribution of residues mod  $q$ .

- 1 A Finding Problem
- 2 A Counting Problem
- 3 Another Counting Problem

# Packing

- Recall Gallai's result on even subgraphs.

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

# Packing

- Recall Gallai's result on even subgraphs.

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

- What else can we say about partitions and remainders?

# Packing Theorems

## Theorem (A. Scott - 2001)

*Let  $G$  be a graph. Then  $G$  admits a partition into odd subgraphs if and only if every component of  $G$  has even order.*

# Packing Theorems

## Theorem (A. Scott - 2001)

*Let  $G$  be a graph. Then  $G$  admits a partition into odd subgraphs if and only if every component of  $G$  has even order.*

## Theorem (A. Scott - 2001)

*With high probability,  $G \sim \mathcal{G}(n, p)$  for  $n$  even admits a partition into three odd subgraphs.*

# Packing Theorems

## Theorem (A. Scott - 2001)

*Let  $G$  be a graph. Then  $G$  admits a partition into odd subgraphs if and only if every component of  $G$  has even order.*

## Theorem (A. Scott - 2001)

*With high probability,  $G \sim \mathcal{G}(n, p)$  for  $n$  even admits a partition into three odd subgraphs.*

## Conjecture (A. Scott - 2001)

*For  $q > 1$  and  $r \leq q$ , there exists a constant  $c_q$  such that  $G \sim \mathcal{G}(n, p)$  admits a partition into  $c_q$  classes such that the degrees in each class are congruent to  $r \pmod{q}$  with high probability.*

# Packing Theorems

## Theorem (A. Ferber, H., M. Krivelevich)

*For  $q > 1$  and  $r \leq q$ ,  $G \sim \mathcal{G}(n, 1/2)$  admits a partition into  $q + 1$  classes such that the degrees in each class are congruent to  $r \pmod{q}$  with high probability.*



# Sketch of Packing Proof

Explain that it's just another second moment argument. Maybe some calculations, diagrams. Explain need for “both ways” lemma.

- 1 A Finding Problem
- 2 A Counting Problem
- 3 Another Counting Problem

# Definitions

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

# Definitions

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

## Definition

Let  $G = (V, E)$  be a graph and let  $\mathcal{C}$  be a sequence of edges. Then  $\mathcal{C}$  is a...

- ...**path** if it joins a sequence of distinct vertices.

# Definitions

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

## Definition

Let  $G = (V, E)$  be a graph and let  $\mathcal{C}$  be a sequence of edges. Then  $\mathcal{C}$  is a...

- ...**path** if it joins a sequence of distinct vertices.
- ...**cycle** if it joints a sequence of vertices  $v_1, v_2, \dots, v_k, v_1$ .

# Definitions

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

## Definition

Let  $G = (V, E)$  be a graph and let  $\mathcal{C}$  be a sequence of edges. Then  $\mathcal{C}$  is a...

- ...**path** if it joins a sequence of distinct vertices.
- ...**cycle** if it joints a sequence of vertices  $v_1, v_2, \dots, v_k, v_1$ .
- ...**Hamiltonian cycle** if it is a cycle that visits every vertex in  $G$  exactly once.

# History and Motivation

- When does a graph have Hamiltonian a cycle (when is it **Hamiltonian**)?

# History and Motivation

- When does a graph have Hamiltonian a cycle (when is it **Hamiltonian**)?
- For any graph  $G$ , let  $\delta(G)$  denote the minimum degree of any vertex.



# History and Motivation

- When does a graph have Hamiltonian a cycle (when is it **Hamiltonian**)?
- For any graph  $G$ , let  $\delta(G)$  denote the minimum degree of any vertex.

Theorem (G. Dirac - 1952)

*A graph on  $n \geq 3$  vertices is Hamiltonian if  $\delta(G) \geq n/2$ .*

# History and Motivation

- When does a graph have Hamiltonian a cycle (when is it **Hamiltonian**)?
- For any graph  $G$ , let  $\delta(G)$  denote the minimum degree of any vertex.

Theorem (G. Dirac - 1952)

*A graph on  $n \geq 3$  vertices is Hamiltonian if  $\delta(G) \geq n/2$ .*

- A graph is Hamiltonian if it is sufficiently dense.

# Counting Hamiltonian Cycles in Dense Graphs - Intuition

- The expected number of Hamiltonian cycles in  $G \sim \mathcal{G}(n, p)$  is

$$\frac{1}{2}(n-1)!p^n.$$

# Counting Hamiltonian Cycles in Dense Graphs - Intuition

- The expected number of Hamiltonian cycles in  $G \sim \mathcal{G}(n, p)$  is

$$\frac{1}{2}(n-1)!p^n.$$

- Fix  $\gamma > 0$ . If  $p > (1/2 + \gamma)$ , then

# Cuckler-Kahn

# Introduce hypergraphs and notion of cycle

# Hypergraph theorems

# Our theorem



# Proof sketch