

# Problems in Extremal Graph and Hypergraph Theory

# Liam Hardiman

December 10, 2020

- 1 Graph substructures
- 2 Counting perfect matchings and Hamiltonian cycles
- 3 Degrees of prescribed remainder

# Basic Definitions

- A **graph**  $G = (V, E)$  consists of a (finite for us) set of **vertices**  $V$  and a set  $E$  of edges where  $E \subseteq \binom{V}{2}$ .

# Basic Definitions

- A **graph**  $G = (V, E)$  consists of a (finite for us) set of **vertices**  $V$  and a set  $E$  of edges where  $E \subseteq \binom{V}{2}$ .
- The number of vertices incident to  $v \in V$  is its **degree**,  $d_G(v)$ . We let  $\Delta(G)$  and  $\delta(G)$  denote the maximum and minimum degree, respectively.

# Basic Definitions

- A **graph**  $G = (V, E)$  consists of a (finite for us) set of **vertices**  $V$  and a set  $E$  of edges where  $E \subseteq \binom{V}{2}$ .
- The number of vertices incident to  $v \in V$  is its **degree**,  $d_G(v)$ . We let  $\Delta(G)$  and  $\delta(G)$  denote the maximum and minimum degree, respectively.
- $H = (V', E')$  is a **subgraph** of  $G$  if  $V' \subseteq V$ ,  $E' \subseteq E$  and each edge in  $E'$  has both ends in  $V'$ .

# Basic Definitions

- A **graph**  $G = (V, E)$  consists of a (finite for us) set of **vertices**  $V$  and a set  $E$  of edges where  $E \subseteq \binom{V}{2}$ .
- The number of vertices incident to  $v \in V$  is its **degree**,  $d_G(v)$ . We let  $\Delta(G)$  and  $\delta(G)$  denote the maximum and minimum degree, respectively.
- $H = (V', E')$  is a **subgraph** of  $G$  if  $V' \subseteq V$ ,  $E' \subseteq E$  and each edge in  $E'$  has both ends in  $V'$ .
- If  $X \subseteq V$ , then the **subgraph induced by  $X$** ,  $G[X]$ , is the graph with vertex set  $X$  and all edges from  $E$  that have both ends in  $X$ .

# Example




Figure: A graph

# Example




Figure: A subgraph



# Example




Figure: An induced subgraph

# An existence question


## Question

*How many edges does a graph on  $n$  vertices need in order to guarantee that it contains a triangle?*

# An existence question

## Question



*How many edges does a graph on  $n$  vertices need in order to guarantee that it contains a triangle?*



# An existence question

## Question

*How many edges does a graph on  $n$  vertices need in order to guarantee that it contains a triangle?*



# Guaranteeing the existence of a subgraph

## Theorem (W. Mantel - 1907)

*If  $G$  is a graph on  $n$  vertices with more than  $\lfloor n/2 \rfloor \lceil n/2 \rceil$  edges, then  $G$  contains a triangle.*

# Guaranteeing the existence of a subgraph

## Theorem (W. Mantel - 1907)

*If  $G$  is a graph on  $n$  vertices with more than  $\lfloor n/2 \rfloor \lceil n/2 \rceil$  edges, then  $G$  contains a triangle.*

- The graph on  $n$  vertices with all possible edges is called the **complete graph** and is denoted  $K_n$ .

# Guaranteeing the existence of a subgraph

## Theorem (W. Mantel - 1907)

*If  $G$  is a graph on  $n$  vertices with more than  $\lfloor n/2 \rfloor \lceil n/2 \rceil$  edges, then  $G$  contains a triangle.*

- The graph on  $n$  vertices with all possible edges is called the **complete graph** and is denoted  $K_n$ .

## Theorem (P. Turán - 1941)

*If  $G$  is a graph on  $n$  vertices with more than  $(1 - \frac{1}{r})\frac{n^2}{2}$  edges, then  $G$  contains a copy of  $K_{r+1}$ .*

# What about other graphs?

- Given a graph  $H$ , the **extremal number** (or **Turán number**)  $\text{ex}(n, H)$  is the largest number of edges in an  $H$ -free graph on  $n$  vertices.



# What about other graphs?

- Given a graph  $H$ , the **extremal number** (or **Turán number**)  $ex(n, H)$  is the largest number of edges in an  $H$ -free graph on  $n$  vertices.

Theorem (P. Erdős, A. Stone - 1946)

Let  $H$  be a nonempty graph and let  $r$  be the **chromatic number** of  $H$  (the smallest number of colors needed to color the vertices of  $H$  so that no two adjacent vertices are of the same color). Then

$$ex(n, H) = \left(1 - \frac{1}{r-1}\right) \binom{n}{2} + o(n^2).$$

# What about other graphs?

- Apparently, the case of a **bipartite graph** (one whose chromatic number is 2) is exceptional. We only know the correct order of magnitude for such graphs in specific cases.

# What about other graphs?

- Apparently, the case of a **bipartite graph** (one whose chromatic number is 2) is exceptional. We only know the correct order of magnitude for such graphs in specific cases.
- If  $T$  is a **tree** (a graph which contains no cycles), then  $\text{ex}(n, T) = \Theta(n)$ .

# What about other graphs?

- Apparently, the case of a **bipartite graph** (one whose chromatic number is 2) is exceptional. We only know the correct order of magnitude for such graphs in specific cases.
- If  $T$  is a **tree** (a graph which contains no cycles), then  $\text{ex}(n, T) = \Theta(n)$ .
- Erdős showed that  $\text{ex}(n, C_{2k}) = O(n^{1+1/k})$  for even cycles  $C_{2k}$ . Matching lower bounds are only known for  $C_4$ ,  $C_6$  and  $C_{10}$ .

# What about other graphs?

- Apparently, the case of a **bipartite graph** (one whose chromatic number is 2) is exceptional. We only know the correct order of magnitude for such graphs in specific cases.
- If  $T$  is a **tree** (a graph which contains no cycles), then  $\text{ex}(n, T) = \Theta(n)$ .
- Erdős showed that  $\text{ex}(n, C_{2k}) = O(n^{1+1/k})$  for even cycles  $C_{2k}$ . Matching lower bounds are only known for  $C_4$ ,  $C_6$  and  $C_{10}$ .
- Problems related to embedding subgraphs based on the number of edges in the host graph are called **Turán-type problems**.

# Larger substructures

- Suppose we have 16 sports teams. Due to travel restrictions, some teams cannot play each other. How many pairs of teams that can play each other do we need to ensure that we can pair off all teams?

# Larger substructures

- Suppose we have 16 sports teams. Due to travel restrictions, some teams cannot play each other. How many pairs of teams that can play each other do we need to ensure that we can pair off all teams?
- If even one team is unable to play with all the other teams, it can't be done.

# Larger substructures

- Suppose we have 16 sports teams. Due to travel restrictions, some teams cannot play each other. How many pairs of teams that can play each other do we need to ensure that we can pair off all teams?
- If even one team is unable to play with all the other teams, it can't be done.
- Say we have a network of cities and roads between them. We have important deliveries to make in each city and we need to come back to where we started at the end of the day. How many roads do we need to guarantee that we need to visit each city only once?



# Larger substructures

- Suppose we have 16 sports teams. Due to travel restrictions, some teams cannot play each other. How many pairs of teams that can play each other do we need to ensure that we can pair off all teams?
- If even one team is unable to play with all the other teams, it can't be done.
- Say we have a network of cities and roads between them. We have important deliveries to make in each city and we need to come back to where we started at the end of the day. How many roads do we need to guarantee that we need to visit each city only once?
- We can't do this if there's only one road into any particular city.

# Larger substructures

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

# Larger substructures

## Definition

Let  $G = (V, E)$  be a graph. A subset of edges  $\mathcal{M} \subseteq E$  is called a **matching** if its edges are vertex-disjoint. The matching  $\mathcal{M}$  is **perfect** if the vertices that comprise it cover all of  $V$ .

## Definition

Let  $G = (V, E)$  be a graph and let  $\mathcal{C}$  be a sequence of edges. Then  $\mathcal{C}$  is a **Hamiltonian cycle** if it is a cycle that visits every vertex in  $G$  exactly once.

# Larger substructures




Figure: A graph

# Larger substructures




Figure: A perfect matching

# Larger substructures




Figure: A Hamiltonian cycle

# Another perspective

- How many edges do we need to force a perfect matching or Hamiltonian cycle?

# Another perspective

- How many edges do we need to force a perfect matching or Hamiltonian cycle?
- This question is a bit different from asking about  $\text{ex}(n, H)$  for some  $H$  since now our subgraph isn't fixed - it depends on  $n$ .



# Another perspective

- How many edges do we need to force a perfect matching or Hamiltonian cycle?
- This question is a bit different from asking about  $ex(n, H)$  for some  $H$  since now our subgraph isn't fixed - it depends on  $n$ .
- The Turán perspective doesn't tell us much about perfect matchings - start with  $K_n$  and delete all edges incident to one fixed vertex. The resulting graph has no perfect matching.

# Another perspective

- How many edges do we need to force a perfect matching or Hamiltonian cycle?
- This question is a bit different from asking about  $\text{ex}(n, H)$  for some  $H$  since now our subgraph isn't fixed - it depends on  $n$ .
- The Turán perspective doesn't tell us much about perfect matchings - start with  $K_n$  and delete all edges incident to one fixed vertex. The resulting graph has no perfect matching.

## Theorem (P. Hall - 1935)

*Let  $G$  be a bipartite graph with parts  $X$  and  $Y$  (all edges in  $G$  have one end in  $X$  and one end in  $Y$ ). Then  $G$  contains a matching that saturates  $X$  if and only if every subset  $W$  of  $X$  has at least  $|W|$  neighbors in  $Y$ .*

## Another perspective

- Similarly, if any vertex has degree less than 2, we cannot have a Hamiltonian cycle.

# Another perspective

- Similarly, if any vertex has degree less than 2, we cannot have a Hamiltonian cycle.

## Theorem (G. Dirac - 1952)

*A graph on  $n \geq 3$  vertices contains a Hamiltonian cycle if  $\delta(G) \geq n/2$  (graphs with such minimum degree are called **Dirac graphs**). In particular, such a graph contains a perfect matching if  $n$  is even.*

# Another perspective

- Similarly, if any vertex has degree less than 2, we cannot have a Hamiltonian cycle.

## Theorem (G. Dirac - 1952)

*A graph on  $n \geq 3$  vertices contains a Hamiltonian cycle if  $\delta(G) \geq n/2$  (graphs with such minimum degree are called **Dirac graphs**). In particular, such a graph contains a perfect matching if  $n$  is even.*

- Apparently, some structures aren't simply related to the number of edges in the host graph, but how they're distributed.

## Another perspective

- Similarly, if any vertex has degree less than 2, we cannot have a Hamiltonian cycle.

### Theorem (G. Dirac - 1952)

*A graph on  $n \geq 3$  vertices contains a Hamiltonian cycle if  $\delta(G) \geq n/2$  (graphs with such minimum degree are called **Dirac graphs**). In particular, such a graph contains a perfect matching if  $n$  is even.*

- Apparently, some structures aren't simply related to the number of edges in the host graph, but how they're distributed.
- Graph embedding problems relating to the degree of the host graph are called **Dirac-type problems**.

- 1 Graph substructures
- 2 Counting perfect matchings and Hamiltonian cycles
- 3 Degrees of prescribed remainder

# From one to many

- If a graph has sufficiently high minimum degree, it contains a Hamiltonian cycle (and a perfect matching if there is an even number of vertices). Could it have many?



# From one to many

- If a graph has sufficiently high minimum degree, it contains a Hamiltonian cycle (and a perfect matching if there is an even number of vertices). Could it have many?
- The **Erdős-Renyi random graph**,  $\mathcal{G}(n, p)$ , is the random variable that outputs a graph on  $n$  vertices, any two of which are independently connected with probability  $p$ . It's one of the most well-studied random structures.

# From one to many - intuition from random graphs

- Idea: if  $p > 1/2$ , then  $G \sim \mathcal{G}(n, p)$  is Dirac whp. Maybe we can use probabilistic tools to estimate the number of perfect matchings/Hamiltonian cycles in  $G$ . Does this tell us anything about deterministic Dirac graphs?

# From one to many - intuition from random graphs

- Idea: if  $p > 1/2$ , then  $G \sim \mathcal{G}(n, p)$  is Dirac whp. Maybe we can use probabilistic tools to estimate the number of perfect matchings/Hamiltonian cycles in  $G$ . Does this tell us anything about deterministic Dirac graphs?

- Expected number of perfect matchings ( $n$  even):

$$\frac{n!}{2^{n/2}(n/2)!} p^{n/2}.$$

# From one to many - intuition from random graphs

- Idea: if  $p > 1/2$ , then  $G \sim \mathcal{G}(n, p)$  is Dirac whp. Maybe we can use probabilistic tools to estimate the number of perfect matchings/Hamiltonian cycles in  $G$ . Does this tell us anything about deterministic Dirac graphs?
- Expected number of perfect matchings ( $n$  even):  

$$\frac{n!}{2^{n/2}(n/2)!} p^{n/2}.$$
- Expected number of Hamiltonian cycles:  $\frac{1}{2}(n-1)!p^n$ .

# From one to many - intuition from random graphs

- Idea: if  $p > 1/2$ , then  $G \sim \mathcal{G}(n, p)$  is Dirac whp. Maybe we can use probabilistic tools to estimate the number of perfect matchings/Hamiltonian cycles in  $G$ . Does this tell us anything about deterministic Dirac graphs?
- Expected number of perfect matchings ( $n$  even):  

$$\frac{n!}{2^{n/2}(n/2)!} p^{n/2}.$$
- Expected number of Hamiltonian cycles:  $\frac{1}{2}(n-1)!p^n$ .
- By Markov's inequality,  $G \sim \mathcal{G}(n, p)$  doesn't have "too many" Hamiltonian cycles (perfect matchings) whp.

# From one to many - intuition from random graphs

- Idea: if  $p > 1/2$ , then  $G \sim \mathcal{G}(n, p)$  is Dirac whp. Maybe we can use probabilistic tools to estimate the number of perfect matchings/Hamiltonian cycles in  $G$ . Does this tell us anything about deterministic Dirac graphs?
- Expected number of perfect matchings ( $n$  even):  
$$\frac{n!}{2^{n/2}(n/2)!} p^{n/2}.$$
- Expected number of Hamiltonian cycles:  $\frac{1}{2}(n-1)!p^n$ .
- By Markov's inequality,  $G \sim \mathcal{G}(n, p)$  doesn't have "too many" Hamiltonian cycles (perfect matchings) whp.
- Glebov and Krivelevich showed that if  $p = \frac{\log n + \log \log n + \omega(1)}{n}$ , then the number of Hamiltonian cycles in  $G \sim \mathcal{G}(n, p)$ , up to a sub-exponential factor, concentrates about its mean.

# From one to many - intuition from random graphs

Theorem (B. Cuckler, J. Kahn - 2009)

If  $G$  is a graph on  $n$  vertices with  $\delta(G) \geq n/2$ , then

$$\#\{\text{perfect matchings in } G\} \geq (1 - o(1))^n \cdot \frac{n!}{2^n (n/2)!}$$

$$\#\{\text{Hamiltonian cycles in } G\} \geq (1 - o(1))^n \cdot \frac{1}{2^n} (n-1)!$$

# From one to many - intuition from random graphs

## Theorem (B. Cuckler, J. Kahn - 2009)

If  $G$  is a graph on  $n$  vertices with  $\delta(G) \geq n/2$ , then

$$\#\{\text{perfect matchings in } G\} \geq (1 - o(1))^n \cdot \frac{n!}{2^n (n/2)!}$$

$$\#\{\text{Hamiltonian cycles in } G\} \geq (1 - o(1))^n \cdot \frac{1}{2^n} (n-1)!$$

- Edge disjoint cycles? Covering? Directed graphs?



## A riddle

*Fifteen young ladies in a school walk out three abreast seven days in succession: it is required to arrange them daily so that no two shall walk twice abreast.*

# A riddle

## Problem (T. Kirkman - 1850)

*Fifteen young ladies in a school walk out three abreast seven days in succession: it is required to arrange them daily so that no two shall walk twice abreast.*

- Consider the set of ladies  $H$  and let  $E$  be the collection of subsets of  $H$  of the form  $\{\ell_1, \ell_2, \ell_3\}$ , where the ladies  $\ell_1$ ,  $\ell_2$  and  $\ell_3$  walk together on some day.

# A riddle

## Problem (T. Kirkman - 1850)

*Fifteen young ladies in a school walk out three abreast seven days in succession: it is required to arrange them daily so that no two shall walk twice abreast.*

- Consider the set of ladies  $H$  and let  $E$  be the collection of subsets of  $H$  of the form  $\{\ell_1, \ell_2, \ell_3\}$ , where the ladies  $\ell_1$ ,  $\ell_2$  and  $\ell_3$  walk together on some day.
- Sounds like a graph, except edges are triples!

# A riddle

## Problem (T. Kirkman - 1850)

*Fifteen young ladies in a school walk out three abreast seven days in succession: it is required to arrange them daily so that no two shall walk twice abreast.*

- Consider the set of ladies  $H$  and let  $E$  be the collection of subsets of  $H$  of the form  $\{\ell_1, \ell_2, \ell_3\}$ , where the ladies  $\ell_1, \ell_2$  and  $\ell_3$  walk together on some day.
- Sounds like a graph, except edges are triples!
- Kirkman's problem is asking us to find something resembling a collection of perfect matchings.

# A generalization - hypergraphs

- A **hypergraph**  $H = (V, E)$  consists of a set of vertices  $V$  and a set of edges  $E \subseteq 2^V$ . We say that  $H$  is  $k$ -uniform if every edge consists of exactly  $k$  vertices (note that a graph is a 2-uniform hypergraph).

# A generalization - hypergraphs

- A **hypergraph**  $H = (V, E)$  consists of a set of vertices  $V$  and a set of edges  $E \subseteq 2^V$ . We say that  $H$  is  $k$ -uniform if every edge consists of exactly  $k$  vertices (note that a graph is a 2-uniform hypergraph).
- We can talk about Turán or Dirac-type problems in this setting too.

# A generalization - hypergraphs

- A **hypergraph**  $H = (V, E)$  consists of a set of vertices  $V$  and a set of edges  $E \subseteq 2^V$ . We say that  $H$  is  $k$ -uniform if every edge consists of exactly  $k$  vertices (note that a graph is a 2-uniform hypergraph).
- We can talk about Turán or Dirac-type problems in this setting too.
- More generalized notion of degree: if  $S$  is any subset, then  $d(S)$  is the number of edges that contain  $S$ .  $\delta_t(H)$  is the minimum of  $d(S)$  over all subsets of size  $t$ .

# Generalizing the substructures

- Can we extend the notions of matchings and cycles to hypergraphs?



# Generalizing the substructures

- Can we extend the notions of matchings and cycles to hypergraphs?
- Matchings seem fine.

# Generalizing the substructures

- Can we extend the notions of matchings and cycles to hypergraphs?
- Matchings seem fine.
- Cycles are a bit less clear since the edges in graph cycles overlap at the ends. How can we account for this in hypergraphs?

# Generalizing the substructures

- Can we extend the notions of matchings and cycles to hypergraphs?
- Matchings seem fine.
- Cycles are a bit less clear since the edges in graph cycles overlap at the ends. How can we account for this in hypergraphs?
- We say that a  $k$ -uniform hypergraph  $H$  contains a **Hamiltonian  $\ell$ -cycle** if there is a cyclic ordering of the vertices of  $H$  such that the edges of the cycle are segments of length  $k$  in this ordering and any two consecutive edges  $f_i$ ,  $f_{i+1}$  share exactly  $\ell$  vertices.

# Example




Figure: A hypergraph

# Example




Figure: Top:  $k = 6, \ell = 2$ . Bottom:  $k = 6, \ell = 4$ .

# Matchings and cycles in hypergraphs

- Notice that if  $\ell = 0$ , then a Hamiltonian  $\ell$ -cycle is a perfect matching.

# Matchings and cycles in hypergraphs

- Notice that if  $\ell = 0$ , then a Hamiltonian  $\ell$ -cycle is a perfect matching.
- If  $\ell > k/2$ , *three* consecutive edges in an  $\ell$ -cycle overlap. This ostensibly makes counting these more nuanced.

# Matchings and cycles in hypergraphs

- Notice that if  $\ell = 0$ , then a Hamiltonian  $\ell$ -cycle is a perfect matching.
- If  $\ell > k/2$ , *three* consecutive edges in an  $\ell$ -cycle overlap. This ostensibly makes counting these more nuanced.
- For  $\ell \leq k/2$ , the number of Hamiltonian  $\ell$ -cycles we expect to see in a random hypergraph with edge probability  $p$  is

$$(n-1)! \cdot \frac{k-\ell}{2} \cdot \left( \frac{p}{\ell!(k-2\ell)!} \right)^{\frac{n}{k-\ell}}.$$



# Matchings and cycles in hypergraphs

Theorem (A. Ferber, M. Krivelevich, B. Sudakov - 2016)

*Let  $0 \leq \ell < k/2$  and let  $1/2 < p \leq 1$ . If  $(k - \ell) \mid n$  and  $\delta_{k-1} \geq pn$ , then the number of Hamiltonian  $\ell$ -cycles in the  $k$ -uniform hypergraph  $H$  is at least*

$$(1 - o(1))^n \cdot n! \cdot \left( \frac{p}{\ell!(k - 2\ell)!} \right)^{\frac{n}{k - \ell}}.$$

# Matchings and cycles in hypergraphs

Theorem (A. Ferber, M. Krivelevich, B. Sudakov - 2016)

*Let  $0 \leq \ell < k/2$  and let  $1/2 < p \leq 1$ . If  $(k - \ell) \mid n$  and  $\delta_{k-1} \geq pn$ , then the number of Hamiltonian  $\ell$ -cycles in the  $k$ -uniform hypergraph  $H$  is at least*

$$(1 - o(1))^n \cdot n! \cdot \left( \frac{p}{\ell!(k - 2\ell)!} \right)^{\frac{n}{k - \ell}}.$$

- Notice that the minimum **co-degree**,  $\delta_{k-1}(H)$  plays a role analogous to the minimum degree in a graph. A **Dirac hypergraph** on  $n$  vertices is one whose minimum co-degree is at least  $n/2$ .

1. *Journal of the American Medical Association*, 2000; 284: 2689-2694.

# Matchings and cycles in hypergraphs - current work

- Extending the work of Ferber, Krivelevich and Sudakov to the case  $\ell < k - 1$ . Joint with A. Mond (University of Cambridge)
- Future work: a tight (up to subexponential factor) bound for tight ( $\ell = k - 1$ ) cycles; number of *disjoint*  $\ell$ -cycles.

- 1 Graph substructures
- 2 Counting perfect matchings and Hamiltonian cycles
- 3 Degrees of prescribed remainder

# A motivating problem

## Exercise (L. Lovász, T. Gallai - 1979)

*Assume that at each vertex of a graph there is a light that is turned on. Toggling any light toggles all of its neighbors as well. Turn off all of the lights.*

# A motivating problem

## Exercise (L. Lovász, T. Gallai - 1979)

*Assume that at each vertex of a graph there is a light that is turned on. Toggling any light toggles all of its neighbors as well. Turn off all of the lights.*

- Solution: find some  $X \subseteq V(G)$  where all degrees in  $G[X]$  are even and every  $v \notin X$  has odd degree into  $X$ . Then toggle every light in  $X$ .

# A motivating problem

## Exercise (L. Lovász, T. Gallai - 1979)

*Assume that at each vertex of a graph there is a light that is turned on. Toggling any light toggles all of its neighbors as well. Turn off all of the lights.*

- Solution: find some  $X \subseteq V(G)$  where all degrees in  $G[X]$  are even and every  $v \notin X$  has odd degree into  $X$ . Then toggle every light in  $X$ .
- Every light in  $V \setminus X$  flips an odd number of times (each of its neighbors in  $X$ ) and every light in  $X$  flips an odd number of times (each of its neighbors in  $X$  and itself).



# Even degrees

## Theorem (L. Lovász, T. Gallai - 1979)

*(Stated as an exercise in Lovász' book). Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

# Even degrees

## Theorem (L. Lovász, T. Gallai - 1979)

*(Stated as an exercise in Lovász' book). Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

- Use this theorem to prove that our solution is possible.

# Even degrees

## Theorem (L. Lovász, T. Gallai - 1979)

*(Stated as an exercise in Lovász' book). Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

- Use this theorem to prove that our solution is possible.
- This theorem contains two statements: one about a large induced subgraph and another about partitioning into particular subgraphs.

# Odd subgraphs

- What about odd subgraphs?

# Odd subgraphs

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

# Odd subgraphs

- What about odd subgraphs?

## Theorem (A. Scott - 1992)

*Every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq \frac{n}{900 \log n}$  and  $G[W]$  has all degrees odd.*

## Conjecture (A. Scott - 2001)

*There exists some constant  $c > 0$  such that every graph  $G(V, E)$  on  $n$  vertices, none of which are isolated, contains a set  $W \subseteq V(G)$  such that  $|W| \geq cn$  and  $G[W]$  has all degrees odd.*

# Other moduli and remainders

## Theorem (A. Scott - 1992)

*Let  $G \sim \mathcal{G}(n, 1/2)$ . Then with high probability,  $G$  has an induced subgraph on at least  $0.7729n$  vertices with all degrees odd.*

# Other moduli and remainders

## Theorem (A. Scott - 1992)

*Let  $G \sim \mathcal{G}(n, 1/2)$ . Then with high probability,  $G$  has an induced subgraph on at least  $0.7729n$  vertices with all degrees odd.*

## Conjecture (A. Scott - 2001)

*For any positive integer  $q$  at least 2, there exists a constant  $c_q$  so that every graph on  $n$  vertices without isolated vertices has an induced subgraph on at least  $c_q n$  vertices with all degrees  $1 \pmod{q}$ .*



# Other moduli and Remainders

Theorem (A. Ferber, H., M. Krivelevich - 2020+)

*Let  $q \geq 2$  and let  $r$  be an integer. Then there exists a constant  $c_q$  such that, with high probability, the random graph  $G \sim \mathcal{G}(n, 1/2)$  has an induced subgraph on at least  $c_q n$  vertices with all degrees  $r \pmod{q}$ .*

# Key idea behind proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.

# Key idea behind proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.
- In this case, the  $i$ -th entry of  $M\mathbf{1}$  is the degree of vertex  $i$  in  $G$ , where  $\mathbf{1}$  is the length  $n$  all-1 vector.

# Key idea behind proof

- If  $G \sim \mathcal{G}(n, 1/2)$ , then its adjacency matrix  $M$  ( $M_{ij} = 1$  if and only vertices  $i$  and  $j$  are connected and  $M_{ij} = 0$  otherwise) is a random symmetric  $n \times n$  matrix whose above-diagonal entries are iid  $\text{Bern}(1/2)$  random variables.
- In this case, the  $i$ -th entry of  $M\mathbf{1}$  is the degree of vertex  $i$  in  $G$ , where  $\mathbf{1}$  is the length  $n$  all-1 vector.

## Observation

*$G$  contains an induced subgraph with all degrees  $r \pmod{q}$  if and only if its adjacency matrix contains a principal submatrix  $B$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .*

# Proof (sketch) of theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

# Proof (sketch) of theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

- Let  $k = c_q n$  for some  $c_q > 0$  and let  $X_k$  be the number of  $k \times k$  principal submatrices  $B$  of the adjacency matrix of  $G$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .

# Proof (sketch) of theorem

## Theorem (Chebyshev's Inequality)

*Let  $X$  be a nonnegative integer-valued random variable with finite variance. Then*

$$\Pr[X > 0] \geq 1 - \frac{\text{Var}[X]}{(\mathbb{E}[X])^2}.$$

- Let  $k = c_q n$  for some  $c_q > 0$  and let  $X_k$  be the number of  $k \times k$  principal submatrices  $B$  of the adjacency matrix of  $G$  satisfying  $B\mathbf{1} \equiv r\mathbf{1} \pmod{q}$ .
- Show that  $c_q$  can be chosen so that  $\text{Var}[X_k] = o(\mathbb{E}[X_k]^2)$ . Then  $G$  has an induced subgraph of size  $c_q n$  with high probability.

# Key computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables. If  $\delta_0(x) = 1$  if and only if  $x \equiv 0 \pmod{q}$  and  $\delta_0(x) = 0$  otherwise, then



# Key computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables. If  $\delta_0(x) = 1$  if and only if  $x \equiv 0 \pmod{q}$  and  $\delta_0(x) = 0$  otherwise, then

$$\Pr[x_1 + \dots + x_t \equiv r] = \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)]$$

# Key computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables. If  $\delta_0(x) = 1$  if and only if  $x \equiv 0 \pmod{q}$  and  $\delta_0(x) = 0$  otherwise, then

$$\begin{aligned} \Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right] \end{aligned}$$

# Key computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables. If  $\delta_0(x) = 1$  if and only if  $x \equiv 0 \pmod{q}$  and  $\delta_0(x) = 0$  otherwise, then

$$\begin{aligned} \Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right] \\ &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \prod_{j=1}^t \mathbb{E} e^{2\pi i \ell x_j / q} \end{aligned}$$

# Key computation

Fix a positive integer  $q \geq 2$  and let  $r$  be an integer. Let  $x_1, \dots, x_t$  be iid  $\text{Bern}(1/2)$  random variables. If  $\delta_0(x) = 1$  if and only if  $x \equiv 0 \pmod{q}$  and  $\delta_0(x) = 0$  otherwise, then

$$\begin{aligned}
 \Pr[x_1 + \dots + x_t \equiv r] &= \mathbb{E}[\delta_0(x_1 + \dots + x_t - r)] \\
 &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} \mathbb{E} \exp \left[ \frac{2\pi i \ell}{q} (x_1 + \dots + x_t - r) \right] \\
 &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \prod_{j=1}^t \mathbb{E} e^{2\pi i \ell x_j / q} \\
 &= \frac{1}{q} \sum_{\ell \in \mathbb{Z}_q} e^{-2\pi i r \ell / q} \left( \frac{1 + e^{2\pi i \ell / q}}{2} \right)^t.
 \end{aligned}$$

# Key computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

# Key computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

# Key computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

- Calculus trick:  $|\cos(\pi\ell/q)| \leq e^{-2/q^2}$  for all  $\ell = 1, \dots, q-1$ .

# Key computation

- Isolate the  $\ell \equiv 0$  term, apply the triangle inequality and Euler's identity.

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{1}{q} \sum_{\ell=1}^{q-1} |\cos(\pi\ell/q)|^t.$$

- Calculus trick:  $|\cos(\pi\ell/q)| \leq e^{-2/q^2}$  for all  $\ell = 1, \dots, q-1$ .

$$\left| \Pr[x_1 + \cdots + x_t \equiv r] - \frac{1}{q} \right| \leq \frac{q-1}{q} e^{-2t/q^2}.$$



# Key computation

- The distribution of a random Bernoulli sum modulo  $q$  is asymptotically uniform!

# Key computation

- The distribution of a random Bernoulli sum modulo  $q$  is asymptotically uniform!
- Using this, we can show that if  $M$  is the adjacency matrix of a random graph, then  $M\mathbf{1} \pmod{q}$  is (asymptotically) distributed uniformly over all possible values it can take.

# Partitioning

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

# Partitioning

## Theorem (L. Lovász, T. Gallai - 1979)

*Let  $G = (V, E)$  be any graph. Then  $G$  admits a partitioning of its vertex set into two parts,  $V = V_1 \cup V_2$ , so that each vertex in  $G[V_1]$  and each vertex in  $G[V_2]$  has even degree. In particular, any graph on  $n$  vertices has an even subgraph of order at least  $n/2$ .*

## Theorem (A. Ferber, H., M. Krivelevich, 2020+)

*For  $q > 1$  and  $r \leq q$ ,  $G \sim \mathcal{G}(n, 1/2)$  admits a partition into  $q + 1$  classes such that the degrees in each class are congruent to  $r \pmod{q}$  with high probability.*

# Future work - discrete Fourier analysis

## Theorem (A. Ferber - 2020)

*Let  $M_n$  denote an  $n \times n$  symmetric  $\pm 1$  matrix chosen uniformly from all such matrices. Let  $p(n)$  denote the probability that  $M_n$  is singular. Then there exists some constant  $C > 0$  for which*

$$p(n) = O\left(\frac{\log^C n}{\sqrt{n}}\right).$$

# Future work - discrete Fourier analysis

## Theorem (R. Meshulam - 1990)

*Let  $G$  be a finite abelian group and let  $s(G)$  denote the maximal  $s$  for which there exists a sequence  $a_1, \dots, a_s \in G$  such that  $\sum_{i \in I} a_i \neq 0$  for any nonempty  $I \subseteq [s]$ . Then if  $m$  is the maximum order of elements in  $G$ , we have*

$$s(G) \leq m \left( 1 + \log \frac{|G|}{m} \right).$$

# Future work - discrete Fourier analysis

Conjecture (N. Alon, N. Linial, R. Meshulam - 1988)

*For every prime  $p$ , there exists a constant  $c_p$  such that for any  $n \geq 1$ , the union of any  $c_p$  linear bases for  $\mathbb{Z}_p$  forms an additive basis for  $\mathbb{Z}_p$ .*

# Future work - discrete Fourier analysis

Conjecture (N. Alon, N. Linial, R. Meshulam - 1988)

*For every prime  $p$ , there exists a constant  $c_p$  such that for any  $n \geq 1$ , the union of any  $c_p$  linear bases for  $\mathbb{Z}_p$  forms an additive basis for  $\mathbb{Z}_p$ .*

- The best known bound is at most  $c_p \log n$  many linear bases.



*Thank you.*