

Problem Statement

Consider the integro-differential equation

$$\frac{d^2 h}{dx^2} + h + \frac{i}{2} \int_{-\pi/2}^{\pi/2} H_0^1(|x-y|) h(y) dy = e^{iax} \quad (1)$$

for $-\pi/2 \leq x \leq \pi/2$. The complex dependent variable $h(x)$ satisfies the boundary conditions $h(\pi/2) = h(-\pi/2) = 0$. This equation describes the interface displacement resulting from the scattering of an acoustic wave from an interface separating two fluids. Here H_0^1 is a Hankel function of the first kind.

1. Describe a numerical method to solve this differential equation.
2. Solve this equation numerically for $a = 0.1$. Do a numerical convergence check to show convergence of your numerical scheme and use these results to estimate the rate of convergence of your scheme. Plot the solution.
3. Solve the equation for $a = 1.0$.
4. Explain the differences between the $a = 0.1$ case and the $a = 1.0$ case. When can such differences be expected?

Methods

General Strategy

We will solve the equation by discretizing the domain as well as the region of integration uniformly in grid space. Given a resolution N , we define

$$x_i, y_i = \pi \left(\frac{i}{N} - \frac{1}{2} \right) \quad (2)$$

where $i = 0, 1, 2, \dots, N$ including both boundary points. The second-order center finite difference operator at a point x_i is given by

$$h''(x_i) \approx \frac{h(x_{i+1}) - 2h(x_i) + h(x_{i-1}))}{\Delta x^2} \quad (3)$$

where the grid spacing $\Delta x = \Delta y = \frac{\pi}{N}$. Accordingly we define a finite difference matrix operator D whose entries satisfy equation 3 for the inside points. The integral term will be approximated via repeated trapezoid product integration quadrature. This is the crux of the problem. When this is done properly, we can approximate the terms on the left hand side of equation 1 as a matrix acting on some unknown solution vector \mathbf{h} . The system can then be solved readily via Gaussian elimination. Special care must be taken when dealing with the integral term to ensure fast convergence.

The Integrand

First we reproduce a few well-established properties of the Hankel function, namely

$$H_n^{(1)}(y) = J_n(y) + iY_n(y) \quad (4)$$

where $J_n(y)$ and $Y_n(y)$ are Bessel functions of the first and second kinds respectively [1]. Recall that J_n has series expansion

$$J_n(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k+n} k! (n+k)!} z^{2k+n} \quad (5)$$

given by [2]. We also reproduce the series expansion of Y_0 , given by [3] as

$$Y_0(z) = \frac{2}{\pi} J_0(z) \log\left(\frac{z}{2}\right) - \frac{2}{\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{k! k!} \left(\frac{z}{2}\right)^{2k} \psi(k+1) \quad (6)$$

where ψ is the digamma function. We note the singular point at $x = 0$. We continue by decomposing the complex functions into their real and imaginary components, substituting into equation 1, and equating the real and imaginary components. Letting $h(x) = u(x) + iv(x)$, we have that

$$\frac{d^2(u(x) + iv(x))}{dx^2} + (u(x) + iv(x)) + \frac{i}{2} \int_{-\pi/2}^{\pi/2} \left(J_0(|x-y|) + iY_0(|x-y|) \right) (u(y) + iv(y)) dy = e^{iax} \quad (7)$$

becomes

$$\frac{d^2u}{dx^2} + u - \frac{1}{2} \int_{-\pi/2}^{\pi/2} J_0(|x-y|) v(y) + Y_0(|x-y|) u(y) dy = \cos(ax) \quad (8)$$

$$\frac{d^2v}{dx^2} + v + \frac{1}{2} \int_{-\pi/2}^{\pi/2} J_0(|x-y|) u(y) - Y_0(|x-y|) v(y) dy = \sin(ax) \quad (9)$$

$$(10)$$

With this in mind we now outline how the integral itself will be approximated.

Quadrature

Recall that a smooth function's integral can be approximated to second order as

$$\int_{-\pi/2}^{\pi/2} f(x) dx = \frac{\Delta x}{2} f_0 + \Delta x \sum_{k=1}^{N-1} f_k + \frac{\Delta x}{2} f_N \quad (11)$$

where the shorthand f_k refers to $f(x_k)$. The nonsingular terms can all be dealt with in this way whereas the singular term requires a bit more care. We apply the product integration method. For some "fixed" x_i have have that

$$\int_{-\pi/2}^{\pi/2} \log\left(\frac{|x_i - y|}{2}\right) g(y) dy = \int_{-\pi/2}^{x_i} \log\left(\frac{x_i - y}{2}\right) g(y) dy + \int_{x_i}^{\pi/2} \log\left(\frac{y - x_i}{2}\right) g(y) dy. \quad (12)$$

Replacing the integrands with the relevant expressions and the integrals with some yet-unknown kernel yields

$$\int_{-\pi/2}^{\pi/2} \log\left(\frac{|x_i - y|}{2}\right) J_0(x_i - y) f(y) dy = \sum_{j=0}^{j=i} w_{ij} f_j = \int_{-\pi/2}^{x_i} \log\left(\frac{x_i - y}{2}\right) J_0(x_i - y) f(y) dy \quad (13)$$

$$+ \sum_{j=i}^{j=N} w_{ij} f_j + \int_{x_i}^{-\pi/2} \log\left(\frac{y - x_i}{2}\right) J_0(y - x_i) f(y) dy \quad (14)$$

Where the integrands are meant to describe the nonlinear contributions of equation 1. In practice $f(y)$ is some unknown function related to the solution's real or imaginary component. Note that we include the endpoints in the summations. Though it appears at first sight as though we are over-constraining the kernel weights w_{ij} with the above, this is in fact not the case due to the bounds on the summations. We must require, however that w_{ij} agree at $j = i$. Next we discretize such that our kernel is of the same dimension as the finite difference matrix. Recall that $x_j = y_j$. We then have

$$= \sum_{j=0}^{i-1} \int_{y_j}^{y_{j+1}} \log\left(\frac{x_i - y}{2}\right) J_0(x_i - y) f(y) dy \quad (15)$$

$$+ \sum_{j=i}^{N-1} \int_{y_j}^{y_{j+1}} \log\left(\frac{y - x_i}{2}\right) J_0(y - x_i) f(y) dy \quad (16)$$

To evaluate this integral “analytically” we would need f , which is related to our solution. Instead we will pretend we know the value of f at each of our grid points. Using interpolation, we have evaluate the integral while keeping the problem linear. There are a huge number of interpolation methods (cubic splines, Bézier curves, B-splines, etc... this could be a course of its own). We will be lazy and use linear interpolation, as this ought to agree with the anticipated second-order convergence of the trapezoidal method. At some arbitrary y value we would then have

$$f(y) \approx \left(\frac{y_{j+1} - y}{\Delta y}\right) f_j + \left(\frac{y - y_j}{\Delta y}\right) f_{j+1} \quad (17)$$

where we take j to be such that $y_j < y < y_{j+1}$.

Substitution then gives

$$\sum_{j=0}^{j=i} w_{ij} f_j = \sum_{j=0}^{i-1} \int_{y_j}^{y_{j+1}} \log\left(\frac{x_i - y}{2}\right) J_0(x_i - y) \left(\left(\frac{y_{j+1} - y}{\Delta y}\right) f_j + \left(\frac{y - y_j}{\Delta y}\right) f_{j+1}\right) dy \quad (18)$$

$$+ \sum_{j=i}^{j=N} w_{ij} f_j + \sum_{j=i}^{N-1} \int_{y_j}^{y_{j+1}} \log\left(\frac{y - x_i}{2}\right) J_0(y - x_i) \left(\left(\frac{y_{j+1} - y}{\Delta y}\right) f_j + \left(\frac{y - y_j}{\Delta y}\right) f_{j+1}\right) dy \quad (19)$$

Setting $i = 1, 2, 3, \dots$ allows us to obtain an expression for each of the weights

$$w_{ij} = \frac{2}{\pi \Delta y} \int_{y_{j-1}}^{y_j} \ln\left(\frac{y_i - y}{2}\right) J_0(y_i - y) (y - y_{i-1}) dy \quad (20)$$

$$+ \frac{2}{\pi \Delta y} \int_{y_j}^{y_{j+1}} \ln\left(\frac{y - y_1}{2}\right) J_0(y - y_1) (y_{i+1} - y_i) dy. \quad (21)$$

Then, for convenience, we define the linear operator A such that

$$\frac{1}{2} \sum_{j=0}^N w_{ij} f_j = A \mathbf{f} \cdot \hat{\mathbf{e}}_i. \quad (22)$$

where the leading $1/2$ factor is to reflect the $i/2$ factor on the integral term in equation 1 (multiplication of i has already been accounted for). A corresponding B exists for the imaginary-valued subproblem. Finally we define the real and imaginary component vectors \mathbf{b} and \mathbf{c} of the nonhomogeneous terms where

$$b_i = \cos(ax) \quad (23)$$

$$c_i = \sin(ax). \quad (24)$$

Our problem can then be collapsed into a single linear equation which approximates the real and imaginary components of equation 1 as

$$\begin{bmatrix} D + I - A & -B \\ B & D + I - A \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix} = \begin{bmatrix} \mathbf{b} \\ \mathbf{c} \end{bmatrix}. \quad (25)$$

From here we need only use a standard linear solver to obtain the approximate solution.

References

- [1] Weisstein, Eric W. "Hankel Function of the First Kind." From MathWorld—A Wolfram Web Resource. <https://mathworld.wolfram.com/HankelFunctionoftheFirstKind.html>
- [2] Weisstein, Eric W. "Hankel Weisstein, Eric W. "Bessel Function of the First Kind." From MathWorld—A Wolfram Web Resource. <https://mathworld.wolfram.com/BesselFunctionoftheFirstKind.html>
- [3] Lambers, Jim. "Bessel Functions of the Second Kind" <http://www.math.usm.edu/lambers/mat415/lecture16.pdf>