

# Marginally-Stable Thermal Equilibria of Rayleigh-Bénard Convection

Liam O'Connor<sup>1,2</sup>

<sup>1</sup>*Department of Engineering Sciences and Applied Mathematics,  
Northwestern University, Evanston, IL 60208 USA and*

<sup>2</sup>*Center for Interdisciplinary Exploration and Research in Astrophysics,  
Northwestern University, Evanston, IL, 60201 USA*

Natural convection exhibits turbulent flows which are difficult or impossible to resolve in direct numerical simulations. In this work, we investigate a quasilinear form of the Rayleigh-Bénard problem which describes the bulk one-dimensional properties of convection without resolving the turbulent dynamics. We represent perturbations away from the mean using a sum of marginally-stable eigenmodes. By constraining the perturbation amplitudes, the marginal stability criterion allows us to evolve the background temperature profile under the influence of multiple eigenmodes representing flows at different length scales. We find the quasilinear system evolves to an equilibrium state where advective and diffusive fluxes sum to a constant. These marginally-stable thermal equilibria (MSTE) are exact solutions of the quasilinear equations. The mean MSTE temperature profiles have thinner boundary layers and larger Nusselt numbers than thermally-equilibrated 2D and 3D simulations of the full nonlinear equations. MSTE solutions exhibit a classic boundary-layer scaling of the Nusselt number  $Nu$  with the Rayleigh number  $Ra$  of  $Nu \sim Ra^{1/3}$ . When an MSTE is used as initial conditions for a 2D simulation, we find that  $Nu$  quickly equilibrates without the burst of turbulence often induced by purely conductive initial conditions, but we also find that the kinetic energy is too large and viscously attenuates on a long viscous time scale.

## I. INTRODUCTION

Rayleigh-Bénard convection plays a foundational role in astrophysical and geophysical settings. The resulting buoyancy-driven flows regulate heat transfer and generate large-scale vortices [1]. Turbulent convection, which

is associated with large Rayleigh numbers  $Ra$ , is difficult to simulate. State of the art simulations performed by [2] have reached  $Ra \sim 10^{14}$  but estimates for the sun's convective zone and earth's interior are  $Ra \sim 10^{16} - 10^{20}$  and  $Ra \sim 10^{20} - 10^{30}$  respectively [3, 4].

- 
- [1] L.-A. Coustou, D. Lecoanet, B. Favier, and M. Le Bars, Shape and size of large-scale vortices: A generic fluid pattern in geophysical fluid dynamics, *Phys. Rev. Research* **2**, 023143 (2020).
  - [2] X. Zhu, V. Mathai, R. J. Stevens, R. Verzicco, and D. Lohse, Transition to the ultimate regime in two-dimensional rayleigh-bénard convection, *Physical Review Letters* **120**, 10.1103/physrevlett.120.144502 (2018).
  - [3] M. Ossendrijver, The solar dynamo, *The Astronomy and astrophysics review* **11**, 287 (2003).
  - [4] D. Gubbins, The rayleigh number for convection in the earth's core, *Physics of the Earth and Planetary Interiors* **128**, 3 (2001), dynamics and Magnetic Fields of the Earth's and Planetary Interiors.