

# Marginally-Stable Thermal Equilibria of Rayleigh-Bénard Convection

Liam O'Connor<sup>1</sup>, Daniel Lecoanet<sup>1,2</sup>, and Evan H. Anders<sup>2</sup>

<sup>1</sup>*Department of Engineering Sciences and Applied Mathematics,  
Northwestern University, Evanston, IL 60208 USA and*

<sup>2</sup>*Center for Interdisciplinary Exploration and Research in Astrophysics,  
Northwestern University, Evanston, IL, 60201 USA*

Natural convection is ubiquitous throughout the physical sciences and engineering, yet many of its important properties remain elusive—particularly in the turbulent regime. In this investigation, we derive and solve a quasilinear form of the Rayleigh-Bénard problem by representing the perturbations in terms of marginally stable eigenmodes. The amplitude of each eigenmode is determined by requiring that the background state maintains marginal stability. The background temperature profile evolves due to the advective flux of each eigenmode, as well as diffusion. The entire calculation is one-dimensional, and can be run on a workstation. We find the background temperature field evolves to an equilibrium state, where the advective flux from the marginally-stable eigenmodes and the diffusive flux sum to a constant. These marginally-stable thermal equilibria (MSTE) are exact solutions of the quasilinear equations. The mean temperature profile has thinner boundary layers and larger Nusselt numbers than thermally-equilibrated 2D and 3D simulations of the full nonlinear equations. We find the Nusselt number scales like  $Nu \sim Ra^{1/3}$ . When an MSTE is used as initial conditions for a 2D simulation, we find that  $Nu$  quickly equilibrates without the burst of turbulence often induced by purely conductive initial conditions, but we also find that the kinetic energy is too large and viscously attenuates on a long viscous timescale.

## I. INTRODUCTION

Rayleigh-Bénard convection plays a foundational role in astrophysical and geophysical settings. The resulting buoyancy-driven flows regulate heat transfer and generate large-scale vortices [1]. Turbulent convection, which is associated with large Rayleigh numbers  $Ra$ , is difficult to simulate. State of the art simulations performed by [2] have reached  $Ra \sim 10^{14}$  but estimates for the sun’s convective zone and earth’s interior are  $Ra \sim 10^{16} - 10^{20}$  and  $Ra \sim 10^{20} - 10^{30}$  respectively [3, 4]. The scaling behavior of the Nusselt number  $Nu \sim Ra^\beta$  in the asymptotic ultimate regime is of particular interest. There exists a substantial body of work pertaining to this specific topic with no general consensus [5–11].

Absent solid evidence from direct numerical simulation, other methods have been developed to try to infer large- $Ra$  behavior or otherwise gain insight. In the presence of other physical effects (e.g., rotation, magnetic fields), one can sometimes derive an asymptotically consistent set of reduced equations [12, 13]. Reduced models are potentially useful in this context because they may allow us to study the problem with less expensive computations. Another approach relates to unstable exact coherent states (ECS) [14–17]. Simulations and analysis performed by [18, 19] suggest that chaotic solution trajectories might “visit” these ECS. Should that be the case, it is crucial that we discover and classify such equilibria.

Others have turned to studying quasilinear systems. The quasilinear approximation starts with a decomposition of all variables into a background and perturbations about this background. This approximation neglects the influence of nonlinear interactions between the perturbations on the perturbations themselves [20]. This renders the perturbation equations linear. Although the quasi-

linear approximation greatly simplifies the problem, an additional condition must be imposed to determine the amplitude of the perturbations. In [21], researchers compute ECS in parallel shear flows by deriving and solving a quasilinear formulation of the Navier-Stokes equations via multi-scale asymptotic arguments. They assume the background velocity evolves on a slow timescale, and to determine the perturbation amplitudes they require marginal stability at each timestep. A similar strategy is employed by [22] to studying acoustic streaming. In that work, an analytic expression for the first-order perturbation’s amplitude is found by deriving a solvability condition.

In this paper we derive and solve a quasilinear form of the Rayleigh-Bénard convection problem using a similar time-scale separation argument. In section II we recall the underlying equations, and in section III we describe how we evolve the background temperature profile while maintaining marginal stability. Section IV describes the properties of the marginally-stable thermal equilibria, in particular how the Nusselt number and characteristic wavenumbers vary with the Rayleigh number. Finally, we describe the results of simulations initialized with marginally-stable thermal equilibria in section V, and conclude in section VI.

## II. MODEL SETUP

We begin with the Boussinesq approximation for Rayleigh-Bénard Convection, nondimensionalized on the freefall timescale. The domain  $\mathcal{D}$  is 2-dimensional, rectangular, and horizontally periodic with spatial dimensions  $0 \leq x < 4$  and  $-1/2 < z < 1/2$ . The fluid of interest is constrained between two flat boundaries at

$z = -1/2$  and  $z = 1/2$  with fixed temperatures  $1/2$  and  $-1/2$  respectively. At both boundaries we specify impenetrable, no-slip conditions, such that the velocity  $\mathbf{u} = u\hat{x} + w\hat{z} = \mathbf{0}$  at  $z = \pm 1/2$ , where  $\hat{x}, \hat{z}$  are the unit vectors in the  $x$  and  $z$  directions. The equations of motion are then given by

$$\nabla \cdot \mathbf{u} = 0 \quad (1)$$

$$\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + T\hat{z} + \mathcal{R}\nabla^2 \mathbf{u} \quad (2)$$

$$\frac{\partial T}{\partial t} + \mathbf{u} \cdot \nabla T = \mathcal{P}\nabla^2 T \quad (3)$$

where  $p$  is pressure and  $T$  is temperature. For completeness, we specify a final boundary condition  $p = p_0$  at  $z = \pm 1/2$ . Any system of this form can be characterized by its dimensionless Rayleigh number  $\text{Ra} = \frac{g\alpha L^3 \Delta T}{\nu \kappa}$  and Prandtl number  $\text{Pr} = \frac{\nu}{\kappa}$ , where  $g, \alpha, L, \Delta T, \nu, \kappa$  are the gravitational acceleration, coefficient of thermal expansion, domain height, opposed temperature difference, kinematic viscosity, and thermal diffusivity respectively. In this paper, we fix  $\text{Pr} = 1$ . For convenience, we define

$$\mathcal{R} = \sqrt{\frac{\text{Pr}}{\text{Ra}}}, \quad \mathcal{P} = \frac{1}{\sqrt{\text{Pr Ra}}}. \quad (4)$$

To derive the quasilinear form, we posit that an arbitrary field  $f$  can be represented as the sum of a mean profile (denoted by  $\bar{f}$ ) and a perturbation function (denoted by  $f'$ ).

$$\mathbf{u}(x, z, t) = \mathbf{u}'(x, z, t) \quad (5)$$

$$= u'(x, z, t)\hat{x} + w'(x, z, t)\hat{z} \quad (6)$$

$$T(x, z, t) = \bar{T}(z, t) + T'(x, z, t) \quad (7)$$

$$p(x, z, t) = \bar{p}(z, t) + p'(x, z, t). \quad (8)$$

where the mean-velocity components vanish due to incompressibility and symmetry. Perturbations are defined to have no horizontal-average

$$\langle f'(x, z, t) \rangle_x \equiv \int_0^4 f'(x, z, t) dx = 0. \quad (9)$$

Substituting (7) into (3) and taking the horizontal-average reduces the system to a simple initial value problem (IVP) for  $\bar{T}$

$$\frac{\partial \bar{T}}{\partial t} + \frac{\partial}{\partial z} \langle w'T' \rangle_x = \mathcal{P} \frac{\partial^2 \bar{T}}{\partial z^2}, \quad (10)$$

with associated boundary conditions  $\bar{T}(-1/2, t) = 1/2$  and  $\bar{T}(1/2, t) = -1/2$ . It should be noted that we could obtain a similar IVP for  $u$  by breaking symmetry and considering some nontrivial mean horizontal flow  $\bar{u}(z, t)$ . However we must have  $\bar{w}(z, t) = 0$  due to incompressibility.

Substituting (8) into (2) and taking the horizontal average reveals that the mean pressure field  $\bar{p}(z)$  must satisfy

$$0 = -\frac{\partial \bar{p}}{\partial z} + \bar{T} \quad (11)$$

To solve (10) numerically, we need an expression for the perturbation so we can calculate the advective heat flux. Here we will make the quasilinear approximation, dropping the  $\mathbf{u}' \cdot \nabla \mathbf{u}'$  and  $\mathbf{u}' \cdot \nabla T'$  terms from the evolution equations for the perturbations. Substituting (5)–(8) into (1)–(3) followed by subtracting (10) and (11) from the resulting temperature and  $\hat{z}$  momentum equations gives

$$\nabla \cdot \mathbf{u}' = 0 \quad (12)$$

$$\frac{\partial \mathbf{u}'}{\partial t} = -\nabla p' + T'\hat{z} + \mathcal{R}\nabla^2 \mathbf{u}' \quad (13)$$

$$\frac{\partial T'}{\partial t} + \frac{\partial \bar{T}}{\partial z} w' = \mathcal{P}\nabla^2 T' \quad (14)$$

with Dirichlet boundary conditions

$$T'|_{z=\pm 1/2} = 0, \quad u'|_{z=\pm 1/2} = 0, \quad p'|_{z=\pm 1/2} = 0. \quad (15)$$

This is now a linear problem in  $\mathbf{u}'$  and  $T'$  which can be solved as an eigenvalue problem.

In his groundbreaking report [23], Lord Rayleigh observed that (12)–(14) can be manipulated into a separable form with generalized solutions

$$w'(x, z, t) = A \Re \left[ W(z) e^{i(k_x x - st)} \right], \quad (16)$$

$$u'(x, z, t) = A \Re \left[ U(z) e^{i(k_x x - st)} \right], \quad (17)$$

$$T'(x, z, t) = A \Re \left[ \theta(z) e^{i(k_x x - st)} \right], \quad (18)$$

$$p'(x, z, t) = A \Re \left[ P(z) e^{i(k_x x - st)} \right], \quad (19)$$

where  $A$  is the (undetermined) mode amplitude,  $s = \omega + i\sigma$  and  $k_x$  is constrained, by periodicity, to the countably infinite set (spectrum) of wavenumbers

$$k_x \in \left\{ \frac{n\pi}{2} \mid n \in \mathbb{N} \right\}. \quad (20)$$

We normalize the eigenmodes to have

$$\langle |\theta|^2 \rangle_{\mathcal{D}} = 1 \quad (21)$$

where  $\langle \cdot \rangle_{\mathcal{D}}$  denotes the spatial mean over the entire domain.

For each  $k_x$ , we can assess the stability of the perturbations by solving for the eigenvalue  $s$ , whose imaginary component  $\sigma$  plays the role of an exponential growth rate. Positive eigenvalues indicate that the system is unstable to small disturbances of wavenumber  $k_x$ , while negative eigenvalues indicate stability. A complete linear stability analysis requires solution over the full spectrum of wavenumbers. The prototypical case is used to demonstrate that the critical Rayleigh number  $\text{Ra}_c = 1708$  when  $\frac{\partial \bar{T}}{\partial z} = -1$ .

To calculate the advective heat flux in equation 10, we can sum  $\langle w'T' \rangle_x$  from each horizontal wavenumber individually. In this way, the heat flux from the perturbations influence the evolution of  $\bar{T}$ . But the evolution of  $\bar{T}$  also influences the perturbations, as equation 14 depends on  $\partial_z \bar{T}$ . Thus, the mean temperature and perturbations fields are coupled.

### III. PERTURBATION EVOLUTION

The linearized system (12)–(14) does not constrain the amplitude of the eigenmodes,  $A$ . However, the advective heat flux is proportional to  $A^2$ , so we need to specify the amplitude in order to solve equation 10. To evolve  $\bar{T}$ , we assume the perturbations evolve on a much faster timescale than the mean temperature, as in [22]. Stable modes ( $\sigma < 0$ ) decay away rapidly. Unstable modes ( $\sigma > 0$ ) will not persist on the slow timescale because the advective term  $\langle w'T' \rangle_x$  tends to stabilize  $\bar{T}$ , thereby creating a negative feedback loop. Only marginally stable modes can be maintained on the slow timescale. Therefore the amplitude  $A$  must satisfy

$$\max_{k_x} \{\sigma\} = 0. \quad (22)$$

For various  $Ra$  and fixed  $Pr = 1$ , we seek marginally-stable thermal equilibria (MSTE) satisfying  $\frac{\partial \bar{T}}{\partial t} = 0$  according to (10). We employ the *Dedalus* pseudo-spectral python framework [24] to solve the EVP outlined in Section II as well as the IVP (10). We represent each field with Chebyshev polynomials use the 3/2 dealiasing rule to calculate the advective heat flux. The necessary number of basis functions increases with  $Ra$  as the eigenfunctions include increasingly small-scale features. We use the *Eigentools* package [25] to manipulate the eigenfunctions and calculate the advective heat flux  $\langle w'T' \rangle_x$ .

At iteration 0 we construct a marginally-stable initial temperature profile  $\bar{T}(z, 0)$  whose equation is given in appendix A. At an arbitrary iteration beginning at  $t = t_0$ , we seek to evolve  $\bar{T}(z, t_0)$  into a new marginally-stable profile  $\bar{T}(z, t_0 + \Delta t)$  according to (10). To achieve this, we use a second-order, two-stage IMEX Runge-Kutta method. We initialize (10) with the previous iteration's solution to (10) and maintain marginal stability at each iteration. The eigenfunctions and amplitude are assumed to be constant over the timestep. We can solve the IVP over long periods of time by stopping regularly (at each iteration) to adjust the eigenfunctions and their amplitude  $A$ . It is essential that we pick the correct eigenfunction amplitude when calculating the advective term to maintain marginal stability. We will now illustrate our method of finding the appropriate  $A$  through an example.

An iteration is performed as follows. Consider a marginally-stable temperature profile  $\bar{T}(z, t_0)$ . By definition, its maximum eigenvalue is 0. Diffusing  $\bar{T}(z, t_0)$  tends to increase its eigenvalues while ignoring the diffusive term and evolving according to advection tends to stabilize the system.  $A$  must be selected such that these two influences are equal and opposite. We can measure the effects of diffusion and advection on the maximum eigenvalue by solving two new IVPs

$$\frac{\partial \bar{T}_{\text{diff}}}{\partial t} = \mathcal{P} \frac{\partial^2 \bar{T}_{\text{diff}}}{\partial z^2} \quad (23)$$

$$\frac{\partial \bar{T}_{\text{adv}}}{\partial t} + \frac{\partial}{\partial z} \langle w'T' \rangle_x = 0 \quad (24)$$




FIG. 1. Eigenvalue spectra for  $Ra = 10^5$ . The spectrum of an “initial” marginally-stable mean temperature profile  $\bar{T}(z, t_0)$  has a maximum eigenvalue of 0. Given a small fixed timestep  $\Delta t$ , diffusion destabilizes the system, increasing its eigenvalues. Advection tends to stabilize the system, decreasing its eigenvalues. We find the eigenfunction amplitude  $A^2$  such that the combination of diffusion and advection yields a new, “evolved,” marginally-stable mean temperature profile  $\bar{T}(z, t_0 + \Delta t)$ . In this case  $A^2 \approx 1$  due to approximately equal magnitudes of the diffused and advected eigenvalues.

where  $\bar{T}_{\text{diff}}$  and  $\bar{T}_{\text{adv}}$  denote the diffused and advected temperature profiles. We initialize these IVPs with the marginally stable temperature of the current iteration. Suppose  $k_x$  is the wavenumber of the marginally-stable mode. Then, using the diffused and advected profiles, we compute the eigenvalues  $\sigma_{\text{diff}}$  and  $\sigma_{\text{adv}}$  at  $k_x = k_x$ . The appropriate amplitude  $A^2$  can then be approximated by

$$A^2 \approx A_0^2 \equiv -\frac{\sigma_{\text{diff}}}{\sigma_{\text{adv}}} \quad (25)$$

where the square in  $A_0^2$  is due to the quadratic nonlinearity of the eigenfunctions in the advective term and the minus sign is due to the fact that  $\sigma_{\text{adv}} < 0$  and  $\sigma_{\text{diff}} > 0$  in general. We illustrate these trends in Figure 1. The subscript 0 denotes a preliminary estimate which we later refine.  $\bar{T}(z, t)$  is then evolved according to (10) and another eigenvalue solve is performed. Given a fixed timestep  $\Delta t$ , we assume the dominant eigenvalue can be described by a continuous function  $\sigma_{\text{max}}(A^2)$  which is locally differentiable. We use Newton’s method to find an amplitude which satisfies our marginal stability tolerance criterion  $|\sigma_{\text{max}}(A^2)| < 10^{-9}$ . Marginally-stable modes do not oscillate in time, i.e.  $\sigma = 0$  implies  $\omega = 0$ . This agrees with the conventional notion of exchange of stabilities [26]. Crucially, we do not assume the  $k_x$  of the marginally-stable mode is fixed. In section III A we specify procedures for the treatment of multiple simultaneously marginal modes.

### A. Treatment of Multiple Marginally-Stable Modes

In most cases, we encounter eigenvalue spectra with multiple simultaneously marginal modes. To accommodate this we generalize the advective term in (10) to accommodate  $N$  simultaneously marginal modes

$$\langle w'T' \rangle_x = \sum_{n=1}^N 2A_n^2 \Re[W_n \theta_n^*] \quad (26)$$

where  $W_n$  and  $\theta_n$  are the eigenmodes associated with  $k_x = \frac{n\pi}{2}$ . There are now  $N$  modes, each having their own amplitude to solve for and eigenvalue to keep marginally-stable. Given a small fixed time step  $\Delta t$ , let  $\mathbf{A}^2$  be the amplitude vector and  $\boldsymbol{\sigma}(\mathbf{A}^2)$  be the dependent eigenvalues. We expect a function  $\boldsymbol{\sigma} : \mathbb{R}^N \rightarrow \mathbb{R}^N$  to have isolated roots  $\tilde{\mathbf{A}}^2$  (should they exist). A preliminary approximation  $\mathbf{A}_0^2$  for the appropriate amplitude vector  $\tilde{\mathbf{A}}^2$  is given by a generalized form of (25):

$$\tilde{\mathbf{A}}^2 \approx \mathbf{A}_0^2 = -\Sigma_{\text{adv}}^{-1} \boldsymbol{\sigma}_{\text{diff}}. \quad (27)$$

Here  $\boldsymbol{\sigma}_{\text{diff}} = \boldsymbol{\sigma}(\mathbf{0})$  refers to the eigenvalues after a brief  $\Delta t$  period of diffusion. We account for the influence of  $N$  advection terms on  $N$  eigenvalues (one for each mode) by constructing an eigenvalue matrix  $\Sigma_{\text{adv}} \in \mathbb{R}^{N \times N}$ . For example, the element of  $\Sigma_{\text{adv}}$  at row  $i$  and column  $j$  is given by the  $i$ th mode's eigenvalue due to advection by the  $j$ th mode's eigenfunctions. At the point of our preliminary estimate  $\mathbf{A}_0^2$ , we approximate the Jacobian matrix

$$J = \begin{bmatrix} \nabla \sigma_1(A_1, A_2, \dots, A_N) \\ \nabla \sigma_2(A_1, A_2, \dots, A_N) \\ \vdots \\ \nabla \sigma_N(A_1, A_2, \dots, A_N) \end{bmatrix} \quad (28)$$

via first-order finite differences, requiring  $N^2$  more individual EVP solves. A once-refined estimate  $\mathbf{A}_1^2$  is then given by

$$\tilde{\mathbf{A}}^2 \approx \mathbf{A}_1^2 = -J^{-1} \boldsymbol{\sigma}(\mathbf{A}_0^2). \quad (29)$$

We then use each subsequent guess to adjust the Jacobian via Broyden's method for root-finding in multi-dimensional functions [27]. We find  $\boldsymbol{\sigma}(\mathbf{A}^2)$  does indeed have a unique root provided the time step is not too large and there are no numerical instabilities, as outlined in appendix B. Presumably this is due to the coupling of  $\langle w'T' \rangle_x$  with  $\bar{T}$ . Over the course of a large time step,  $\bar{T}$  evolves according to (10) and eventually the original eigenfunctions cease to provide a stabilizing influence. It is precisely for this reason why we must halt the IVP regularly to adjust the eigenfunctions and their amplitudes.

Difficulty arises when transitioning between different numbers of marginal modes, particularly when stable modes become marginally-stable mid-iteration. We facilitate these transitions by defining an adjustable candidate tolerance  $\varepsilon_{\text{cand}} \in [10^{-6}, 10^{-8}]$ . A mode which meets




FIG. 2. Mean temperature profiles  $\bar{T}$  for  $Ra = 10^8$ . The initial profile is given by (A1). We evolve this background temperature profile according to section III until we reach a marginally-stable thermal equilibrium (MSTE). The DNS curve is obtained from a 2D nonlinear simulation of (1)–(3) with *Dedalus*. DNS temperature data are horizontally- and time-averaged. The initial profile has the narrowest boundary layer, while the DNS profile has the widest boundary layer. The MSTE profile exhibits prominent dips, nested alongside the boundary regions.

the candidate tolerance can be included as a candidate marginal mode. Candidate modes are rejected when the root-finding algorithm converges on a negative amplitude  $A^2 < 0$ . Otherwise the mode becomes marginally-stable.

After several time steps,  $\bar{T}$  tends to asymmetricize due to numerical noise. Though asymmetric pairs of solutions may exist, we further constrain  $\bar{T}$  by setting the coefficients of its even Chebyshev basis functions to zero at the start of each iteration.

### IV. PROPERTIES OF THERMALLY EQUILIBRATED STATES

We evolve  $\bar{T}$  as described above until  $\max |\partial_t \bar{T}| < 10^{-5}$ . In this marginally-stable thermal equilibrium,  $\bar{T}$  does not evolve in time, and the perturbations also do not evolve in time, as they are marginally stable. Thus, these configurations are exact solutions to the quasilinear equations (equations 10–14). They differ from the usual ECS in that ECS are fixed points of the full nonlinear problem (1)–(3). Such definitions are not mutually exclusive, but in general we can assume that MSTE and ECS are not steady with respect to their counterparts' equations. We compute symmetric MSTE for  $Ra$  in the range  $10^5 - 10^9$ .

Figure 2 gives temperature profiles for  $Ra = 10^8$  where the initial profile, whose construction is outlined in appendix A, is employed at iteration 0. Direct numerical simulations (DNS) are performed by solving (1)–(3) with *Dedalus*, followed by horizontal- and time- averaging. The DNS curve is more diffuse than the MSTE curve, which in turn, is more diffuse than the initial curve. Per-

forming an eigenvalue solve by setting  $\bar{T}$  equal to the DNS profile yields unstable eigenvalues. This suggests that MSTE might maximize boundary layer thickness, subject to the marginal stability constraint.

The most resilient and unexpected feature of MSTE temperature profiles are the pronounced dips adjacent to the boundary layers. These dips appear in every solution, regardless of Ra. Physically, they correspond to thin layers in which the mean temperature gradient reverses, contradicting an important hypothesis of [5, 6]. This counter-diffusion, which opposes overall heat transfer, is overcome by the coinciding advective flux, shown in Figure 3. We do not understand the source of these dips, but similar temperature gradient reversals were reported by [17] along the midlines of 2D convective cellular solutions at  $Ra \sim 10^6$ . In that case, the reversals were due to nonlinear advection, which is not present in our quasilinear model.

In Figure 3, where we give heat flux profiles and eigenvalue spectra for two cases:  $Ra = 2 \times 10^5$  (top) and  $Ra = 10^9$  (bottom). For  $Ra = 2 \times 10^5$ , there is a single marginal mode at  $k_x = 1.5\pi$  whose advective flux occupies the bulk of the domain. These states have wide boundary layers which gradually subside as advection becomes the dominant flux component. Transitional regions occur over a smaller length scale for  $Ra = 10^9$  where the shift from diffusion to advection is sharp. At  $Ra = 10^9$  we find five marginally-stable modes are necessary to reach an MSTE. Thin advection profiles, belonging to high-wavenumber modes with  $k_x = 23.5\pi, 24\pi$ , hug the boundary layer. Closer to the bulk of the domain, we see wider advection profiles corresponding to modes in a second group of marginal modes  $k_x = 6\pi, 6.5\pi$ . The  $k_x = 1.5\pi$  mode forms the large-scale convective cell structures observed in DNS, again occupying the bulk of the domain. The pairs of modes  $k_x = 6\pi, 6.5\pi$  and  $k_x = 23.5\pi, 24\pi$  are each associated with a single maximum in our plots of growth rate  $\sigma$  as a function of  $k_x$  (lower right panel of Figure 3). If we allowed wavenumbers to vary continuously, there would be an unstable mode between these pairs of wavenumbers. However, since we have fixed the horizontal size of our domain, we are left with pairs of discrete margin modes.

MSTE for large Ra tend to have a diverse combination of marginal modes. In every case, the  $k_x = 1.5\pi$  mode is included. In Figure 4 we give the wavenumbers  $k_x$  of marginal modes. For some values of Ra we obtain states with pairs marginally-stable modes, with wavenumbers separated by  $\pi/2$ . Like  $k_x = 6\pi, 6.5\pi$  and  $k_x = 23.5\pi, 24\pi$  for the  $Ra = 10^9$  MSTE, these are due to the discretization of wavenumbers from our domain of width 4. We think of the pairs of modes as acting together as part of a single maximum of the growth rate as a function of the wavenumber. When wavenumbers are adjacent, we plot them in the same color and denote the larger mode with an x and the smaller with a +. For  $Ra \geq 10^6$ , a second branch of marginal modes is shown in light green. Least-squares regression gives

$\max\{k_x\} \propto Ra^{0.300}$  with  $R^2 = 0.998$  for this maximum branch. For large Ra, the advective fluxes of this maximum branch opposes the strong diffusion of the thin MSTE boundary layers. At  $Ra \geq 10^8$ , a third branch appears (shown in blue), splitting the widening gap between the other two. For these points regression gives  $k_x \propto Ra^{0.155}$  with  $R^2 = 0.988$ . The blue branch is associated with moderately wide advection profiles, filling a niche in the total flux by uniting the thin profiles of the maximum branch with those of the bulk-domain-oriented minimum branch.

The largest marginal wavenumber  $\max\{k_x\}$  (represented by the light green branch in Figure 4) serves as an inverse minimum length-scale in the  $x$  direction. The finest vertical structures in  $\bar{T}(z)$  appear near the boundaries, requiring more basis functions (resolution) at large Ra. Naturally, this provides a complementary minimum length-scale for  $z$ . We define the boundary layer width  $\delta$  as the distance from the boundary where  $\partial_z T$  equals zero, i.e.,

$$\left. \frac{\partial \bar{T}}{\partial z} \right|_{z=-\frac{1}{2}+\delta} = 0. \quad (30)$$

This height corresponds to the local extrema of the MSTE temperature profile, e.g., in Figure 2. In Figure 5, we show  $\max\{k_x\}$  is proportional to  $\delta^{-1}$  over our range of Ra. Least-squares regression gives  $\delta^{-1} = 1.71 \max\{k_x\} - 2.13$  with  $R^2 = 0.996$ . We can assume from this length scale agreement that the mean-squared  $x$  and  $z$  components of the temperature and velocity gradients are proportional

$$\begin{aligned} \left\langle \frac{\partial T}{\partial x} \cdot \frac{\partial T}{\partial x} \right\rangle_{\mathcal{D}} &\propto \left\langle \frac{\partial T}{\partial z} \cdot \frac{\partial T}{\partial z} \right\rangle_{\mathcal{D}} \\ \left\langle \frac{\partial \mathbf{u}}{\partial x} \cdot \frac{\partial \mathbf{u}}{\partial x} \right\rangle_{\mathcal{D}} &\propto \left\langle \frac{\partial \mathbf{u}}{\partial z} \cdot \frac{\partial \mathbf{u}}{\partial z} \right\rangle_{\mathcal{D}}. \end{aligned} \quad (31)$$

This is consistent with the mean squared gradient assumptions of [5].

We find the MSTE can be characterized by their boundary layer height. In Figure 6, we illustrate the scaling behavior of the boundary layer height  $\delta = Ra^{-1/3}$ . This is consistent with Malkus' classical marginal-stability theory, a scaling argument which perceives the boundary regions as subdomains which are themselves marginally-stable [5].

The Nusselt number, which measures convective performance is given by

$$Nu = \frac{\langle \langle w'T' \rangle_x - \mathcal{P} \frac{\partial \bar{T}}{\partial z} \rangle_z}{\langle -\mathcal{P} \frac{\partial \bar{T}}{\partial z} \rangle_z}. \quad (32)$$

There is no general consensus surrounding the scaling behavior of Nu for high Ra systems, which are of particular importance in astrophysical and geophysical systems. In Figure 7 we report Nu for MSTE, "steady rolls" ECS [16], and DNS [28, 29]. We find that MSTE satisfy  $Nu \sim Ra^{1/3}$ , consistent with our finding that the




FIG. 3. Heat fluxes (left) and eigenvalue spectra (right) of equilibrated states  $Ra = 2 \times 10^5$  (top) and  $Ra = 10^9$  (bottom). The heat flux profiles are symmetric about  $z = 0$ , so we only plot for  $z < 0$ . Advection profiles belong to marginally-stable modes. For low  $Ra$ , a single mode with  $k_x = 1.5\pi$  is sufficient to oppose boundary layer diffusion and facilitate heat flux throughout the bulk of the domain. For large  $Ra$ , high-wavenumber modes contribute pronounced small-scale advection profiles which tightly hug the thin boundary layers. A combination of progressively wider advection profiles is necessary to transition to the  $k_x = 1.5\pi$  mode.

boundary layer width scales like  $\delta \sim Ra^{1/3}$ . The Nusselt numbers of the ECS are somewhat lower and the DNS Nusselt numbers are yet lower still. In both cases, the  $Ra$  dependence appear slightly more shallow than for the MSTE. [16] hypothesized that the  $Nu$  of all ECS which admit classical Malkus scaling must always exceed the  $Nu$  of turbulent convection. If we generalize this notion to include quasilinear equilibria, our findings agree; MSTE have larger  $Nu$  than 2D and 3D DNS. This might be due to the chaotic transitions among the unstable periodic orbits outlined by [18, 19] inhibiting heat flux. We might also anticipate the existence of similar equilibria with smaller  $Nu$ , occupying complementary nodes in the Markov chain whose behavior agrees with DNS.

## V. SIMULATIONS WITH THERMALLY EQUILIBRATED INITIAL CONDITIONS

This investigation is partially motivated by the prospect of decreasing DNS runtimes by employing MSTE as initial conditions. One common choice of initial

conditions for DNS of equations (1)–(3) are

$$\begin{aligned} T(x, z)|_{t=0} &= 0.5 - z + N \\ \mathbf{u}(x, z)|_{t=0} &= \mathbf{0} \\ p(x, z)|_{t=0} &= 0 \end{aligned} \quad (33)$$

where  $N$  is low-amplitude random noise. Here we instead initialize using the MSTE,

$$\begin{aligned} T(x, z)|_{t=0} &= \bar{T}(z) + \sum_{n=1}^N A_n \Re[\theta_n(z) e^{ik_{x_n} x}] + N \\ \mathbf{u}(x, z)|_{t=0} &= \sum_{n=1}^N A_n \Re[(U_n(z) \hat{x} + W_n(z) \hat{z}) e^{ik_{x_n} x}] \\ p(x, z)|_{t=0} &= \sum_{n=1}^N A_n \Re[P_n(z) e^{ik_{x_n} x}] \end{aligned} \quad (34)$$

where  $\theta_n(z), U_n(z), W_n(z), P_n(z); A_n$ ; and  $k_{x_n}$  refer to the complex eigenfunctions, amplitude, and wavenumber at the  $n$ th marginal mode respectively. Note that although the MSTE is an equilibrium of the quasilinear equations, it is not an equilibrium of the full nonlinear






FIG. 4. Wavenumbers of marginally-stable modes in thermally equilibrated states. Marginal modes often appear in adjacent pairs, which we denote with a common color. Within each of these pairs, the largest and smallest  $k_x$  are denoted with an + and x respectively. For example, the spectrum corresponding to  $Ra = 10^5$  has adjacent marginal wavenumbers  $k_x = \pi, 1.5\pi$ . The  $Ra = 10^9$  spectrum, shown in lower right corner of Figure 3, has three groups of maxima, with a single marginal mode in the first group ( $k_x = 1.5\pi$ ), two adjacent marginal modes in the second group ( $k_x = 6\pi, 6.5\pi$ ), and two adjacent marginal modes in the third group ( $k_x = 23.5\pi, 24\pi$ ). The largest wavenumbers of the green branch obey a power-law relationship  $\max\{k_x\} \propto Ra^{0.300}$  with  $R^2 = 0.998$ .




FIG. 5. For  $Ra \geq 10^6$ , the maximum marginally-stable wavenumber (corresponding to the green x markers in Figure 4) are inversely related to the boundary layer height  $\delta$ .  $(\max\{k_x\})^{-1}$  gives a minimum  $x$  length scale for the perturbations, and consequently, the advection. For large  $Ra$ , the boundary layers admit small scale features, requiring more vertical basis functions (higher resolution). The boundary layer width gives an estimate of the minimum vertical length-scale in the problem. This suggests that the minimum horizontal and vertical lengthscales are proportional to each other over a wide range of  $Ra$ .




FIG. 6. Boundary layer height  $\delta$  of MSTE. We define the boundary layer height based off the location where  $\frac{\partial T}{\partial z} = 0$  (see equation 30). Plotting on a log-log scale, we find that  $\delta$  and  $Ra$  obey a power-law relationship. We also demonstrate that  $Ra^{1/3}\delta$  is approximately constant with respect to  $Ra$  which is consistent with [5]

equations. Accordingly the simulation state would evolve on initialization absent a random noise term. Here we include noise as a source of asymmetries.

Simulations initialized with the conductive equilibrium plus low-amplitude thermal noise have a large peak in  $Nu$  early on in their evolution (Figure 8). This is due to a burst of turbulence which occurs when the convective motions first become nonlinear. A simulation initialized with the MSTE, however, does not exhibit this transient burst of turbulence, as the large-scale anatomy of convective cells exists on initialization. Simulation of this transitional period is prohibitive [30]. For high  $Ra$  experiments, researchers often “bootstrap” data by initializing simulations with the results of similar  $Ra$  runs [28, 31]. MSTE can be perceived as a set of initial conditions, designed for avoiding the simulation of transient high Reynolds number flows.

MSTE are laminar, lacking the small-scale structures associated with moderate to high  $Ra$  experiments. This is an apparent consequence of the quasilinear assumptions. If we perceive MSTE as background states, DNS suggest that plumes, vortex sheets, and other unstable turbulent features inhibit total heat transfer. This perspective agrees with conventional models of transitions to turbulent flows, such as Boussinesq’s turbulent-viscosity hypothesis [32]. The emergence of small-scale velocity structures tends to increase total shear [26, 33, 34], thereby impeding buoyancy-driven flows and decreasing advection in the bulk of the domain. We could also attribute the diffuse DNS temperature profile in Figure 2 with unsteady boundary-layer penetration and mixing that MSTE do not exhibit.

We also find that the average kinetic energies of simulations initialized with MSTE are significantly larger than those found in simulations initialized with the conductive state plus noise, as shown Figure 9. This is because the




FIG. 7. Nusselt numbers are shown for MSTE, aspect-ratio-optimized “steady rolls” ECS [16], as well as statistically-steady 2D and 3D DNS [28, 29]. All datasets obey power-law relationships, with the MSTE and ECS scaling like  $Nu \sim Ra^{1/3}$ . MSTE have greater  $Nu$  than the ECS, which in turn, have greater  $Nu$  than the DNS. This can be explained by the contrasting boundary layer geometries shown in Figure 2.




FIG. 8. Nusselt numbers of simulations performed at  $Ra = 10^8$ , initialized with the conductive equilibrium plus thermal noise (black) and the MSTE (blue). MSTE simulations do not undergo a convective-transient period because the characteristic large-scale convective cell structure exists on initialization.

MSTE contain strong large-scale flows, which decay on a viscous time-scale

$$t_\nu \sim \sqrt{\frac{Ra}{Pr}}.$$

Consequently, MSTE initial conditions do not reduce the simulation time required to achieve a statistically-steady state—rather they increase it considerably! This suggests the MSTE background state perspective is partially flawed, as a more useful background state would approximate the kinetic energy with more fidelity.




FIG. 9. Average kinetic energies are reported for the same simulations illustrated in Figure 8 ( $Ra = 10^8$ ). The eigenfunctions belonging to MSTE have significantly more kinetic energy than the statistically-steady state. Kinetic equilibrium is achieved on the viscous time-scale  $t_\nu \sim \sqrt{Ra/Pr}$ .

## VI. DISCUSSION

In this paper we describe a new way to study Rayleigh–Bénard convection. We compute marginally-stable thermal equilibria (MSTE), which are equilibria of the quasilinear equations. To compute MSTE, we construct a marginally-stable mean temperature profile and evolve it according to the advective flux of its marginally-stable eigenfunctions, and its own diffusion. We assume that at least some modes are always in a marginally-stable configuration. The marginal stability constraint then fixes the ratio between advection and



diffusion (eigenfunction amplitude  $A^2$ ). We use standard root-finding algorithms to solve for the appropriate  $A^2$  at each iteration, until, the fixed combination of diffusion and advection sum to a constant flux. The MSTE calculation is a one-dimensional problem, combining eigenvalue solves, and the time evolution of the one-dimensional mean temperature profile. Thus, they can be calculated on a single workstation.

The MSTE retain several key features that are prevalent in experiments and simulations:  $Nu \sim Ra^{1/3}$  scaling, large-scale convective cell structures, and minimum length scale agreement. They also exhibit unique and unexpected features: mean temperature gradient-reversals/dips, high kinetic energy flows, and a larger  $Nu$  than other time-invariant solutions. When initializing with different mean temperature profiles, we find the same MSTE, suggesting these equilibria might be unique.

Simulations initialized with the MSTE (34) do not undergo an early convective transient period, but have faster flows when compared with DNS. From a dynamical systems perspective, unstable orbits depart from MSTE and approach the global attractor on a viscous time-scale. This requires more computational effort to achieve relaxation when compared to the conventional conductive initial condition (33).

Using the mean temperature in a statistically-steady DNS as a background state for an eigenvalue problem yields positive eigenvalues: the system is in a perpetual state of instability. Unstable modes tend to stabilize the system rapidly, creating a negative feedback loop whose average state is linearly unstable. We might curtail the disagreement between MSTE and DNS by adjusting our marginal-stability criterion. Should the fast and slow time scales not be entirely separate, we might anticipate the long-term persistence of moderately unstable modes. To find MSTE, we initialize the time-evolution algorithm with the analytic temperature profile derived by [35]. This involves modifying the boundary layer thickness  $\delta_0$  to achieve marginal stability.

Instead of imposing marginal stability, we can use quasilinear model to thermally-equilibrate and expand 1D approximations into 2D. As previously noted, there is no shortage of theories pertaining to the scaling behavior  $Nu \sim Ra^\beta$  [5–11]. For large  $Ra$ , we can approximate  $\delta_0 \approx Nu^{-1}$  and construct a 1D temperature profile according to (A1) or some other approximation. In this way, other 2D quasilinear thermal equilibria can be obtained and analyzed.

## ACKNOWLEDGMENTS

The authors thank Geoff Vasil, Greg Chini, and Emma Kaufman for their valuable feedback and suggestions. We also thank the **Dedalus** and **Eigentools** development teams. Computations were conducted with support by the NASA High End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center on Pleiades with allocation GIDs s2276.

## Appendix A: Initial Buoyancy Profile

We initialize the thermal-equilibration algorithm with an analytical thermal boundary layer equation, derived by [35]

$$\begin{aligned} \bar{\tau}_0(\xi) &= \frac{\sqrt{3}}{4\pi} \log \frac{(1 + a\xi)^3}{1 + (a\xi)^3} + \frac{3}{2\pi} \arctan \left( \frac{4\pi}{9} \xi - \frac{1}{\sqrt{3}} \right) + \frac{1}{4} \\ \xi &= \frac{z + 1/2}{\delta_0} \geq 0, \quad a = \frac{2\pi}{3\sqrt{3}} \end{aligned} \quad (A1)$$

where  $\delta_0$  is the boundary layer height. This function is meant to describe the temperature near  $z = -1/2$ . It does not pass through the origin and it is undefined for  $z > 0$ . An appropriate initial mean temperature profile  $\bar{T}_0$  must be odd-symmetric, i.e.  $\bar{T}_0(-z) = -\bar{T}_0(z)$ . Due to continuity, this implies  $\bar{T}_0(0) = 0$ . Accordingly, we construct  $\bar{T}_0(z)$  by translating  $\bar{\tau}_0(z)$  vertically to pass through the origin. We then take its odd-extension and include a unique scaling coefficient to satisfy the boundary conditions  $\bar{T}_0(-1/2) = 1/2$  and  $\bar{T}_0(1/2) = -1/2$ . The initial mean temperature profile is therefore given by

$$\bar{T}_0(z) = \frac{1}{2} \begin{cases} 1 - \frac{\bar{\tau}_0\left(\frac{z+1/2}{\delta_0}\right)}{\bar{\tau}_0\left(\frac{1}{2\delta_0}\right)} & -1/2 \leq z \leq 0 \\ -1 + \frac{\bar{\tau}_0\left(\frac{1/2-z}{\delta_0}\right)}{\bar{\tau}_0\left(\frac{1}{2\delta_0}\right)} & 0 < z \leq 1/2. \end{cases}$$

We expect each  $Ra$  to be associated with a unique  $\delta_0$  for which  $\bar{T}_0(z)$  is marginally-stable. It should be noted that when experimenting with various initial profiles (tanh, erf, etc.), we obtain indistinguishable equilibrated states. Therefore these initial states lie in the MSTE basin of attraction. This might also suggest that solutions are unique. An example of (A1) is given by the blue curve in Figure 2.

## Appendix B: Timestep Management

Timesteps don't manage themselves! ■

### Appendix C: MSTE Metrics

Ra	$N_z$	Nu	$\delta_0$	$k_x$ marginals
1.00E+05	256	5.9343	0.15746	$1\pi, 1.5\pi$
2.00E+05	256	7.45986	0.12341	$1.5\pi$
4.00E+05	256	8.92597	0.10395	$1.5\pi$
1.00E+06	256	11.6689	0.07922	$1.5\pi, 2.5\pi, 3\pi$
2.00E+06	256	14.8463	0.06345	$1.5\pi, 3.5\pi$
4.00E+06	256	18.9433	0.04933	$1.5\pi, 4.5\pi$
1.00E+07	256	25.6821	0.03632	$1.5\pi, 5.5\pi$
2.00E+07	256	32.4531	0.02820	$1.5\pi, 6.5\pi, 7\pi$
4.00E+07	512	40.7925	0.02289	$1.5\pi, 8\pi, 8.5\pi$
1.00E+08	512	55.4383	0.01690	$1.5\pi, 4.5\pi, 10.5\pi, 11\pi$
2.00E+08	512	69.8349	0.01318	$1.5\pi, 5\pi, 13.5\pi, 14\pi$
4.00E+08	512	87.8525	0.01053	$1.5\pi, 5.5\pi, 17.5\pi$
1.00E+09	768	119.318	0.00817	$1.5\pi, 6.0\pi, 6.5\pi, 23.5\pi, 24\pi$

TABLE I: Control parameters are given for the timestepping algorithm.  $N_z$  denotes the number of Chebyshev basis functions employed. The remaining quantities are derived directly from MSTE solutions.

- 
- [1] L.-A. Couston, D. Lecoanet, B. Favier, and M. Le Bars, *Phys. Rev. Research* **2**, 023143 (2020).
- [2] X. Zhu, V. Mathai, R. J. Stevens, R. Verzicco, and D. Lohse, *Physical Review Letters* **120** (2018), 10.1103/physrevlett.120.144502.
- [3] M. Ossendrijver, *Astronomy & Astrophysics Reviews* **11**, 287 (2003).
- [4] D. Gubbins, *Physics of the Earth and Planetary Interiors* **128**, 3 (2001), dynamics and Magnetic Fields of the Earth's and Planetary Interiors.
- [5] W. V. R. Malkus, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **225**, 196 (1954).
- [6] L. N. Howard, in *Applied Mechanics*, edited by H. Görtler (Springer Berlin Heidelberg, Berlin, Heidelberg, 1966) pp. 1109–1115.
- [7] R. H. Kraichnan, *Physics of Fluids* **5**, 1374 (1962).
- [8] E. A. Spiegel, *Journal of Geophysical Research* **67**, 3063 (1962).
- [9] B. Castaing, G. Gunaratne, L. Kadanoff, A. Libchaber, and F. Heslot, *Journal of Fluid Mechanics* **204**, 1 (1989).
- [10] S. Grossmann and D. Lohse, *Journal of Fluid Mechanics* **407**, 27–56 (2000).
- [11] G. Ahlers, S. Grossmann, and D. Lohse, *Rev. Mod. Phys.* **81**, 503 (2009).
- [12] K. Julien and E. Knobloch, *Journal of Mathematical Physics* **48**, 065405 (2007).
- [13] K. Julien, E. Knobloch, A. M. Rubio, and G. M. Vasil, *Phys. Rev. Lett.* **109**, 254503 (2012).
- [14] F. Waleffe, A. Boonkasame, and L. Smith, *Physics of Fluids* **27**, 051702 (2015).
- [15] D. Sondak, L. M. Smith, and F. Waleffe, *Journal of Fluid Mechanics* **784**, 565 (2015), arXiv:1507.03151 [physics.flu-dyn].
- [16] B. Wen, D. Goluskin, and C. Doering, (2020).
- [17] G. P. Chini and S. Cox, *Physics of Fluids* **21**, 083603 (2009).
- [18] G. Yalniz, B. Hof, and N. Burak Budanur, arXiv e-prints, arXiv:2007.02584 (2020), arXiv:2007.02584 [physics.flu-dyn].
- [19] P. Cvitanović and J. Gibson, *Physica Scripta* **2010**, 014007 (2010).
- [20] J. B. Marston, G. P. Chini, and S. M. Tobias, *Phys. Rev. Lett.* **116**, 214501 (2016), arXiv:1601.06720 [physics.flu-dyn].
- [21] C. Beaume, G. P. Chini, K. Julien, and E. Knobloch, *Physical Review E* **91** (2015), 10.1103/phys-rev.91.043010.
- [22] G. Michel and G. P. Chini, *Journal of Fluid Mechanics* **858**, 536–564 (2019).
- [23] L. R. O. F.R.S., *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **32**, 529 (1916).
- [24] K. J. Burns, G. M. Vasil, J. S. Oishi, D. Lecoanet, and B. P. Brown, *Physical Review Research* **2**, 023068 (2020), arXiv:1905.10388 [astro-ph.IM].
- [25] J. Oishi, K. Burns, S. Clark, E. Anders, B. Brown, G. Vasil, and D. Lecoanet, “Eigentools: Tools for studying linear eigenvalue problems,” (2021), ascl:2101.017.
- [26] P. G. Drazin and W. H. Reid, *Hydrodynamic Stability*, 2nd ed., Cambridge Mathematical Library (Cambridge University Press, 2004).
- [27] C. Broyden, *Mathematics of Computation* **19**, 577 (1965).
- [28] H. Johnston and C. R. Doering, *Phys. Rev. Lett.* **102**, 064501 (2009).
- [29] J. D. Scheel and J. Schumacher, *Journal of Fluid Mechanics* **802**, 147–173 (2016).
- [30] E. H. Anders, B. P. Brown, and J. S. Oishi, *Phys. Rev. Fluids* **3**, 083502 (2018).

- [31] R. Verzicco and R. Camussi, *Physics of Fluids* **9**, 1287 (1997), <https://doi.org/10.1063/1.869244>.
- [32] J. Boussinesq, (1877).
- [33] D. Lecoanet, M. McCourt, E. Quataert, K. Burns, G. Vasil, J. Oishi, B. Brown, J. Stone, and R. O’Leary, *Monthly Notices of the Royal Astronomical Society* **455**, 4274 (2016), publisher Copyright: © 2015 The Authors.
- [34] S. B. Pope, *Turbulent Flows* (Cambridge University Press, 2000).
- [35] O. Shishkina, S. Horn, S. Wagner, and E. S. C. Ching, *Phys. Rev. Lett.* **114**, 114302 (2015).