

Light Curve Simulation and Shape Inversion for Human-Made Space Objects

Liam Robinson

Purdue Space Information Dynamics Group

11/13/2023

Outline

- 1 Introduction
- 2 Light Curve Simulation
- 3 Direct Shape Inversion
- 4 Discussion
- 5 Conclusion

Research Objectives

- ▶ Developing a high-fidelity light curve simulation framework to simulate the physics of real measurements
- ▶ Adapting direct shape inversion techniques for human-made space objects
- ▶ Designing a new method for inverting shape with noisy measurements and estimating nonconvex geometry

State of The Art: Simulation

- ▶ Light curve simulation literature often assumes:
 - Uniform material properties
 - Self-shadowing is negligible
 - Object geometry is simple [1]
 - Measurement noise is negligible or Gaussian [2]


State of The Art: Shape Inversion

- ▶ Direct shape inversion
 - Require *a priori* knowledge of material properties and attitude
 - Highest fidelity shape estimates
- ▶ Filter-based methods
 - Attempt to estimate shape and attitude simultaneously
 - Limited to simple geometries
- ▶ Deep learning
 - Trains models to classify objects by their light curves
 - Unpredictable behavior outside the training set


This work presents improvements to direct shape inversion

Direct Inversion Literature Timeline

Asteroid Inversion



Human-made Object Inversion



Outline

- 1 Introduction
- 2 Light Curve Simulation
- 3 Direct Shape Inversion
- 4 Discussion
- 5 Conclusion

Attitude Motion

- ▶ Environmental torques do exist on orbit, but can be neglected on the scale of minutes to hours
- ▶ Torque-free attitude motion for rigid bodies is described by Euler's equations of motion for an inertia tensor J and body frame angular velocity ω :

$$\dot{\omega} = J^{-1} [(J\omega) \times \omega]$$

- ▶ These EOMs are integrated with a quaternion $\mathbf{q} = [q_1, q_2, q_3, q_4]^T$ to track the orientation of the body in inertial space:

$$\begin{bmatrix} \dot{\epsilon}_1 \\ \dot{\epsilon}_2 \\ \dot{\epsilon}_3 \\ \dot{\epsilon}_4 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} \epsilon_4 & -\epsilon_3 & \epsilon_2 & \epsilon_1 \\ \epsilon_3 & \epsilon_4 & -\epsilon_1 & \epsilon_2 \\ -\epsilon_2 & \epsilon_1 & \epsilon_4 & \epsilon_3 \\ -\epsilon_1 & -\epsilon_2 & -\epsilon_3 & \epsilon_4 \end{bmatrix} \begin{bmatrix} \omega_1 \\ \omega_2 \\ \omega_3 \\ 0 \end{bmatrix}.$$

Types of Torque-Free Motion

Bidirectional Reflectance Distribution Functions (BRDFs)

- ▶ BRDFs describe how light is reflected from a surface
 - $f_r(L, O)$ describes the fraction of incident light L reflected in the direction of an observer O
- ▶ Many formulations exist, but to be relevant to this work, they must [20]:
 - Conserve energy for all L :

$$\int_{O \in \mathbb{S}^2} f_r(L, O) d\mathbb{S}^2 \leq 1$$

- Be nonnegative: $f_r(L, O) \geq 0$ for all L and O
 - Be reciprocal: $f_r(L, O) = f_r(O, L)$ for all L and O
- ▶ Popular relevant BRDFs include:
 - Lambertian [20]
 - Phong [21]
 - Cook-Torrance [22]

BRDFs in Action

DIFFUSE



OREN-NAYAR



PHONG



BLINN-PHONG



ASHIKHMIN-SHIRLEY



COOK-TORRANCE



GLOSSY



Figure: BRDFs implemented for this work

Outline

- 1 Introduction
- 2 Light Curve Simulation
- 3 Direct Shape Inversion
- 4 Discussion
- 5 Conclusion

Outline

- 1 Introduction
- 2 Light Curve Simulation
- 3 Direct Shape Inversion
- 4 Discussion
- 5 Conclusion

Outline

- 1 Introduction
- 2 Light Curve Simulation
- 3 Direct Shape Inversion
- 4 Discussion
- 5 Conclusion

The Earth

center



EGI Optimization

Equation

- [1] David Vallverdú Cabrera, Jens Utzmann, and Roger Förstner.
Inversion of the shape of space debris from non-resolved optical measurements within spook.
In Proceedings of the 22nd Advanced Maui Optical and Space Surveillance Technologies Conference, pages 1–18, 2021.
- [2] Siwei Fan.
The Light Curve Simulation and Its Inversion Problem for Human-Made Space Objects.
PhD thesis, Purdue University, August 2020.
- [3] Henry Norris Russell.
On the light-variations of asteroids and satellites.
The Astrophysical Journal, 24(1):1–18, 1906.
- [4] M. Kaasalainen and J. Torppa.
Optimization methods for asteroid lightcurve inversion: I. shape determination.
Icarus, 153(1):24–36, 2001.
- [5] Mikko Kaasalainen, Johanna Torppa, and Karri Muinonen.
Optimization methods for asteroid lightcurve inversion: II. the complete inverse problem.
Icarus, 153(1):37–51, 2001.
- [6] M Kaasalainen, J Torppa, and J Piironen.
Models of twenty asteroids from photometric data.
Icarus, 159(2):369–395, 2002.

- [7] J Durech and M Kaasalainen.
Photometric signatures of highly nonconvex and binary asteroids.
Astronomy & Astrophysics, 404(2):709–714, 06 2003.
- [8] LiangLiang Yu, Jianghui Ji, and Su Wang.
Shape, thermal and surface properties determination of a candidate spacecraft target asteroid (175706) 1996 fg3.
Monthly Notices of the Royal Astronomical Society, 439(4):3357–3370, 2014.
- [9] Chee-Kheng Chng, Michele Sasdelli, and Tat-Jun Chin.
Globally optimal shape and spin pole determination with light-curve inversion.
Monthly Notices of the Royal Astronomical Society, 513(1):311–332, 01 2022.
- [10] Katsushi Ikeuchi.
Recognition of 3-d objects using the extended gaussian image.
In *IJCAI*, pages 595–600, 1981.
- [11] James J. Little.
Extended gaussian images, mixed volumes, shape reconstruction.
In *Proceedings of the First Annual Symposium on Computational Geometry*, SCG '85, pages 15–23, New York, NY, USA, 1985. Association for Computing Machinery.
- [12] Richard Gardner and Peyman Milanfar.
Reconstruction of convex bodies from brightness functions.
Discrete and Computational Geometry, 29:279–303, 07 2003.

- [13] Doyle Hall, Brandoch Calef, Keith Knox, Mark Bolden, and Paul Kervin.
Separating attitude and shape effects for non-resolved objects.
In *The 2007 AMOS Technical Conference Proceedings*, pages 464–475. Maui
Economic Development Board, Inc. Kihei, Maui, HI, 2007.
- [14] Ben K. Bradley and Penina Axelrad.
Lightcurve inversion for shape estimation of geo objects from space-based
sensors.
In *Proceedings of the International Symposium on Space Flight Dynamics*, pages
1–20, 2014.
- [15] Siwei Fan and Carolin Frueh.
A direct light curve inversion scheme in the presence of measurement noise.
The Journal of the Astronautical Sciences, 67, 08 2019.
- [16] Alex M Friedman.
Observability Analysis for Space Situational Awareness.
PhD thesis, Purdue University, Apr 2020.
- [17] S Fan and C Frueh.
Multi-hypothesis light curve inversion scheme for convex objects with minimal
observations.
In *Proceedings of the 8th European Conference on Space Debris*, pages 1–7. ESA
Space Debris Office, 2021.
- [18] Liam Robinson and Carolin Frueh.

Light curve inversion for reliable shape reconstruction of human-made space objects.

In *Proceedings of the 32nd AIAA/AAS Astrodynamics Specialist Conference*, pages 1–19, 09 2022.

- [19] Alex M. Friedman and Carolin Frueh.
Observability of light curve inversion for shape and feature determination exemplified by a case analysis.
Journal of the Astronautical Sciences, 69(2):537–569, April 2022.
- [20] Rosana Montes Soldado and Carlos Ureña Almagro.
An overview of brdf models.
2012.
- [21] Bui Tuong Phong.
Illumination for Computer Generated Pictures, pages 95–101.
Association for Computing Machinery, New York, NY, USA, 1998.
- [22] R. L. Cook and K. E. Torrance.
A reflectance model for computer graphics.
ACM Trans. Graph., 1(1):7–24, jan 1982.