

# ADA HW10

*Liangquan Zhou lz2377*

*November 13, 2014*

Consider the Mayo Clinic Lung Cancer Data in R package survival: `data(lung)` or `data(cancer)`: including the variables

inst: Institution code

time: Survival time in

status: censoring status 1=censored, 2=dead

age: Age in years

sex: Male=1 Female=2, etc.


```
library(survival)
dat = lung
```

**1. Estimate and plot the survival curves for time BY sex using the following methods:**

**Kaplan-Meier**

```
fit1 <- survfit(Surv(time, status) ~ sex, data=dat)
plot(fit1, lty=2:3, main = "Kaplan-Meier Estimates for Lung Cancer Data")
legend("bottomleft", c("male", "female"), lty = c(2,3))
```


## Kaplan–Meier Estimates for Lung Cancer Data



### Fleming-Harrington

```
fit2 <- survfit(Surv(time, status) ~ sex, type = "fh2", data=dat)
plot(fit2, lty=2:3, main = "Fleming-Harrington Estimates for Lung Cancer Data")
legend("bottomleft", c("male", "female"), lty = c(2,3))
```

## Fleming-Harrington Estimates for Lung Cancer Data



2. For each case in 1, estimate the median survival time, using the estimated survival curves.

```
print(fit1)
```

```
## Call: survfit(formula = Surv(time, status) ~ sex, data = dat)
##
##      records n.max n.start events median 0.95LCL 0.95UCL
## sex=1      138   138   138   112   270     212     310
## sex=2       90    90    90    53   426     348     550
```

```
print(fit2)
```

```
## Call: survfit(formula = Surv(time, status) ~ sex, data = dat, type = "fh2")
##
##      records n.max n.start events median 0.95LCL 0.95UCL
## sex=1      138   138   138   112   270     218     320
## sex=2       90    90    90    53   426     348     641
```

We can see that the the median survival time for male and female using Kplan-Meier Estimates are 270 and 426;

The median survival time for male and female using Fleming-Harrington Estimates are also 270 and 426;