



Review Article

Hydrogen peroxide sensing, signaling and regulation of transcription factors



H. Susana Marinho^a, Carla Real^a, Luísa Cyrne^a, Helena Soares^{a,b,c}, Fernando Antunes^{a,*}

^aDepartamento de Química e Bioquímica, Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

^bInstituto Gulbenkian de Ciência, Oeiras, Portugal

^cEscola Superior de Tecnologia da Saúde de Lisboa, IPL, Lisboa, Portugal

ARTICLE INFO

Article history:

Received 11 January 2014

Received in revised form 19 February 2014

Accepted 21 February 2014

Keywords:

Redox signaling

Localized H₂O₂ concentrations

Rate constants

Thiol reactivity

Cytosol–nuclear traffic

DNA binding and transactivation

ABSTRACT

The regulatory mechanisms by which hydrogen peroxide (H₂O₂) modulates the activity of transcription factors in bacteria (OxyR and PerR), lower eukaryotes (Yap1, Maf1, Hsf1 and Msn2/4) and mammalian cells (AP-1, NRF2, CREB, HSF1, HIF-1, TP53, NF-κB, NOTCH, SP1 and CREB-1) are reviewed. The complexity of regulatory networks increases throughout the phylogenetic tree, reaching a high level of complexity in mammals. Multiple H₂O₂ sensors and pathways are triggered converging in the regulation of transcription factors at several levels: (i) synthesis of the transcription factor by upregulating transcription or increasing both mRNA stability and translation; (ii) stability of the transcription factor by decreasing its association with the ubiquitin E3 ligase complex or by inhibiting this complex; (iii) cytoplasm–nuclear traffic by exposing/masking nuclear localization signals, or by releasing the transcription factor from partners or from membrane anchors; and (iv) DNA binding and nuclear transactivation by modulating transcription factor affinity towards DNA, co-activators or repressors, and by targeting specific regions of chromatin to activate individual genes. We also discuss how H₂O₂ biological specificity results from diverse thiol protein sensors, with different reactivity of their sulfhydryl groups towards H₂O₂, being activated by different concentrations and times of exposure to H₂O₂. The specific regulation of local H₂O₂ concentrations is also crucial and results from H₂O₂ localized production and removal controlled by signals. Finally, we formulate equations to extract from typical experiments quantitative data concerning H₂O₂ reactivity with sensor molecules. Rate constants of 140 M^{−1} s^{−1} and $\geq 1.3 \times 10^3$ M^{−1} s^{−1} were estimated, respectively, for the reaction of H₂O₂ with KEAP1 and with an unknown target that mediates NRF2 protein synthesis. In conclusion, the multitude of H₂O₂ targets and mechanisms provides an opportunity for highly specific effects on gene regulation that depend on the cell type and on signals received from the cellular microenvironment.

© 2014 The Authors. Published by Elsevier B.V.
This is an open access article under the CC BY license
(<http://creativecommons.org/licenses/by/3.0/>).

Introduction

Hydrogen peroxide (H₂O₂) is a ubiquitous oxidant present in all aerobic organisms. Since its first identification in a living cell, H₂O₂ was considered a toxic byproduct of aerobic metabolism, something that cells had to remove [1]. If H₂O₂ detoxification catalyzed by catalases and peroxidases was not adequate, H₂O₂ would diffuse and oxidize biological targets causing cellular malfunctions responsible for several pathologies and aging. Favoring this paradigm was the discovery that neutrophils use H₂O₂ toxicity and produce massive amounts of H₂O₂ during the oxidative burst to kill invading pathogens. In the

70s some isolated observations already supported a role for H₂O₂ as a signaling molecule, e.g. H₂O₂ was found to mimic insulin action [2] or to activate guanylate cyclase [3]. Apparently, these observations remained mostly unnoticed in the field of oxidative stress, but at the end of the 80s some key discoveries built up on them. In 1987, it was found that H₂O₂ at micromolar levels elicits arterial pulmonary relaxation mediated by the activation of guanylate cyclase [4] and in 1989, H₂O₂ was found to potentiate tyrosine phosphorylation during insulin signaling [5] and to stimulate cell proliferation at low concentrations [6]. Also in 1989, OxyR was identified as the transcription factor (TF) targeted by H₂O₂ in the adaptive response of *Escherichia coli* (*E. coli*) and *Salmonella typhimurium* (*S. typhimurium*) [7], and in 1990 NF-κB was identified as a redox regulated TF [8]. In the following year, the activation of NF-κB by H₂O₂ was discovered in a publication [9] that had a profound impact in the field, with near 3500 citations so far. Also in 1991, NADPH oxidases were identified in non-phagocytic cells as H₂O₂ producing systems [10,11]. If H₂O₂ was

Abbreviations: AD, activation domain; ER, endoplasmic reticulum; GPx, glutathione peroxidases; NES, nuclear exporting signal; NLS, nuclear localization signal; PHD, prolyl hydroxylase; Prxs, peroxiredoxins; TF, transcription factor; Ub, Ubiquitin.

* Corresponding author.

E-mail address: fantunes@fc.ul.pt (F. Antunes).

2213-2317/\$ - see front matter © 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

<http://dx.doi.org/10.1016/j.redox.2014.02.006>

a toxic species, why were cells intentionally producing this species by a complex regulated mechanism? Concomitantly, several clinical trials based on the notion that oxidants were toxic and antioxidants were beneficial for cancer prevention were largely unsuccessful as reviewed in [12]. Nowadays, redox biology is an established field and the essential regulating role played by H_2O_2 *in vivo* with important implications in health and disease is unquestionable. However, there are still a lot of unanswered questions regarding our understanding of redox-dependent regulation of gene expression. What makes a good H_2O_2 sensor? What are the common chemical and kinetic principles that govern H_2O_2 signaling? Is it possible to obtain an integrative view of H_2O_2 regulation of TFs?

In this review, we will start by discussing what characteristics an H_2O_2 sensor should have; we review the chemistry of H_2O_2 , mainly its reaction with thiols. The aim is to give a brief overview of basic chemical and kinetic principles that govern H_2O_2 signaling. Next, we describe briefly the TFs reviewed here, which include bacterial (OxyR and PerR), yeast (Yap1, Msn2/4, Maf1, and Hsf1), and mammalian (AP-1, NRF2, CREB, TP53, NOTCH, NF- κ B, SP1, HIF-1, SREBP-1 and HSF1) TFs. The main body of this article describes the redox regulation of these TFs by H_2O_2 . A detailed review on each of the TFs listed is not intended, as there are many excellent reviews that do so. We aim to give an integrative review of their regulation by H_2O_2 at several steps: synthesis and stability of the TF, cytoplasm-nuclear trafficking and DNA binding and transactivation, so that the reader is made aware of the diversity of mechanisms by which H_2O_2 regulates TFs and also what the common themes in H_2O_2 -regulated signaling pathways are.

What makes a good sensor for H_2O_2 ?

The characteristics of a good sensing molecule for H_2O_2 can be derived from basic concepts taken from information theory and chemistry. Low-molecular weight thiols react slowly with H_2O_2 , as exemplified by the rate constants for H_2O_2 reaction with cysteine and reduced glutathione (GSH), which are respectively $2.9\text{ M}^{-1}\text{ s}^{-1}$ and $0.87\text{ M}^{-1}\text{ s}^{-1}$ (pH 7.4, see Table 1). The reaction of thiols with H_2O_2 involves a nucleophilic attack of the thiolate on H_2O_2 and, as such, thiol reactivity is driven by the pKa of the sulfhydryl (–SH) group. Since the pKa of the SH group in cysteine is 8.3 only about 10% of free cysteine is ionized at the physiological pH. In proteins, the electrostatic environment around the SH group of cysteine residues may render these groups more acidic and, therefore, they may have an increased reactivity towards H_2O_2 , since a higher fraction will be in the thiolate form. Nucleophilicity is also an important factor and, in several proteins, a lower stabilization of the thiolate in cysteine residues increases nucleophilicity of the thiolate [13] and increases, by several orders of magnitude, the rate constants with H_2O_2 (see Table 1). The concept of redox signaling by H_2O_2 was proposed following the discovery of proteins involved in signaling, such as phosphatases, kinases and transcription factors, that contain cysteine residues whose SH groups are oxidized (Fig. 1) causing a change of their biological activity. According to this paradigm upon an increase in the concentration of H_2O_2 , these proteins are specifically oxidized, and a cascade of molecular events ensues. Unfortunately, the wealth of data identifying reactive SH groups, *i.e.* groups that are oxidized upon exposure to an oxidant, contrasts with the near absence of quantitative kinetic data characterizing this reactivity. The few rate constants listed in Table 1 show that the reactivity with H_2O_2 of signaling proteins like the phosphatases Cdc25B and PTB1B is much lower than the reactivity of peroxiredoxins (Prxs), of the selenocysteine residues present in glutathione peroxidases (GPx), or of the heme center present in catalase. In addition, the cellular abundance of antioxidant enzymes like GPx, Prxs and catalase is much larger than that of signaling proteins like phosphatases or TFs. This is important since in the reaction of H_2O_2 with thiols we are dealing with second order rate constants, *i.e.* the rate of reaction is proportional to the concentrations of H_2O_2 and

the thiol. The consequence is that signaling molecules cannot compete with known protein antioxidant systems that remove H_2O_2 . In addition, existing data show that several types of GPx (at least eight isoenzymes) and Prxs (six isoenzymes) coexist [14,15]. If these enzymes had only an antioxidant function, why is there such a variety? For all these reasons, it was concluded that a signaling protein like PTP1B that is redox regulated by H_2O_2 [16–18] but has a low reactivity towards H_2O_2 [19], could not be a direct sensor of H_2O_2 [13,20–22]. Also, antioxidant systems like Prxs and GPx would constitute a kinetic bottleneck that avoids any significant reaction of H_2O_2 with signaling low-reactive proteins [13]. Instead, a high reactive protein, like a peroxiredoxin or a glutathione peroxidase, would be the initial H_2O_2 sensor, which through a thiol-disulfide reshuffling transfer reaction would then oxidize the target protein. This paradigm was inspired in the activation mechanism of the OxyR TF in bacteria [23]. However, these kinetic considerations do not tell the whole story.

- (1) Different H_2O_2 signaling pathways are triggered by different H_2O_2 concentrations and occur with different kinetics. For example exposure of H4IIEC hepatocytes to extracellular H_2O_2 (25–50 μM) for 3 h decreased insulin-stimulated AKT phosphorylation, and increased the phosphorylation of both JNK and its substrate c-JUN, while lower concentrations of H_2O_2 (5–10 μM) enhanced insulin-stimulated phosphorylation of AKT [24]. In addition, H_2O_2 exerts often biphasic responses in which one effect is reversed in a narrow range of concentration such as in H_2O_2 regulation of fatty acid synthase [25–27]. If the initial target is a high-reactive molecule, it is hard to imagine such quantitative diversity in H_2O_2 response.
- (2) More importantly, information is not mass. That antioxidant systems impose a kinetic bottle-neck for the flux of H_2O_2 , and that a rate of oxidation of a sensor is vastly outcompeted by the rate of oxidation of antioxidant systems is irrelevant for a sensing mechanism. In Fig. 2, we simulate a situation where an antioxidant system outcompetes the reaction of H_2O_2 with PTP1B by nine orders of magnitude and, in spite of that, PTP1B is oxidized with a half-life of 5.7 min, a time scale typical of a signaling response. The role of a sensor is to interact selectively with the signaling molecule and to produce an effect that can be measured by a transducer. So, its main role is to transmit information and not, *e.g.* to be a bulk catalyst in a biochemical pathway. What is important is that a variation of H_2O_2 concentration is sensed and this information is transmitted downstream the signaling cascade. By sensing we mean the rate of oxidation of the sensor increases/decreases upon an increase/decrease in the H_2O_2 concentration (the signal). If the rate of oxidation of the sensor is many orders of magnitude lower than the rate of production of H_2O_2 or the rate of H_2O_2 consumption by antioxidant systems, this is actually a good characteristic for a sensor. An ideal sensor does not change the intensity of the signal, it just responds to a change in the signal. For example, a thermometer in a water bath senses changes in the temperature, and does not decrease or increase the temperature of the water. One biochemical illustration of this is the HIF system sensing O_2 . In this system, a prolyl hydroxylase (PHD) catalyzes the hydroxylation of the subunit HIF-1 α by O_2 , which is then subsequently marked for degradation [28]. The fraction of O_2 consumed by PHD compared with the overall cellular O_2 consumption is small, but this does not prevent it from being an O_2 sensor.

Thus, a putative target with reactivity towards H_2O_2 much lower than other molecules also present in the system does not, *per se*, exclude it from being a sensor. Next, we evaluate whether the known characteristics of low-reactivity thiol proteins are compatible with a

Table 1

Rate constants for the reactions between H_2O_2 and several thiol and metal proteins. The intracellular steady-state H_2O_2 concentrations needed to obtain a response time of 30 s, 5 min and 1 h were calculated with Eq. (7).

	Rate constant ($\text{M}^{-1} \text{s}^{-1}$)	$[\text{H}_2\text{O}_2]$ needed for a response time of 30 s (μM)	$[\text{H}_2\text{O}_2]$ needed for a response time of 5 min (μM)	$[\text{H}_2\text{O}_2]$ needed for a response time of 1 h (μM)
Thiol-protein				
GSH	0.87	2.7×10^4	2.7×10^3	220
Thioredoxin	1.05	2.2×10^4	2.2×10^3	180
PTP1B	20	1.2×10^3	120	9.6
KEAP1	140	170	17	1.4
Cdc25B	160	140	14	1.2
GAPDH	500	46	4.6	0.39
Peroxioredoxin-5	3.0×10^5	0.077	7.7×10^{-3}	6.4×10^{-4}
Peroxioredoxin-2	1.0×10^7	2.3×10^{-3}	2.3×10^{-4}	1.9×10^{-5}
Metal-protein				
PerR	1.0×10^5	0.23	2.3×10^{-2}	1.9×10^{-3}
Catalase	2.0×10^7	1.2×10^{-3}	1.2×10^{-4}	9.6×10^{-6}

The rate constant for the reaction of H_2O_2 with KEAP1, was estimated in this work from data from [189]. The other rate constants are for pH 7.4–7.6 at 37°C unless noted otherwise. References: GSH [361] GAPDH [362] estimated for 37°C from $100 \text{ M}^{-1} \text{s}^{-1}$ measured at 0°C ; thioredoxin [363]; PTP1B [19]; Cdc25B [364]; human peroxiredoxin 2 [365] and human peroxiredoxin 5 [366] both at $20\text{--}25^\circ\text{C}$; PerR [300] at pH 7; and catalase [367].




Fig. 1. Oxidative modifications of cysteine (**A** and **B**) and histidine (**C**) residues in proteins induced by H_2O_2 . In cells, sulfhydryl (SH) groups of cysteine residues with low pKa may ionize forming thiolates. Thiolates are good nucleophiles and form a sulfenic acid (SOH) upon reaction with H_2O_2 (reaction 1). Once formed, the SOH can be reduced to a disulfide by a reaction with the SH group of another cysteine residue either in the same (reaction 7) or in a second protein (reaction 5). Alternatively, a SOH can react with the low molecular weight thiol glutathione (GSH) (reaction 4) to form a mixed disulfide in a reaction known as S-glutathionylation or S-thiolation. In an event where a neighboring cysteine residue or GSH is absent, the amide nitrogen of a neighboring amino acid residue can attack the SOH to form a sulfenamide (reaction 6). This reaction occurs in PTP1B. The SOH can also react further with H_2O_2 to generate more oxidized forms of sulfur, the sulfenic acid (SO_2H) (reaction 2) and sulfonic acid SO_3H (reaction 3). Disulfides can be reduced back to thiols using the thioredoxin/thioredoxin reductase and glutaredoxin/GSH/glutathione reductase systems. Sulfenic acids in 2-cys Prxs, but not other proteins, can be reduced to thiols using the enzyme sulfiredoxin [372]. No known enzyme is able to catalyze the reduction of sulfonic acids in proteins. In proteins containing iron metal centers such as PerR, histidine residues can be oxidized by H_2O_2 in a Fenton-like reaction possibly involving the formation of the hydroxyl radical as an intermediate, to form 2-oxo-histidine.

role as H_2O_2 sensors. A chemical sensor should have the following characteristics.

- (1) It does not consume the chemical signal it is responding to,





Fig. 2. PTP1B signaling by H_2O_2 when in the presence of an antioxidant kinetic bottleneck that outcompetes the rate of PTP1B oxidation. The following reactions were included: a rate of H_2O_2 production of $1.2 \times 10^{-2} \text{ M s}^{-1}$; H_2O_2 consumption via glutathione peroxidase ($V_{\text{GPx}} = k_{\text{GPx}} \times [\text{GPx}] \times [\text{H}_2\text{O}_2]$), via PTP1B ($V_{\text{PTP1B}} = k_{\text{PTP1B}} \times [\text{PTP1B}_{\text{rd}}] \times [\text{H}_2\text{O}_2]$) and via non-enzymatic reaction with GSH ($V_{\text{GSH}} = k_{\text{GSH}} \times [\text{GSH}] \times [\text{H}_2\text{O}_2]$). $k_{\text{GPx}} = 6 \times 10^7 \text{ M}^{-1} \text{s}^{-1}$, $[\text{GPx}] = 2 \times 10^{-6} \text{ M}$, $k_{\text{PTP1B}} = 20 \text{ M}^{-1} \text{s}^{-1}$, $[\text{PTP1B}_{\text{tot}}] = 8.3 \times 10^{-9} \text{ M}$, $k_{\text{GSH}} = 0.87 \text{ M}^{-1} \text{s}^{-1}$, $[\text{GSH}] = 5 \times 10^{-3} \text{ M}$. With these parameters, the steady state obtained was $[\text{H}_2\text{O}_2] = 1 \times 10^{-4} \text{ M}$. $[\text{PTP1B}_{\text{tot}}] = [\text{PTP1B}_{\text{rd}}] + [\text{PTP1B}_{\text{ox}}]$, where the subscripts tot, rd and ox, refer to the total amount of PTP1B, to the reduced and to the oxidized forms of PTP1B, respectively.

which, as we have seen, is verified for low-reactivity thiol proteins. Sensor functions may be combined with other functions, as is the case of peroxiredoxin 1 in the AKT signaling pathway, in which H_2O_2 sensing and control of H_2O_2 are combined in the same molecule [29].

- (2) It should be sensitive to changes in the concentration of the chemical signal it is sensing. Visiting again O_2 sensing by the HIF system, the K_m towards O_2 of PHD is $100 \mu\text{M}$ [30], much higher than the endogenous concentration of O_2 (approximately $30 \mu\text{M}$), and so the HIF system responds to O_2 changes in the operational O_2 concentration range *in vivo*. When the O_2 concentration falls, the rate of hydroxylation of HIF-1 α decreases and, consequently, HIF is not degraded and triggers gene expression. In the case of protein thiols, the reaction between the thiol and H_2O_2 is a second-order reaction, and so the rate of reaction depends on H_2O_2 concentration.
- (3) Finally, a sensor should have dynamic characteristics that suit its function. The reactivity of the sensor has to be such that before the H_2O_2 signal is terminated the sensor is activated, *i.e.* it is oxidized by H_2O_2 . To analyze this issue the reactivity of

thiol proteins towards H_2O_2 needs to be evaluated.

To help this analysis, a minimal mathematical model can be set up according to the following two reactions:



For these two reactions the rate laws are defined as follows:

- For the activation step (1) $v_1 = k_{\text{activation}} \times [\text{Target}_{\text{reduced}}]$, where $k_{\text{activation}} = k_{\text{target} + \text{H}_2\text{O}_2} \times [\text{H}_2\text{O}_2]$. $k_{\text{target} + \text{H}_2\text{O}_2}$ is the rate constant for the direct reaction between H_2O_2 and the thiol protein.
- For the switch-off step (2), in which the oxidized protein is regenerated back to the reduced form, $v_2 = k_{\text{switchoff}} \times [\text{Target}_{\text{oxidized}}] = k_{\text{switchoff}} \times ([\text{Target}]_{\text{total}} - [\text{Target}_{\text{reduced}}])$, assuming that the total concentration of the target protein is constant ($[\text{Target}]_{\text{total}} = [\text{Target}_{\text{oxidized}}] + [\text{Target}_{\text{reduced}}]$).

With this, the following differential equation is set up, where $\text{Target}_{\text{reduced}}$ is the fraction of the target thiol protein in the reduced state:

$$\frac{d\text{Target}_{\text{reduced}}}{dt} = k_{\text{switch off}}(1 - \text{Target}_{\text{reduced}}) - k_{\text{activation}}\text{Target}_{\text{reduced}} \quad (3)$$

The analytical solution of Eq. (3) is the following:

$$\text{Target}_{\text{reduced}}|_t = \frac{k_{\text{switch off}}}{k_{\text{switch off}} + k_{\text{activation}}} + e^{-(k_{\text{switch off}} + k_{\text{activation}})t} \times \left(\text{Target}_{\text{reduced}}|_0 - \frac{k_{\text{switch off}}}{k_{\text{switch off}} + k_{\text{activation}}} \right) \quad (4)$$

$$\text{Target}_{\text{reduced}}|_t = 1 - \text{Target}_{\text{oxidized}}|_t \quad (5)$$

Some useful information can be taken from Eqs. (4) and (5):

- If the activation of the thiol protein is not switched-off (i.e., $k_{\text{switchoff}} = 0$), Eq. (4) simplifies to a simple exponential decay (Eq. (6)) and the response time, defined as half of the total response, is given by Eq. (7):

$$\begin{aligned} \text{Target}_{\text{reduced}}|_t &= e^{-k_{\text{activation}}t}(\text{Target}_{\text{reduced}}|_0) \\ \Leftrightarrow \ln\left(\frac{\text{Target}_{\text{reduced}}|_t}{\text{Target}_{\text{reduced}}|_0}\right) &= -k_{\text{activation}} \times t \end{aligned} \quad (6)$$

$$\tau_{1/2} = \frac{\ln(2)}{k_{\text{activation}}} = \frac{\ln(2)}{k_{\text{target} + \text{H}_2\text{O}_2} \times [\text{H}_2\text{O}_2]} \quad (7)$$

Eq. (7) can be manipulated to calculate not only the $\tau_{1/2}$, but also the $[\text{H}_2\text{O}_2]$ or the $k_{\text{target} + \text{H}_2\text{O}_2}$, provided two of these parameters are known. In this case, at the end of the response all protein will be activated since there is not an operating switch-off mechanism.

- If the thiol protein is switched-off ($k_{\text{switchoff}} > 0$), the steady-state fraction of protein present in the oxidized form is given by Eq. (8) and the response time, defined as half of the total response, is given by Eq. (9):

$$\text{Target}_{\text{oxidized}}|_{\text{steadystate}} = 1 - \frac{k_{\text{switch off}}}{k_{\text{switch off}} + k_{\text{activation}}} \quad (8)$$

$$\tau_{1/2} = \frac{\ln(2)}{k_{\text{activation}} + k_{\text{switch off}}} \quad (9)$$

In this article we will apply only Eqs. (6) and (7) because there are no quantitative data available for the switch-off mechanism that regenerates the H_2O_2 sensor. This approximation is acceptable for time courses where the switch-off mechanisms are not operating at a significant rate. In Table 1, we show H_2O_2 concentrations needed to have a response time of 30 s, 5 min and 1 h, calculated by applying Eq. (7) for those proteins whose rate constant with H_2O_2 is known. In other words, the intracellular steady-state H_2O_2 concentrations indicated are those necessary to oxidize the listed proteins by 50% after exposing it to H_2O_2 for 30 s, 5 min and 1 h. As can be observed, if a fast response is necessary (30 s), or if the H_2O_2 transient signal lasts only 30 s, only Prxs, PerR and catalase are sufficiently sensitive targets as to provide the desired response. For other targets, like PTP1B, the H_2O_2 signaling concentration needed to trigger the response during the 30 s of the duration of the signal would be too high, 1.2 mM. However, if cells require a slow response (1 h), or if the H_2O_2 transient signal lasts for 1 h, even a low reactive sensor, such as Cdc25B, will be sufficient to mediate the signaling pathway, as exposure to a 1.2 μM H_2O_2 concentration during 1 h would be enough to activate the response, i.e. to oxidize Cdc25B by 50%. Thus, the duration of the transient H_2O_2 signal is an important experimental observation that gives a hint on whether a sensor with high or low reactivity is operating. In this regard, for example, H_2O_2 production triggered by EGF peaks at 5 min, and returns to baseline after 20 min [31] or 60 min [32]. While a short H_2O_2 transient signal excludes the possibility that a low-reactive-sensor is operating, a long transient signal is compatible with both a high and low-reactivity sensor. The same kind of information can be inferred from the time course of the signaling pathway: a very fast response is incompatible with a low-reactive sensor, while a slow response may be the result of either a low-reactive sensor that takes time to respond or, alternatively, the result of a high-reactive sensor that responds rapidly, but then oxidizes slowly an effector molecule. Using PTP1B as an example to analyze these two scenarios, analysis of experimental data where recombinant PTP1B inactivation was studied as a function of H_2O_2 concentration *in vitro* [33] revealed a $k_{\text{target} + \text{H}_2\text{O}_2} = 22 \text{ M}^{-1} \text{ s}^{-1}$ (Fig. 3), which is near the published values (see Table 1) establishing this protein as a low reactivity thiol sensor protein. Also, in the same work, *in vivo* activation by EGF caused a 35% inactivation of PTP1B after 5 min. If a direct oxidation of PTP1B by H_2O_2 with a rate constant $k_{\text{target} + \text{H}_2\text{O}_2} = 22 \text{ M}^{-1} \text{ s}^{-1}$ is assumed, a local concentration of H_2O_2 near 66 μM would be needed. However, if we consider that a high-reactive thiol sensor protein reacts with H_2O_2 , and then relays the signal to PTP1B, a lower H_2O_2 local concentration would be needed. An extra layer of uncertainty is whether the rate constants determined *in vitro* are the same operating for the reaction *in vivo* and whether H_2O_2 derivatives, like peroxymonophosphate [34] and peroxymonocarbonate [35,36], which have higher reactivity towards PTP1B, operate *in vivo*.

Another important parameter to take into account when discussing sensors is the intensity of the H_2O_2 signal, and hence the notion of localized H_2O_2 concentrations should also be considered. The extracellular H_2O_2 threshold concentration that triggers apoptosis in Jurkat T-cells is 7 μM [37], which considering the H_2O_2 gradient across the plasma membrane, converts to an intracellular H_2O_2 concentration probably lower than 1 μM [38]. However, cells may tolerate relatively high localized H_2O_2 concentrations for a short period of time. In recent years it became clear that cells have developed several strategies to insure signaling H_2O_2 concentrations are reached only in localized compartments near the site of its production [39]. For example, H_2O_2 -dependent redox regulation of PTP1B requires colocalization of PTP1B with the NADPH oxidase Nox4 in the endoplasmic reticulum, with cytosolic PTP1B being insensitive to overexpression of endoplasmic-reticulum Nox4 [40].




Fig. 3. Application of Eq. (6) to estimate rate constants between cellular targets and H_2O_2 . Plot of the fraction of PTP1B activity observed *in vitro* after 10 min (A) and of the reduced form of KEAP1 observed in HeLa cells after 5 min (B) of incubation with the indicated H_2O_2 concentrations. Experimental data are taken from [33] and [189], respectively for PTP1B and KEAP1. If a simple exponential decay is considered, that is no regeneration of sensor occurs, the slope of these plots is $k_{\text{target} + \text{H}_2\text{O}_2} \times t$ (Eq. (6)) and, therefore, the rate constants between PTP1B and KEAP1 with H_2O_2 are estimated at $22 \text{ M}^{-1} \text{ s}^{-1}$ and $20 \text{ M}^{-1} \text{ s}^{-1}$, respectively. If a gradient between extracellular and intracellular H_2O_2 of 6.8 is considered in HeLa cells [42], the rate constant for H_2O_2 reaction with KEAP1 is estimated at $140 \text{ M}^{-1} \text{ s}^{-1}$.

Upon activation of receptor activated kinases, H_2O_2 is produced either in specific endosomes or in localized sites near the plasma membrane depending on the cell type. Biomembranes constitute a permeability barrier to H_2O_2 [38,41–43] and may help maintain higher H_2O_2 concentrations near its local of production. Because the permeability of the plasma membrane is regulated by H_2O_2 [44–46], it may be hypothesized that membrane domains near the site of H_2O_2 production are altered in order to have a lower permeability towards H_2O_2 . Indeed, plasma membrane permeability towards H_2O_2 may range from near complete permeable in yeast mutants of the ergosterol pathway [44] to a near complete impermeability in human spermatozoa [47]. Furthermore, aquaporins also regulate H_2O_2 transport across biomembranes [48] and mediate intracellular H_2O_2 signaling [49], providing an additional potential control step.

In addition to a localized production of H_2O_2 , the local inhibition of antioxidant systems, like Prxs, also contributes for a localized increase in the concentration of H_2O_2 [50,51]. Such strategy is reminiscent of the signaling mediated by phosphorylation in which both activation of kinases and inhibition of phosphatases occur. Concerning the inactivation of Prxs, two strategies have been proposed. (i) The so-called floodgate hypothesis in which an overoxidation of the catalytic cysteine residues of Prxs results in the inhibition of the peroxiredoxin-catalyzed reduction of H_2O_2 . Overoxidation of peroxiredoxin is observed with a high concentration of H_2O_2 , but recent studies showed that H_2O_2 levels reached during signaling are not enough to overoxidize peroxiredoxin [29,50]. (ii) An alternative strategy is the inhibition of peroxiredoxin activity by its phosphorylation [50,15] (Fig. 4). The work of Woo et al. [15,50] showed that upon binding of a ligand to a membrane receptor an SRC family kinase is activated. This SRC kinase activates NADPH oxidase in the plasma membrane, which leads

to the production of superoxide that dismutates into H_2O_2 , and also catalyzes Prx1 phosphorylation at a tyrosine residue. This leads to inactivation of Prx1, due to a decreased reactivity of its catalytic cysteine residue with H_2O_2 , and to a transient accumulation of H_2O_2 around membranes, where signaling components are concentrated. The increased levels of H_2O_2 promote further phosphorylation and inactivation of Prx1 both by activating SRC kinases and by inactivating PTPs. There is no cellular toxicity because this increase in H_2O_2 concentration occurs locally and any H_2O_2 that diffuses from this region will be degraded by active Prx1 and other peroxidases present in the cytoplasm. It should be mentioned that not all antioxidant systems may be present at the site of H_2O_2 production; for example glutathione peroxidase is not found in the sub-membrane fraction where H_2O_2 is produced [50]. The lag time in Prx1 catalysis caused by its phosphorylation is inversely proportional to the concentration of H_2O_2 . This suggests that reactivation of Prx1 resumes when H_2O_2 levels rise beyond a certain threshold contributing to the termination of the signaling process [52]. Recently two other kinases, Mst1 and Mst2, which are both activated by H_2O_2 , were shown to inhibit Prx1 [53]. Both these studies have an important general implication for H_2O_2 -dependent redox regulation since they also suggest that phosphorylation/dephosphorylation of thiol proteins can alter their reactivity with H_2O_2 and so, we could speculate that an H_2O_2 sensor with low reactivity can become a high reactivity sensor and vice-versa, depending on its phosphorylation state.

A note should be made regarding the possible role GSH may have in mediating or modulating H_2O_2 signaling. GSH is several orders of magnitude more abundant than a low-reactive thiol protein such as PTP1B, and has a rate constant for the reaction with H_2O_2 , $0.87 \text{ M}^{-1} \text{ s}^{-1}$ at pH 7.4, that is about 20–40 times lower than that of PTP1B. Taking into account these data, can GSH be considered a sensor molecule for H_2O_2 or can it inhibit H_2O_2 signaling mediated by PTP1B? The answer in both cases is no. In terms of reaction with H_2O_2 , GSH certainly outcompetes PTP1B (Fig. 2), but the non-enzymatic reaction of H_2O_2 with GSH is negligible when compared with enzymatic systems removing H_2O_2 , like catalase, GSH peroxidase or peroxiredoxins. Thus, the non-enzymatic reaction of GSH with H_2O_2 does not affect significantly the intensity of the H_2O_2 signal, and does not inhibit PTP1B-mediated H_2O_2 signaling. Concerning the non-enzymatic oxidation of GSH, this does not represent a signaling event because the product of this reaction does not relay information into a signaling pathway. It could be argued that an increased GSSG concentration would affect signaling by changing the ratio $2 \times [\text{GSH}]/[\text{GSSG}]$, but the contribution of the non-enzymatic oxidation of GSH towards this ratio is negligible when compared with the enzymatic oxidation of GSH.

So far we have been discussing H_2O_2 signaling mediated by its direct reaction with thiol proteins, but alternative mechanisms have been described, involving H_2O_2 -dependent formation of other second-messengers. For example, H_2O_2 formed in the mitochondria may initiate lipid peroxidation to produce reactive electrophilic lipid oxidation products that can act as second messengers leading to the activation of mitogen-activated protein kinases [54]. This initiation of lipid peroxidation may be mediated by heme proteins such as cytochrome c [55] and, in general, because of the high-reactivity of heme iron with H_2O_2 , heme proteins could potentially act as H_2O_2 sensors. We could also speculate that the localized production of H_2O_2 in entrapped membrane compartments during signaling could initiate lipid peroxidation and form reactive lipid species, which has been suggested to have a signaling role [56].

From the data in Table 1 it becomes obvious that H_2O_2 signaling can operate either mediated by localized high transient levels of H_2O_2 that activate sensors with a low reactivity, or mediated by proteins with high reactivity towards H_2O_2 that work as the initial sensor that subsequently activate a low reactivity protein. Experimental support of both these mechanisms does exist [29,50] (Fig. 4).




Fig. 4. Localized increase of H₂O₂ levels mediated through inhibition of peroxiredoxins activity by its phosphorylation. Peroxiredoxins (Prxs) can act as highly reactive H₂O₂ sensors and transduce the signal to a signaling molecule (SM). Alternatively, upon binding of a ligand to a membrane receptor an SRC family kinase can be activated. This SRC kinase activates NADPH oxidase in the plasma membrane, which leads to the production of H₂O₂, and catalyzes phosphorylation of Prx at a tyrosine residue leading to its inactivation. Prx inactivation leads to a transient accumulation of H₂O₂ around membranes, where signaling components are concentrated. This will promote the direct oxidation of H₂O₂ sensors with intermediate and low reactivity.

Biological functions of transcription factors regulated by H₂O₂

Before addressing the known mechanisms of H₂O₂-regulated TFs we briefly describe their biological functions. Particular attention is given to three evolutionary aspects that allowed H₂O₂ to evolve as a regulatory molecule. First, we address the constraints imposed by the lack of compartmentalization in bacteria, then the appearance of sub-cellular compartments in eukaryotes and, finally, the impact of multicellularity for H₂O₂ signaling. As a result, **as we progress throughout the phylogenetic tree there is an increase in the complexity of the regulatory networks, involving TFs that are able to respond to variations of H₂O₂ levels.**

Bacterial transcription factors as direct sensors of H₂O₂

In organisms without cellular compartments, one way of achieving signaling is to use **highly reactive proteins** able to sense H₂O₂ and trigger a response. This is obligatory if **a fast response is** required. In fact, in the case of bacteria we highlight two TFs, OxyR and PerR, that are both directly regulated by H₂O₂. This duo of TFs, which are highly reactive with H₂O₂, display only one regulatory mechanism layer, which rapidly allows bacteria exposed to increasing levels of H₂O₂ to cope with oxidative damage increasing cell fitness and survival. The small size of bacteria, with the corresponding high ratio between surface area and volume make them particularly susceptible to environmental stresses, including H₂O₂. For example at anoxic/oxic interfaces oxidation reactions involving reduced metal ions and sulfur species that enter in contact with oxygenated waters produce H₂O₂. H₂O₂ is also formed when UV/visible radiation illuminates extracellular chromophores, including photosynthetic pigments that are released by decomposing plants. Also, H₂O₂ may be intentionally produced by competing organisms like lactic acid bacteria [57]. Thus, **the capacity to rapidly respond to increasing concentrations of H₂O₂ will probably provide a survival advantage in various ecosystems.**

OxyR

OxyR is a member of the LysR family of TFs that contains a conserved N-terminal helix–turn–helix DNA binding domain, a central co-inducer recognition and a response domain that senses the regulatory signal, and a C-terminal domain that is required for multimerization and activation [7,58,59]. In *E. coli*, tetrameric OxyR binds to the 5' promoter–operator regions of target genes at a conserved sequence motif that contains four ATAG elements spaced at 10 bp intervals [60]. OxyR binds to DNA, either in its oxidized or in its reduced form, but only activates transcription when oxidized [61]. In the oxidized form OxyR contacts the DNA motif in four adjacent major grooves on one face of the DNA helix while the reduced form of OxyR binds DNA in

two pairs of major grooves separated by one helical turn [60]. **Most of the OxyR up-regulated genes are involved in defense systems against oxidative stress** [7,58].

PerR

The Peroxide Regulon Repressor (PerR) is a metal-dependent TF and a major regulator of the peroxide inducible stress response in bacteria [62–64]. PerR was identified in 1998 and found to be a member of the ferric uptake repressor (Fur) family of proteins [62,65]. Unlike most members in the Fur family, PerR is not involved in metal homeostasis and, like OxyR, is a specific sensor of H₂O₂ [66]. In fact, PerR is a functional equivalent for OxyR and substitutes OxyR in many Gram-positive bacteria, although it may also coexist with OxyR [67]. However, like the other Fur family proteins, PerR DNA binding is also activated by a metal ion, either Fe²⁺ or Mn²⁺. PerR interacts with DNA at the *per* box, a specific palindromic consensus sequence (TTATAATNATTATAA) residing within and near the promoter sequences of PerR-controlled genes. In *Bacillus subtilis*, PerR, when bound to DNA, represses the genes coding for proteins involved in the oxidative stress response (*kata*, *ahpC*, and *mrgA*) [62,68], metal homeostasis (*hemAXCDBL*, *fur*, and *zsaA*) [68–70] and its own synthesis (*perR*) [69]. Most PerR-regulated genes are de-repressed in cells treated with low levels of extracellular H₂O₂ (8 μM) [64] or cells cultured under conditions of iron and manganese ions deficiency [69].

The challenge of cellular compartmentalization in lower eukaryotes

In yeast, a eukaryote but still a unicellular organism, we will focus on the analysis of four TFs, namely, Yap1, Maf1, Hsf1, and Msn2/4. In this group Yap1 is regulated by H₂O₂ at the level of cytoplasm/nucleus traffic, which creates a new layer on the regulatory mechanisms when compared with OxyR and PerR. Thus, though Yap1, like OxyR and PerR, essentially allows cells to deal with oxidative stress response, the complexity of its regulatory mechanism already reflects cell compartmentalization. This regulatory layer is also found in Maf1, but the traffic regulator partners are now replaced by post-translational modifications (PTMs) that create different intracellular pools of the protein and determine its subcellular localization. In the case of Hsf1 and Msn2/4, both TFs allow cells to respond to a variety of different environmental stresses from heat shock to starvation and oxidative stress. Thus, in yeast, the response to oxidative stress is also part of a more general cellular response to stress, probably making this response to oxidative damage more robust.

Compartmentalization creates new opportunities to generate new levels of regulation and confine certain pathways and metabolic pathways to specific compartments. Moreover, yeast cells are larger than bacteria and this may have contributed to create endogenous

H₂O₂ gradients between distinct compartments [44]. In the case of cytoplasm/nucleus the appearance of these two compartments allowed DNA to be less susceptible to oxidative damage contributing to distinguish H₂O₂ toxic and regulatory responses, thus facilitating evolution of H₂O₂ as a regulatory molecule. Also, endogenous sources of H₂O₂, such as dismutation of superoxide produced by NADPH oxidase (NOX) enzymes began to be established. Up until recently, it was believed that the genomes of *Saccharomyces cerevisiae* (*S. cerevisiae*) and *Schizosaccharomyces pombe* (*S. pombe*) did not contain genes encoding NOX enzymes. Recently, Rinnerthaler et al. [71] showed that one of the *S. cerevisiae* ORF encodes an authentic NOX, which is located in the endoplasmic reticulum (ER) membrane and produces superoxide in a NADPH-dependent fashion. Interestingly, most NOX enzymes are plasma membrane bound but, NOX4 has been localized in endomembranes of ER, the nucleus and, in spite of some contradictory data, in the mitochondria [72]. This clearly shows that the appearance of the intracellular membranes associated to new compartments contributed to the evolution of a regulatory role for H₂O₂.

Yap1

Yap1 (yeast AP-1) is one of the members of the yeast *S. cerevisiae* activator protein (Yap) family that comprises eight members [73]. All members display a significant sequence similarity at the DNA-binding domain, the basic leucine zipper (b-ZIP domain) in the N-terminus [74]. Structurally the Yap1 factor has two cysteine residues rich domains (CRD), the nCRD (Cys303, Cys310 and Cys315) and cCRD (Cys598, Cys620 and Cys629) located in the N- and C-terminal, respectively [75]. Under oxidative stress, nuclear export of Yap1 is decreased and Yap1 is retained in the nucleus where it can regulate its target genes [76]. Yap1 has a key role in the oxidative stress response, redox homeostasis and electrophilic response, regulating the transcription of genes encoding antioxidant and detoxification enzymes.

Maf1

Maf1 is a transcriptional repressor of RNA polymerase III (Pol III) that was originally discovered in *S. cerevisiae* [77]. However, Maf1 is also found in human, animals and plants [78]. In yeast, Pol III is responsible for the transcription of around 300 different genes, mostly tRNA genes [79]. Maf1 does not bind directly to DNA; instead it binds to Pol III clamp and rearranges the subcomplex C82/34/31, which is required for transcription initiation [80]. In this repressive complex, Maf1 impairs recruitment of Pol III to a complex of promoter DNA with the initiation factors TFIIB and thus prevents formation of a closed-complex.

Maf1 is a hydrophilic protein conserved from yeast to human and it contains three signature domains not found in any other polypeptide: A, B and C boxes [81]. Yeast Maf1, contrary to human MAF1, contains two conserved nuclear localization signals (NLS) [77]. Maf1 activity is regulated by means of its phosphorylation state-dependent cellular localization [78].

Hsf1

In eukaryotic cells, the heat shock response is primarily mediated by a homotrimeric DNA-binding TF, the heat shock factor 1 (HSF1), a member of the heat shock TF family that binds to *cis*-acting promoter elements in target genes, called heat shock elements (HSE) [82]. Each HSE contains two, or more, contiguous inverted repeats of the 5-bp sequence nGAAn. HSF1 is regulated differently in mammalian cells and in yeast [83].

The *S. cerevisiae* HSF1 homologue, Hsf1, is constitutively a trimer, and is localized in the nucleus where it associates with high-affinity HSEs under normal conditions and additional HSEs during stress [83,84]. The structure of Hsf1 is highly conserved with its mammalian homologue [82]. Besides being involved in the heat shock response several lines of evidence have shown that Hsf1 is also involved in the yeast oxidative stress response [85]. Target genes of Hsf1 encode for

molecular chaperones such as heat shock proteins (HSPs), metabolic enzymes, and cell wall proteins [86].

Msn2/4

Msn2 and Msn4 (Msn2/4) are homologous and functionally redundant Cys2His2 zinc finger yeast TFs [87]. In *S. cerevisiae*, disruption of both *MSN2* and *MSN4* genes results in a higher sensitivity to different environmental stresses, including carbon source starvation, heat shock and severe osmotic and oxidative stresses. Msn2/4 are required for activation of several yeast genes, whose induction is mediated through the presence of a stress responsive element (STRE) consisting of a pentameric core of CCCCT, such as *CTT1*, coding for cytosolic catalase, and *HSP12* [87].

The challenge of multicellularity in higher eukaryotes

The complexity of the next step in eukaryotes evolution is attained by acquisition of multicellularity and, therefore, the appearance of different cell types and tissues. Cells are now able to differentiate and gain very specialized functions, their proliferative organization is diverse and cells must integrate cell–cell and cell–matrix connections. Some cells maintain their ability to be totipotent, and multicellularity is now upstream of cell compartmentalization. Cells are now exposed to a new environment and information is received from this environment mainly *via* receptors in the plasma membrane.

Interestingly, in this new scenario, H₂O₂, despite still having the capacity to cause damage, acquires a prominent role as a regulatory molecule. Now, cells do not only respond to environmental H₂O₂, but H₂O₂ evolved as a second messenger necessary for many signaling pathways. Therefore, although the stress-response strategies found in yeast for the regulation of transcriptional factors by H₂O₂ can be detected in multicellular organisms (for example, HSF1 and NRF2), now evolution expanded this to an incredible complexity and invented new regulatory mechanisms and combinations between the pre-existent mechanisms allowing each protein to have different code bars and, therefore, integrate different signaling pathways and compartments. In a few cases, the TF is even part of a membrane receptor and is released upon activation, playing membranes a critical role.

In multicellular organisms we focused our attention in nine different transcriptional factors, namely AP-1, NRF2, CREB, HSF1, HIF-1, TP53, NF-κB, NOTCH, SP1 and CREB-1. It should be noted that some of these TFs (e.g. TP53, NF-κB, CREB and CREB-1) that are relevant to metazoan multicellularity evolved prior to the emergence of the metazoan stem lineage, and they can be found in some transition organisms, such as choanoflagellates or *Capsaspora owczarzaki*, but not in other lower unicellular eukaryotes such as yeast [88]. Noteworthy, most of them are involved in the regulation of cell damage response, cell proliferation (cell cycle regulation), differentiation and apoptosis (AP-1, CREB, TP53, NOTCH, NF-κB, and SP1). Therefore, they are closely linked to cell survival and development, and their deregulation is in the basis of different pathophysiological stages such as cancer. HIF-1 and SREBP-1 seem to have a narrow range of actions essentially controlling lipid metabolism and O₂ levels at the cellular and systemic level. Like it happens in yeast, HSF1 still orchestrates the cellular response to a variety of cellular stresses, and H₂O₂ is still under the surveillance of general cell protecting mechanisms. The ability of certain TFs for transactivating response genes still recalls OxyR since they directly sense H₂O₂ as is the case of human HSF1.

The way H₂O₂ regulates the activity of these transcriptional factors is diverse but clearly explores the existence of different cellular compartments, and enrichment in different biochemical forms of both the TF protein and its partners, by PTMs and different forms of processing. In all these different levels of regulation we find targets for the regulatory action of H₂O₂.

AP-1

Activator protein-1 (AP-1) is a TF that regulates several cellular processes, including cell proliferation, apoptosis, survival, and differentiation. Such functional diversity derives primarily from its structural and regulatory complexity [89]. The term AP-1 describes a collection of dimeric bZIP proteins, mainly from the Jun (v-Jun, c-JUN, JUND, and JUNB), Fos (v-Fos, c-FOS, FRA-1, FRA-2, and FOSB), ATF/CREB (CREB, ATF1, ATF2, ATF4, ATF5, ATF6a, ATF6b, ATF7, ATF3/LRF1, B-ATF, and ATFa0), JDP (JDP1/2), small MAF (MAFG, MAFF, and MAFK) and large MAF (cMAF, MAFB, MAFA, and NRL) sub-families that usually form heterodimers that bind to a TPA-responsive element (TRE, 5'-TGAG/CTCA-3') or cAMP response elements (CRE, 5'-TGACGTCA-3') [90,91]. In these TFs the basic region of the bZIP domain mediates DNA binding, whereas the leucine zipper is responsible for dimerization with the partner bZIP factor.

NRF2

The NRF protein family constituted by NRF1, NRF2, and NRF3, which are also bZIP proteins, regulate electrophilic xenobiotic detoxification and oxidative stress response. The main activators of this TF are electrophile agents but H₂O₂ also activates NRF2 [92] by a multitude of mechanisms. NRF proteins bind to the electrophile response element (EpRE, 50-(A/G)TGACNNNGC(A/G)-30) in the target genes, cannot form homodimers, and the typical partners are small MAF proteins, although c-JUN has been reported to heterodimerize with NRF2 [90,93]. In this review, we will focus on NRF2.

CREB

The cAMP response element-binding protein (CREB) is one of the three members of the cAMP responsive TFs family occurring in mammals (CREB, CREM and ATF-1). These TF play important roles in the nuclear responses to a variety of external signals, by binding different promoters of genes encoding proteins involved in transcription, metabolism, cell proliferation, differentiation, apoptosis, and the secretory pathway [94]. CREB binds as a homodimer to the CRE conserved TGACGTCA sequence [95]. This 43 kDa TF has a dimerization and DNA binding, and an N-terminal activation domain (AD) with two independent regions: the phosphorylation box (P box) and a second region comprising two glutamine-rich domains, Q1 and Q2, which flank the P box [96]. The P box contains a cluster of sites phosphorylated by various kinases that regulate the transactivation potential of this protein [95]. Phosphorylation of CREB, mainly at Ser133, enables the recruitment of the co-activators CBP/p300 and stimulates CREB-dependent transcription [95]. However, CREB activity depends on other regulatory partners that are required for recruitment of the transcriptional apparatus to the promoter. More than 20 different protein kinases, members of distinct signaling pathways, have been described as CREB kinases [95,97]. The activity of CREB as a TF can be regulated by other PTMs such as acetylation, ubiquitination, sumoylation and glycosylation [95].

TP53

TP53 (Tumor protein TP53, TTP53, Li-Fraumeni syndrome 1) has been studied for nearly three decades, and is best known for its potent ability to be a tumor suppressor [98]. In fact, this protein is encoded by the *TP53* gene that commonly has lost its function by mutations in the majority (75% of the cases) of human cancers.

TP53 is a DNA-binding TF that both activates and represses a broad range of target genes constituting a critical hub that integrates a huge variety of signals and allows a complex set of cellular responses to DNA damaging agents, oxidative stress, oncogene activation (deregulated growth signals), mitotic spindle damage, hypoxia, nutrient deprivation, telomere erosion, ribosomal stress and is involved in cellular senescence [99–102]. The TP53 protein comprises different structural and functional domains. The N-terminal domain corresponds to the transactivation domain required for transcriptional regulation of

the target genes. This domain contains a proline-rich region that plays a role in the regulation of TP53 stability by the negative regulator protein murine double minute 2 (MDM2) [103]. The central domain binds in a zinc-dependent manner [104,105] to a consensus site that shows an internal symmetry and is composed by two copies of the sequence 5'-PuPuPuC(A/T) (T/A)GPyPyPy-3' separated by 0–13 bp [106,107]. The C-terminal region of the protein contains the oligomerization domain (tetramerization) where specific signals for nucleus import/export are localized [108]. This domain catalyzes DNA annealing and strand transfer and displays a strong preference for damaged DNA by ionizing radiation, having thus specialized functions [109].

NOTCH

The receptors and ligands in the NOTCH signaling pathway are membrane proteins that imply cell–cell contact for their activation, and they constitute the basis of NOTCH-dependent transcription activation. Mammals express four receptors, NOTCH1, NOTCH2, NOTCH3 and NOTCH4 and two families of ligands, Jagged (Jagged1 and Jagged2) and Delta-like (Dll1, Dll3 and Dll4) [110]. The canonical model for NOTCH signaling activation requires a crucial proteolysis that releases the NOTCH intracellular domain (NICD) from the plasma membrane after ligand activation expressed in a neighboring cell. NICD cleavage is mediated by the γ -secretase complex and is facilitated by the previous proteolytic cleavage of NOTCH extracellular domain (NECD) by a metalloproteinase ADAM17/TACE [111]. NICD is then translocated to the nucleus where it associates with the DNA-binding protein Suppressor of Hairless (SU(H)) and with the nuclear effector Mastermind (MAM) for transcriptional activation [110]. The NOTCH signaling pathway has been implicated in numerous cellular processes including neuron differentiation and blood vessel formation in normal embryo development and in disease [112].

NF- κ B

The NF- κ B/REL family of TFs has key regulatory roles in inflammation, innate and adaptive immune response, proliferation and apoptosis [113]. It consists of homo- and heterodimers of five distinct proteins, the REL subfamily proteins (p65/RELA, RELB, and c-REL), which contain C-terminal transactivation domains (TADs) and the NF- κ B subfamily proteins (p50, and p52, and its precursors p105 and p100, respectively) [113]. All NF- κ B/REL proteins contain a Rel-1-homology domain (RHD) that also harbors an NLS, which is responsible for dimerization, recognition and binding to DNA and also for the interaction with the inhibitory proteins I κ Bs [114]. The I κ B family is composed of I κ B- α , I κ B- β , I κ B- ϵ , I κ B- γ and BCL-3 (B-cell lymphoma 3) possessing typical ankyrin repeats that mediate binding to the RHD and interfere with its NLS function. The most common composition of cytoplasmic NF- κ B/I κ B complex appears to be the p50/p65/I κ B- α [114].

The I κ B proteins bind to NF- κ B in the cytoplasm preventing NF- κ B translocation to the nucleus and its binding to DNA. Therefore, complexation with I κ Bs has to be removed for NF- κ B activation [115]. NF- κ B activators such as tumor necrosis factor α (TNF- α), lipopolysaccharide (LPS) and interleukin-1 (IL-1) activate the I κ B-kinase complex (IKK complex), which catalyzes the phosphorylation of I κ Bs at specific regulatory amino acid residues. As a consequence, the I κ Bs are targeted for degradation by the 26S proteasome thereby freeing NF- κ B, which translocates to the nucleus and binds to the promoter/enhancer regions of target genes, the κ B sites, which have the general consensus sequence GGGRNNYYCC (R is a purine, Y is a pyrimidine, and N is any base) [113]. Target genes include pro-inflammatory cytokines, chemokines, adhesion molecules, growth factors, and enzymes that produce secondary inflammatory regulators such as cyclooxygenase-2, inducible NO synthase, and heme oxygenase [22,116].

SP1

SP1 (Specificity protein 1) was the first TF to be identified, purified and cloned from mammalian cells [117]. SP1 is a member of an extended family of DNA-binding proteins that have three zinc-fingers (Cys2His₂ – type zinc finger), which are required for recognizing GC-rich promoter sequences [118,119]. SP1 contains two glutamine-rich domains that are essential for transcriptional activation. Next to these domains are serine/threonine-rich sequences that may be a target for PTMs [119]. SP1 is an essential TF that can activate or repress transcription in response to physiological and pathological stimuli, such as oxidative stress [120,121]. SP1, besides regulating itself, is also implicated in the regulation of many genes that play important roles in a variety of physiological processes including cell cycle regulation and growth control, hormonal activation, apoptosis, and angiogenesis [122].

SP1 directly interacts with TATA-binding protein associated factors [123] and other factors, such as those binding to cAMP response elements [124], NF- κ B [125] and vascular endothelial growth factor receptor-2 (VEGFR-2) [126]. The activity of SP1 as a TF can be regulated by PTMs such as phosphorylation, acetylation and methylation [120,127] that regulate SP1 protein level, transactivation activity, and DNA binding affinity [128].

HIF-1

HIF-1 (hypoxia-inducible factor) is a TF that has an essential role in the response to hypoxia at systemic and cellular level. This TF has been implicated in the activation of angiogenesis and erythropoiesis [129,130] and in the metabolic adaptation to hypoxia through activation of glycolysis [131]. HIF-1 is a dimeric protein complex formed by an inducible subunit (HIF-1 α) and a constitutive subunit (HIF-1 β) that are basic helix-loop-helix (bHLH) proteins. The HIF-1 α/β dimer binds to a DNA motif (G/ACGTG) in hypoxia-response elements (HREs) of target genes [132].

HIF-1 α contains two hypoxia-dependent degradation domains with two conserved prolyl residues. The hydroxylation of these residues catalyzed by PHD promotes the interaction between HIF-1 α and the Hippel–Lindau tumor suppressor (pVHL), targeting the former for proteasomal degradation [28,133]. Although the activation of HIF-1 has been mainly related with low levels of O₂, this TF can also be activated by a hypoxia-independent mechanism that is mediated by the superoxide radical and by H₂O₂ [134,135].

SREBP-1

Sterol regulatory element binding proteins (SREBPs) are a family of critical TFs that bind the sterol regulatory element (SRE), activating genes encoding the enzymes that regulate the synthesis of cholesterol, lipids and fatty acids and cellular uptake of lipoproteins [136–138]. H₂O₂ has a strong influence on SREBP1 activity in cells with a high sensitivity to insulin, promoting lipid accumulation [139]. There are three SREBP isoforms, designated SREBP-1a, SREBP-1c, and SREBP-2 [138]. SREBP proteins, initially synthesized as a 125 kDa membrane bound precursor, are anchored to the ER [136,140]. They share a similar tripartite structure: an N-terminal region, which has a TF domain of the basic helix-loop-helix-leucine zipper (bHLHZip) family, a central domain, which contains the two transmembrane spans, and a C-terminal regulatory domain that binds tightly to the C-terminal domain of SREBP cleavage activating protein (SCAP) [141,142]. The complex SREBP–SCAP is maintained in the ER via the interaction with the protein INSIG (Insulin-induced gene), which binds directly to the protein SCAP [143]. Despite differences in their transcriptional targets, the proteolytic activation of each SREBP isoform is regulated by cholesterol and oxysterols through a common mechanism [144,145]. In the presence of these compounds, SREBP–SCAP is retained in the ER by binding to Insig, contrary to what happens in the absence of sterols [143] where Insig no longer binds SREBP–SCAP and SREBP–SCAP is translocated to the Golgi [143]. Once in the Golgi, the SREBP

active form is obtained after two sequential proteolytic cleavages of the SREBP precursor form, mediated by distinct site specific proteases, namely Site-1 protease (S1P) and Site-2 protease (S2P) [146], in order to release the amino-terminal TF domain of SREBP from the membrane [142]. The activated N-terminal domain of SREBP translocates into the nucleus to bind the SRE (ATCACCCAC) sequence and the E-box (CAXXTG) sequence of the promoter of target genes and trigger gene expression [139].

HSF1

The mammalian heat shock factor (HSF) family has four members HSF, HSF2, HSF3 and HSF4 but HSF1 is the key stress-responsive regulator of the heat shock response [82]. HSF1 has several functional domains, the N-terminal DNA-binding domain, the oligomerization domain containing heptad repeat regions HR-A and HR-B that regulates trimerization, the regulatory domain and the C-terminal trans-activation domain [82,147]. An additional heptad domain region, HR-C, that maintains HSF1 in an inactive state by suppressing spontaneous trimerization is located between the regulatory and trans-activation domains. The DNA-binding domain is of a looped helix-turn-helix type but, unlike TFs containing similar domains, HSF1 does not make direct contact with DNA [147]. The loop apparently stabilizes the DNA-bound HSF1 trimer by protein–protein interactions. The regulatory domain in HSF1 negatively regulates the transactivation domain and is targeted by several PTMs including phosphorylation, sumoylation and acetylation [147].

HSF1 is constitutively expressed in most tissues and cell lines [82]. In the absence of stress conditions HSF1 exists as a monomeric phosphorylated protein that interacts with HSP90 and is present in both the nucleus and cytoplasm [82]. In mammalian cells, on exposure to diverse stress conditions, including oxidative stress, monomeric HSF1 undergoes a multistep activation process that includes dissociation from the Hsp90 complex, trimerization, nuclear accumulation, PTMs, DNA binding and target gene activation [82,147]. The activation of HSF1 is also regulated by binding to heat shock proteins at different phases of the activation process [82]. For example, elevated levels of both Hsp70 and Hsp90 regulate HSF1 through a negative feedback preventing trimer formation during heat shock [148]. Also, Hsp70 and Hsp40 interact with activated HSF1 trimers to inhibit transactivation [148–150].

Redox regulation of transcription factors by H₂O₂

The activity of TF can be regulated by two mechanisms: (i) by synthesis/degradation, where upon activation they are synthesized *de novo* or their stability increases, and (ii) by controlling the activity of a pre-existent TF. There are different mechanisms by which TFs can be activated from an inactive to an active form, most of them mediated by PTMs: state of oligomerization, by binding to a ligand, dissociation from an inhibitory protein, cleavage of a larger precursor, cellular relocation, and access to promoter regions. In principle, activation of a pre-existent TF allows for a faster response to stimuli in order to alter the activity of cellular TFs and produce alterations in gene expression, but, for example, mobilization of a pre-existent mRNA for *de novo* synthesis is also a rapid process [151]. In what follows we analyze how H₂O₂ regulates TFs at each of these levels – synthesis of TF, degradation of TF, cytoplasm–nuclear trafficking, and DNA binding and transactivation – illustrating with representative examples of known regulatory mechanisms.

Protein synthesis of the transcription factor

H₂O₂ regulates several TFs by upregulating their synthesis at the transcriptional, post-transcriptional and translational levels (Fig. 5A). In fact, H₂O₂ increases the rate of transcription of AP-1, TP53 and



Another TF that is activated at the transcriptional level is HIF-1 α . Activation of HIF-1 α transcription by Angiotensin II, in vascular smooth muscle cells, involves the H₂O₂-dependent activation of the phosphatidylinositol 3-kinase (PI3K) pathway [165,166]. The H₂O₂ role in the induction of HIF-1 α transcription might also be mediated by NF- κ B in vascular smooth muscle cells [166]. NF- κ B activates directly HIF-1 α transcription upon recognition of an NF- κ B binding site (at -197/-188 bp upstream of the transcription initiation site) in the HIF-1 α promoter. Both an extracellular bolus addition of H₂O₂ (10–100 μ M) and NOX4 overexpression increase HIF-1 α mRNA levels [166]. Therefore, H₂O₂ acts as a general second messenger for HIF-1-dependent gene expression under normoxia conditions.

There are several studies indicating that TP53 mRNA stability increases in response to cell stress conditions. The 3'-UTR of TP53 mRNA is a target for the RNA-binding protein HUR, which binds and stabilizes TP53 mRNA in response to short-wavelength UV light (UVC) [102]. Exposure of cells to H₂O₂ leads to an increase in cytoplasmic HUR levels [167] and HUR translocates from the nucleus to the cytoplasm and increases IL-6 and IL-8 mRNAs stability [168]. Consequently, it is plausible that H₂O₂ may also regulate TP53 mRNA stability through HUR.

In general, upon exposure to a stress the overall protein synthesis is inhibited, while specific synthesis of stress response factors is upregulated. Protein translation is initiated in eukaryotic cells by two mechanisms: (i) CAP-dependent ribosome scanning, in which the eukaryotic initiation factor 4F complex (eIF4F), recruits the 40S ribosome that scans the 5'-untranslated region (UTR) until it finds the initiation codon AUG; and, (ii) CAP-independent internal ribosome entry mediated by internal ribosomal entry sites (IRESs). Normal physiological protein synthesis is mostly done *via* CAP-dependent mechanisms while protein synthesis of stress response factors is done *via* IRESs, which can induce cells to rapidly produce sufficient amounts of protein in response to the stress [169]. H₂O₂ upregulates both CAP-dependent and CAP-independent translation of TFs (Fig. 5A).

HIF-1 α , increases the rate of translation of Nrf2 and SP1 or may also regulate TP53 mRNA stability.

In addition to the transcriptional activation discussed previously, HIF-1 α expression is also regulated at the translational level. H₂O₂ promotes the activation of a specific kinase for the S6 ribosomal protein, which is a component of 40S ribosomal subunit, p70S6k, increasing HIF-1 α translational rate. This mechanism is induced by insulin and mediated by H₂O₂-dependent activation of MEK/ERK signaling [170].

Concerning NRF2, upregulation of its translation is an important regulatory control exerted by H₂O₂ [171], as a near 50% of maximal response is achieved after 5 min when HeLa cells are exposed to an H₂O₂ steady-state of 12.5 μ M [172]. After applying Eq. (7), this fast response translates into a rate constant of $1.8 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$, for the reaction between H₂O₂ and the target(s) mediating this response. However, if a gradient of 6.8 between extracellular and intracellular H₂O₂ concentrations is considered [42], a rate constant of $1.3 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ is obtained instead. As this estimate is based on NRF2 protein levels, the sensor that triggers this pathway should actually have a higher rate constant, and thus we may speculate that it is a highly reactive H₂O₂ sensor.

The mechanism by which H₂O₂ stimulates NRF2 translation is both cap-dependent and independent [173] (Fig. 5C). CAP-dependent translation may be related to the stimulation of eIF4E phosphorylation at Ser209 by H₂O₂ [173]. The CAP-independent upregulation of NRF2 translation is mediated by an IRES sequence identified within the 5' untranslated region of human NRF2 mRNA containing a highly conserved 18S rRNA binding site (RBS) complementary to the 749–761 bp of human 18S rRNA [173]. In general, IRES activity is regulated by IRES trans-acting factors (ITAFs), localized in the nucleus in the resting state, and which, upon a signal mechanism, translocate to the cytoplasm where they interact with IRES to recruit eIFs and ribosomes to initiate translation [174]. One of such ITAFs, La Autoantigen, was identified as being activated by H₂O₂ in HeLa cells; a treatment with 100 μ M H₂O₂ for 10 min was sufficient to trigger nuclear export of this ITAF, followed by binding to the NRF2 5' UTR and subsequent translation [175]. The mechanism by which H₂O₂ stimulates translocation of La Autoantigen to the cytoplasm is still unknown. Either dephosphorylation of Ser366 [176] or phosphorylation of Thr301 by AKT in mouse glial progenitor cells was reported to promote its cytoplasmic translocation [177], but it was also observed that (de)phosphorylation does not affect the subcellular localization of La Autoantigen [178]. In HeLa cells the phosphorylation status of the La Autoantigen was not observed to be under the influence of H₂O₂ [175].

Another example of H₂O₂-dependent regulation of the translation rate of a TF is SP1 in neurons. H₂O₂ formed extracellularly from D-amino acid oxidase, significantly upregulates both the protein levels and the DNA binding ability of SP1, and of its homologue Sp3, in cortical neurons *in vitro* and *in vivo* [120]. SP1 levels increase significantly 1-, 1.5- and 2-h after the addition of H₂O₂ to neuronal cultures, and H₂O₂ activates the IRES motif present in the 5'-UTR of SP1 mRNA, increasing SP1 levels through enhanced translation of the existing SP1 mRNAs, protecting neurons against ischemic damage [121]. Since SP1 levels auto-regulate its own transcription rate, the high levels of SP1 will lead to a later on increase of SP1 transcriptional rate. It is interesting to note that H₂O₂ activates SP1 translation only in neurons and not in glia cells and it has been proposed that neurons and glial cells probably have different ITAFs that respond differentially to H₂O₂.

The control of translation is a fast-moving area of research and in recent years a lot of information has been obtained. For example, the synthesis of AP-1 is regulated at the translational level, both by cap-dependent and independent mechanisms and by microRNAs [179]. Besides the IRES site present in the 5'-UTR region of NRF2, other regulatory elements are present in NRF2 mRNA. The 3'-UTR region is recognized by microRNAs that repress NRF2 translation [180], while the coding region of NRF2, more specifically in the segment 1159–1815 bp within the 3' portion of the ORF, contains elements that repress NRF2 translation and are responsible for the low-basal

levels of NRF2 synthesis (Fig. 5C) [181]. Finally, IRES sequences have been identified in TP53 [182], YAP1 [183] and c-JUN [179]. Modulation of these modes of control by H₂O₂ is still unknown, but new developments in this area are to be expected in the near future.

Degradation of the transcription factor

Degradation of key modulators of TF function is an important regulatory mechanism in many signaling pathways. The majority of intracellular proteins are degraded via the ubiquitin (Ub)-proteasome pathway, which consists in the degradation of poly-ubiquitinated proteins by a multicatalytic protease called proteasome. Ub-protein ligase (E3) enzymes transfer the activated Ub from an Ub-conjugating protein E2, first to a lysine residue of the target protein and then to lysine residues present in the last added ubiquitin, yielding an Ub chain [184].

Besides the regulation of the general proteasome catalytic activity [185], H₂O₂ is able to modulate the degradation of specific proteins mainly by two mechanisms. In the first mechanism H₂O₂ targets the E3 ligase complex, as are the cases for NRF2 and TP53, and in the second, PTM of the TF either increases (acetylation of AP-1 protein Jra) or decreases (e.g. c-JUN and NRF2 phosphorylation and HIF-1 hydroxylation) its association with the E3 ligase complex (Fig. 6A).

Targeting the E3-complex

Direct targeting The best known example by which H₂O₂ targets the E3-complex involves the factor Kelch-like ECH-associated protein 1 (KEAP1). KEAP1 serves as the substrate adaptor subunit in the E3 holoenzyme in the ubiquitination pathway, leading to NRF2 ubiquitination and degradation [186,187]. In the presence of either electrophilic agents or reactive oxygen species critical cysteine residues in KEAP1 are alkylated or oxidized, leading to a conformational change of KEAP1, due probably to its zinc-finger nature. Such conformational alterations inhibit the binding between NRF2 and KEAP1, thus stopping NRF2 ubiquitination and degradation. There has been an ongoing discussion in the literature whether NRF2 dissociates from oxidized KEAP1 [22]. A recent report based on quantitative fluorescence recovery after photobleaching indicates that such dissociation does not occur [188], and so the NRF2 molecules that translocate to the nucleus are those synthesized *de novo* (Fig. 6B). Human KEAP1 contains 27 cysteine residues, and different agents modify different cysteine residues that translate into specific biological effects [92]. Concerning H₂O₂, in HeLa cells, Cys151 is the critical sensor residue that mediates the formation of an intermolecular disulfide – Cys151–Cys151 – between two KEAP1 molecules [189]. A mutation of this residue impairs NRF2 stabilization in the presence of H₂O₂. In addition, an intramolecular disulfide bond between Cys226 and Cys613 is promoted by H₂O₂, but mutants impairing the formation of this disulfide bond do not show functional alterations [189]. Unfortunately there is no available data concerning the reactivity between KEAP1 cysteine residues and H₂O₂. Based on the data of Fourquet et al. [189], we estimated that the rate constant for this reaction is at $140 \text{ M}^{-1} \text{ s}^{-1}$ (Fig. 3B). There are many assumptions in this estimate, but it is safe to say that cysteine residues in KEAP1 have a relatively low reactivity with H₂O₂.

Mediated targeting Like it happens for NRF2, the cellular TP53 levels are mainly regulated by its ubiquitin-mediated proteasomal degradation [190,191]. In normal cells TP53 is maintained at low levels by interaction with the negative regulator MDM2 [192] a ubiquitin ligase E3 with high specificity for TP53 [193–195]. Regulation of MDM2 activity by H₂O₂ is largely done via PTMs, mainly phosphorylation, that by inhibiting its ubiquitination activity stabilize TP53 (Fig. 6C). The H₂O₂-activated kinases Ataxia telangiectasia mutated (ATM) [196] and c-ABL [197] phosphorylate MDM2 leading to the fast activation of TP53 [198,199].




Fig. 6. Regulation of TFs activity by degradation modulated by H_2O_2 . TF degradation via polyubiquitin (Ub)-proteasome pathway is an important regulatory mechanism in different signaling pathways (A). Proteasome, a multicatalytic protease oligomeric complex, degrades proteins that have been poly-ubiquitinated. Ubiquitin (Ub)-protein ligase (E3) enzymes transfer the activated Ub from an Ub-conjugating protein E2, first to a lysine residue of the substrate protein and the next Ub to lysine residues present in the last added ubiquitin, originating an Ub chain. NRF2 ubiquitination and degradation involves KEAP1 as the substrate adaptor subunit in the E3 holoenzyme (B). In the presence of H_2O_2 critical cysteine residues in KEAP1 are oxidized, changing KEAP1 conformation. This conformational change affects the interaction between NRF2 and KEAP1 inhibiting NRF2 ubiquitination and degradation. In normal cells TP53 is maintained at low levels by interaction with the negative regulator MDM2 a p53 highly specific ubiquitin ligase. H_2O_2 mainly regulates MDM2 activity by activating the ATM and c-ABL kinases involved in MDM2 phosphorylation. MDM2 phosphorylation inhibits its ubiquitination activity and stabilizes TP53 (C). The MDM2 ubiquitin ligase substrate preference for TP53 is enhanced by MDMX. Phosphorylation of MDMX in different residues by c-ABL, AKT and CHK2 kinases inhibit TP53 degradation, while those catalyzed by the CK1 α kinase stimulates TP53 degradation. All the referred kinases activities are regulated by H_2O_2 . The association between the TF and the ubiquitination machinery can also be modulated by PTMs of the transcription factor (D). HIF-1 α is tagged with ubiquitin for degradation after being hydroxylated by PHD in the presence of O_2 (D). H_2O_2 inhibits PHD that leads to HIF-1 α stabilization. Factors colored blue are inhibitors of TF-dependent gene expression; factors colored red are activators of TF-dependent gene expression. Dashed lines indicate activation/inhibition.

A PTM that stabilizes MDM2 and promotes its ability to ubiquitinate TP53 is sumoylation [200]. Sumoylation of MDM2 is downregulated by H_2O_2 through the formation of disulfide bonds between the catalytic cysteine residues of the SUMO E1 subunit Uba2 and the E2-conjugating enzyme Ubc9 [201,202].

The protein MDMX is an additional key partner to the couple TP53/MDM2 and a potential target for H_2O_2 . MDMX is essential for the MDM2-mediated TP53 polyubiquitination [203] as it enhances MDM2 substrate preference towards TP53 [204]. Phosphorylation of MDMX at two residues, Tyr99, catalyzed by c-ABL [205] and Ser367, catalyzed by either AKT [206] or CHK2 [207], activates TP53 by decreasing its MDM2-dependent ubiquitination (Fig. 6C). These PTMs are potentially stimulated by H_2O_2 because H_2O_2 activates the kinases c-ABL [197], AKT [24] and CHK2 [208]. On the other hand, phosphorylation of MDMX at Ser289 catalyzed by casein kinase 1 alpha (CK1 α), stimulates the binding between MDMX and TP53 and leads to an inhibition of TP53 activity. Low levels of H_2O_2 promote the rapid dephosphorylation of CK1 α LS, a nuclear splice form of CK1 α , and enhances the kinase activity [209] (Fig. 6C).

Posttranslational modification of the transcription factor

A more common mechanism by which H_2O_2 stabilizes TFs is, perhaps, mediated by PTMs that modulate the association between the TF and the ubiquitination machinery. Here we describe a few cases involving different PTM of the TFs.

Phosphorylation NRF2 can be tagged for degradation by a KEAP1-independent mechanism that is controlled by glycogen synthase kinase 3 β (GSK-3 β) [210]. Active GSK-3 β catalyzes the phosphorylation of NRF2 in the Neh6 domain, forming a phosphodegron that is recognized by the substrate receptor β -transducin repeat-containing protein (β -TrCP) complex, followed by NRF2 ubiquitination and degradation [211]. GSK-3 β is inhibited by phosphorylation of Ser9 catalyzed by AKT kinase [212]. Thus, H_2O_2 , being an activator of the PI3K/AKT pathway, inhibits GSK-3 β [213,214] activating NRF2 in a KEAP1-independent manner. In fact, inhibition of the PI3K/AKT pathway by LY294002 has been shown to block partially the activation of NRF2 by H_2O_2 [215]. Other kinases that inactivate GSK-3 β are ERK, p38 MAPK and PKC [216], and it has been suggested that these kinases may regulate NRF2 indirectly by inhibiting GSK-3 β [217]. All of these kinases are known to be activated by H_2O_2 , and so they may also

mediate H₂O₂ regulation of NRF2 [218–220]. It should be mentioned that a prolonged exposure to high H₂O₂ concentrations with concomitant cell toxicity activates GSK-3 β [215,221,222], which could be an NRF2 termination signal.

Two additional mechanisms that involve NRF2 phosphorylation causing dissociation of the NRF2/KEAP1 complex, NRF2 stabilization and, ultimately, leading to its nuclear translocation have been identified:

- Phosphorylation catalyzed by protein kinase RNA (PKR)-like ER kinase (PERK), a transmembrane protein kinase that is required for the cellular response to ER stress [223], and that is rapidly activated (15 min) by low extracellular H₂O₂ levels (15 μ M) in HeLa cells [224].
- Phosphorylation at Ser40 catalyzed by PKC- δ [225], a kinase that is well known to be activated by H₂O₂.

H₂O₂ also controls the poly-ubiquitination of the AP-1 members c-JUN and CREB by modulating their PTMs [226,227]. c-JUN phosphorylation, catalyzed either by JNK, at the two clusters Ser63/Ser73 and Thr91/Thr93 [226], or by c-ABL kinase at Tyr170 [228] stabilizes c-JUN, by decreasing c-JUN ubiquitination and proteasomal degradation. Thus H₂O₂ by activating both the apoptosis signal-regulating kinase 1 (ASK1), which activates JNK (see section Modulation of DNA binding by PTM), and c-ABL [197] kinases, may stabilize c-JUN. However, there are conflicting reports concerning the role of Tyr170 in c-JUN stabilization [229]. Concerning CREB, a long-term treatment with a low concentration of H₂O₂ increases CREB-Ser133 phosphorylation and decreases CREB protein abundance via a proteasomal mechanism, but Ser133 phosphorylation is not necessary for CREB degradation [227]. It is possible that either PKD1 (protein kinase D1) regulates CREB phosphorylation at a site other than Ser133 in cardiomyocytes or that the lower levels of CREB are due to the phosphorylation of a different cellular PKD1 substrate able to regulate CREB protein levels indirectly [227].

Hydroxylation Hydroxylation plays a key role controlling HIF activation during hypoxia by increasing HIF-1 α protein degradation. In this system, PHD uses O₂ to catalyze the hydroxylation of the sub-unit HIF-1 α , which is subsequently tagged with ubiquitin for degradation [165,230]. However, the regulatory mechanism leading to HIF-1 activation and stabilization was shown to be more complex than a simple O₂ sensing mechanism. Many other factors were shown to play a role in the regulation of HIF-1 α expression, particularly reactive oxygen species, which have been the center of intense investigation. The molecular mechanism underlining H₂O₂ effect in HIF stabilization has been attributed mainly to an inhibition of PHD activity (Fig. 6D). It has been shown that an increase in the amounts of H₂O₂ due to glucose oxidase addition to the cell medium inhibits hydroxylation of HIF-1 α [231]. Although HIF-1 α has several cysteine residues, their oxidation has never been observed. However, it has been proposed that H₂O₂ is able to oxidize Fe²⁺, present in the catalytic site of PHD, to Fe³⁺ by a Fenton reaction, inactivating the enzyme [232].

In hypoxic conditions, HIF-1 α protein stability and downstream gene target activation were shown to be increased by overexpression of NOX1 [233]. Furthermore, the high levels of mitochondria-derived reactive oxygen species generated by hypoxia were considered to be essential for HIF-1 α stabilization and this effect was reversed upon overexpression of either catalase or glutathione peroxidase 1 [134,234,235]. However, the role of H₂O₂ in HIF-1 α stabilization in hypoxic conditions is still unclear. Recent data has shown that thioredoxin 1 overexpression, although able to block the increase in reactive oxygen species levels under hypoxia, has neither an effect on the levels of HIF-1 α nor in the activation of its downstream targets [236]. Besides H₂O₂, superoxide radical also increases HIF-1 α stabilization and might function as an alternative mechanism [135].

H₂O₂ has also been shown to be an HIF-1 regulator in normoxia. Besides hypoxia, HIF-1 is also upregulated in response to various growth factors, cytokines and hormones in normoxia [237]. The activation of these signaling pathways induces the production of H₂O₂ dependent on the activation of NADPH oxidases that induce HIF-1 α stabilization [238,239].

Acetylation Ubiquitination of Jra, a *Drosophila* Jun protein is facilitated by its acetylation, in a process that is regulated by both Cbp and Sir2 *in vivo* [240]. Because protein acetylation is regulated by H₂O₂ [241], it may be hypothesized that this constitutes another potential pathway for activation of AP-1 by H₂O₂.

Cytoplasm-nuclear trafficking

If the inactive form of a TF rests in the cytoplasm, after activation the TF must be transported from the cytoplasmic pool to the nucleus, where it can affect the transcription of the target genes. There are a few mechanisms by which the accumulation of a TF can be achieved upon exposure to H₂O₂: PTM in the TF itself that either exposes an NLS (mammalian HSF1, Maf1, and Msn2/4) or masks a NES (Yap1, and NRF2); release/association from/with a partner (NF κ B/I κ B, NRF2/KEAP1, and TP53/MDM2). In other cases the TFs precursors are kept in the cytosol associated to membrane anchors; for their activation to occur these factors must be processed, which enables the TFs to enter the nucleus, where the nuclear form activates target genes (SREBP and NOTCH). Below we describe in detail each of these cases where H₂O₂ activates redox-dependent TFs (Fig. 7A).

Activation mediated by post-translational modification in the transcription factor

NLS exposure Mammalian HSF1 provides an interesting example in which trimerization of the TF mediates exposure of an NLS. In non-stressed situations, this TF exists as a monomer and, upon exposure to H₂O₂ forms a trimer, in a reaction involving the formation of disulfide bonds [242]. Redox-regulation of HSF1 multimerization, and its nuclear accumulation are early and linked steps in HSF1 activation. In fact, the mutants HSF1 Cys35Ser, HSF1 Cys105Ser and HSF1 Cys35105Ser, which are defective in stress-induced multimerization *in vivo* and *in vitro*, are also defective for stress-induced nuclear accumulation [242].

Two yeast TFs that in resting conditions are mainly present in the cytosol and whose H₂O₂-dependent activation is mediated by thioredoxins are Maf1 and Msn2/4. Both TFs are controlled by their phosphorylation state and, upon activation, undergo dephosphorylation and translocation to the nucleus. In both cases, rapid nuclear translocation and TF activity are impaired in *trx1* Δ *trx2* Δ cells lacking cytosolic thioredoxins upon exposure to 0.1–0.8 mM extracellular H₂O₂ [243] (Fig. 7B).

The negative regulator Maf1 represses RNA polymerase III activity under carbon source starvation, ER stress (5 mM DTT), and oxidative stress (0.5 mM extracellular H₂O₂) conditions. Maf1 phosphorylation, some of which occurs in the vicinity of an NLS, is mediated by Sch 9 [244], c-AMP-dependent protein kinase A (PKA) [245,246], TORC1 [247] and casein kinase II (CK2) [79]. The cAMP-PKA system is not a key regulator of the H₂O₂-dependent Maf1 nuclear translocation in yeast [243], and it is not known whether the activities of other kinases acting on Maf1 (Sch 9, TORC1 and CK2) are decreased by H₂O₂. What has been established so far is that the activity of protein phosphatase 2A (PP2A) is essential for Maf1 dephosphorylation and nuclear accumulation in cells treated with H₂O₂ [243,248]. This pathway is regulated by oxidized cytosolic thioredoxins because in *trx1* Δ *trx2* Δ cells treated with 0.8 mM extracellular H₂O₂, Maf1 remains in the cytosol and its dephosphorylation is impaired [243] (Fig. 7B).




Fig. 7. Regulation of cytoplasm-nuclear trafficking of TF by H₂O₂. **(A)** Nuclear localization of TF is essential for gene expression activation and H₂O₂ plays a key role in TF cellular trafficking. H₂O₂ modulates NLS exposure by removing PTM or by promoting partner dissociation (P). In certain cases, the association to adaptor proteins might promote NLS exposure. Inversely, NES masking is another mechanism to retain TF in the nucleus that is mediated by H₂O₂. Conformational changes induced by PTM together with the formation of protein complexes make NES inaccessible inducing activation of transcription. Other TF are associated to cellular membranes in their inactive state. The activation of these TF requires proteolytic cleavage and release to the cytoplasm where it is transported to the nucleus. **(B)** Msn2/4 and Maf1 translocation to the nucleus is activated after dephosphorylation, which uncovers NLS. This process is activated by H₂O₂ indirectly through Trx system. Msn2/4 and Maf1 dephosphorylation is dependent upon Trx2 by an unknown mechanism and by PP2A activation, respectively. **(C)** Yap1-dependent gene activation depends on its retention in the nucleus by CRM1 dissociation that occurs in presence of H₂O₂. The oxidation of four Cys residues in Yap1 is responsible for the conformational alterations that prevent NES recognition by CRM1. As for Msn2/4 and Maf1, Yap1 does not react directly with H₂O₂, and its oxidation is mediated by a GPx, Orp1. Trx2 reduces Yap1 inducing its translocation to the cytoplasm, inactivating gene transcription. Factors colored blue are inhibitors of TF-dependent gene expression; factors colored red are activators of TF-dependent gene expression. Dashed lines indicate activation/inhibition.

Msn2 is a direct substrate of protein kinases PKA [245] and Snf1 [249], and most of Msn2 resides in the cytosol when phosphorylated in its NLS. Dephosphorylation of Msn2 may occur due to down-regulation of kinases but the rapid dephosphorylation of Msn2 under glucose depletion suggests that protein phosphatases must also play an important role in the activation of this TF [249], namely PP1 [249] and PP2A [250]. However, contrary to what happens in other stress conditions, for H₂O₂-induced Msn2 nuclear translocation there is no correlation with a decreased PKA activity and translocation is not dependent on PP2A activity [243]. Therefore, the exact mechanism of Msn2/4 activation by H₂O₂ is still unknown, although, like with Maf1 is mediated by thioredoxins. H₂O₂-induced Msn2/4p nuclear localization is not only impaired in *trx1Δtrx2Δ* cells but also in *trr1Δtrx1Δtrx2Δ* cells, which besides lacking cytosolic thioredoxins also have no thioredoxin reductase, and in *tsa1Δ* cells, which lack the most abundant of the yeast Prxs [243]. However, the levels of nuclear Msn2 when H₂O₂ (0.1 mM, 10 min) is added to *trr1Δ* cells, which only lack thioredoxin reductase, are higher than in wild-type cells [243]. These results suggest that H₂O₂-dependent activation of Msn2 involves reaction of H₂O₂ with Tsa1 which is then reduced by the thioredoxin system forming oxidized thioredoxins. How the oxidized thioredoxins transmit the signal to Msn2/4 has not yet been established. However, site directed mutagenesis of the unique Cys residue present in a truncated Msn2 that still exhibits the behavior of the full protein established that there was still thioredoxin-dependent translocation to the nucleus [243]. Therefore, H₂O₂-dependent Msn2 translocation does not involve a direct oxidation of a cysteine residue by the oxidized thioredoxins.

NES masking Yap1 and NRF2 are two TFs that undergo nuclear export mediated by CRM1 (chromosome maintenance region 1). For Yap1 this export is absolutely central for the control of its activation by H₂O₂ and we will describe the process in detail.

Yap1 is found in the cytoplasm under non-stressed conditions but rapidly accumulates in the nucleus following oxidant exposure [75,251]. The cytoplasmic localization is determined by constitutive nuclear export that predominates over constitutive nuclear import [75,252,253]. The activation of Yap1 by H₂O₂ requires the multistep formation of disulfide bonds in the CRD and/or the cCRD domains [254]. The intramolecular disulfide bond between Cys303 (cCRD) and Cys598 (nCRD) formed in Yap1 after H₂O₂ exposure masks the NES present in the cCRD domain, resulting in structural changes in the NES [76]. This allows the Yap1–Crm1 complex to be disrupted and Yap1 accumulates in the nucleus [252,255,251]. It is important to note that, depending on the oxidative stress inducing agent, Yap1 can undergo two distinct conformational changes, both involving disulfide bond formation, and both masking the nuclear export signal, thus abolishing nuclear export by Crm1 [76]. In the case of H₂O₂ cysteine residues in Yap1 are not oxidized by a direct reaction with this agent but require GPx3/Orp1, which acts as sensor protein for H₂O₂ [256]. GPx3/Orp1 has *in vitro* peroxidase activity by a mechanism involving a Cys36–Cys82 catalytic disulfide, which is distinct from those of classical glutathione peroxidases. Also, Orp1 is recycled by thioredoxin and not by GSH. GPx3/Orp1 becomes oxidized at Cys36 which forms a heterodimeric disulfide bond with Cys598 in Yap1 upon H₂O₂ exposure [256]. Two more thiol/disulfide exchange reactions finally result in oxidized Yap1, with formation of a disulfide bond between Cys303 and Cys598, and reduced GPx3/Orp1. The second disulfide bond in Yap1 (Cys310–Cys629) presumably forms through a similar pathway on recruitment of another oxidized GPx3/Orp1 molecule, only after the first disulfide bond has formed [257]. GPx3/Orp1 is reduced by the thioredoxin/thioredoxin reductase system [256]. Yap1 nuclear export is restored when disulfide bonds are reduced by Trx2, whose expression is controlled by Yap1, providing a mechanism for negative

autoregulation [256,251]. Besides GPx3/Orp1, there is another protein, Ybp1, which forms a stress-induced complex with Yap1 *in vivo* and influences the nuclear localization of Yap1 in response to H₂O₂ [258,259]. It has been proposed that the role of Ybp1 might be to increase the efficiency of this process *in vivo* to more rapidly prepare a defensive response to an H₂O₂ challenge (see Fig. 7C).

Concerning NRF2 export from the nucleus, NRF2 possesses both a redox-sensitive [260] and a redox-insensitive [261] NES but, ultimately, both mediate H₂O₂ regulation. The redox-sensitive site is present in the Neh5 transactivation domain of NRF2, where the redox state of Cys183 has a critical role. Mutation of this residue abolishes the rapid activation of NRF2 by 50 μ M extracellular H₂O₂ in HeLa cells, which led to the conclusion that oxidation of Cys183 inhibits binding of NRF2 to the nuclear exporting protein CRM1 [260]. The redox-insensitive NES is present in the leucine zipper (ZIP) domain of NRF2, and phosphorylation of Tyr568 residue present in its vicinity, strengthens interaction with CRM1 to enhance nuclear export of NRF2 [261,262]. The kinase responsible for this PTM is FYN, which is activated by GSK-3 β . [221]. Thus, prolonged exposure to high H₂O₂ concentrations, with concomitant cell toxicity [222], activates GSK-3 β , induces NRF2 phosphorylation at Tyr568 and NRF2 degradation [215,221]. On the other hand, short-time incubation with H₂O₂ inhibits GSK-3 β [213,214], activating NRF2. It has been proposed that phosphorylation at Tyr 568 may mediate a failed antioxidant response, leading to cellular toxicity, and eventually cell death [215] or, alternatively, it may constitute a switch-off mechanism [221]. An additional mechanism by which NRF2 export is controlled is by its binding to small MAF bZIP proteins, which enhances nuclear retention of NRF2 by masking the NES motif in the NRF2 ZIP domain [90].

Activation mediated by association/dissociation from a partner

NF- κ B is a TF whose central activation step involves its release from an inhibitory protein, I κ B, that masks an NLS in NF- κ B. Concerning the most abundant form of NF- κ B, the dimer p65–p50, H₂O₂ is considered a modulator of activation induced by other agents, namely cytokines, and not an inducer *per se* [22,115,263]. Nevertheless, recent evidence indicates that in MCF-7 cells H₂O₂ (25 μ M steady-state) induces c-REL nuclear translocation to values similar to those obtained with physiological levels of TNF- α [42]. The mechanisms by which H₂O₂ promotes degradation of I κ B, thus exposing the NLS of NF- κ B and subsequent NF- κ B activation are complex and are not well defined. I κ B is signaled for proteasome degradation following its phosphorylation catalyzed by the IKKs. The IKK complex consists of a dimer of IKK γ subunits linked through a disulfide bond forming the NF- κ B essential modulator (NEMO) to which IKK α and IKK β bind in the resting state. Activation of IKK is made by phosphorylation catalyzed by an IKK kinase, either TGF β -activated kinase (TAK1) or AKT, in the canonical pathway of activation, or NF- κ B inducing kinase-1 (NIK1) in the alternative pathway, which can all be modulated by H₂O₂ [115,22]. Alternatively, LC8, a substrate of the redoxin TRP 14, may mediate H₂O₂ effects. In the reduced form LC8 binds to I κ B and inhibits its phosphorylation by IKKs and subsequent degradation, and thus H₂O₂ by forming LC8 dimers linked via a disulfide bond [264], promotes LC8 dissociation from I κ B, and NF- κ B activation [265]. On the other hand, inhibition of activation of NF- κ B by H₂O₂ could be mediated by KEAP1-dependent degradation of IKK β [266].

Concerning NRF2, in addition to the mechanism previously described, its cellular localization is also mediated by interaction with its partner, KEAP1. NRF2 contains three NLS, one of which is present in the NEH2 domain of NRF2, the domain that mediates interaction of NRF2 with KEAP1 [267,268]. So, NRF2 that is not bound to KEAP1 tends to translocate to the nucleus [269]. NRF2 can also be exported from the nucleus bound to KEAP1, which binds to CRM1 through a NES [270], being subsequently degraded in the cytosol [271]. KEAP1 does translocate to the nucleus by an unknown mechanism [272], possibly by inhibition of its export from the nucleus. It has been observed

that both mutation of KEAP1 Tyr85 [273] and oxidative modification within an NES of nuclear KEAP1 blocks NRF2-KEAP1 complex export from the nucleus [270].

In the case of TP53, there are several evidences showing that H₂O₂ promotes TP53 nuclear localization [157,274] but the mechanisms involved are not clearly established. The MDM2 protein, which acts in conjunction with CRM1, mainly regulates the TP53 nuclear export through NES [108,275]. Consequently, factors that regulate the specific cellular compartment localization of MDM2 also control TP53 cellular localization. For example, the p14ARF protein causes MDM2 nucleolus permanence, and c-Abl binds to MDM2 preventing its interaction with TP53, both leading to nuclear retention of TP53 [199,276,277]. On the other hand the phosphorylation of MDM2 through the PI3K/AKT pathway promotes MDM2 nuclear import, resulting in TP53 nuclear export [278–280]. Interestingly, HSF1 is required for TP53 nuclear importation and activation, which implies that heat shock factors play a role in the regulation of TP53. As already mentioned MDM2 is regulated by H₂O₂ and, therefore, the nuclear localization of TP53 induced by H₂O₂ may be mediated by MDM2.

The cytoskeleton may also play an important role as a partner in TP53 translocation. TP53 phosphorylated forms, namely at Ser15, show increased binding to microtubules and a lower tendency to nuclear translocation after exposure to chemotherapeutic agents [281]. It is conceivable that H₂O₂ may affect TP53 microtubule binding and nuclear translocation through alterations on TP53 phosphorylation status since several reports describe the phosphorylation of Ser15 in TP53 upon H₂O₂ exposure (see Table 2).

Activation through release from membrane by proteolytic processing

For both NOTCH and SREBP1c the critical activating event is their proteolytic processing and release from the plasma membrane and the ER, respectively. These proteolytic events may be controlled by H₂O₂, activating NOTCH signaling and having cell-type dependent effects for SREBP1c.

In the case of NOTCH signaling, it was observed that colonic tissues of NOX1^{KO} mice present less NICD compared to wild-type animals and a consequent decrease in the activation of NOTCH target genes [282]. The increase of NICD proteolysis by H₂O₂ was also observed for NOTCH3 in human aortic endothelial cells treated with extracellular H₂O₂ (50 μ M, 1 h) or, inversely, with extracellular catalase added to the cultures [283]. These authors also showed a concomitant decrease in NECD in the presence of H₂O₂. ADAM17/TACE, the metalloproteinase responsible for the NECD release, has been demonstrated to be activated by H₂O₂ through the oxidation of critical sulfhydryls of the protein, suggesting its involvement in the redox regulation of NOTCH signaling [284,285].

Concerning SREBP1c, this TF is predominantly produced in the liver and white adipocytes, and controls fatty acid biosynthesis under glucose/insulin stimulation [138,286]. In HepG2 cells, extracellular H₂O₂ (100, 250, and 500 μ M for 3 h) stimulates expression of SREBP-1 gene as well as its target genes involved in lipid accumulation [139], while in COS-7 cells H₂O₂ suppresses SREBP-1c transcriptional activity. The stimulatory effect of H₂O₂ may be mediated by the insulin signaling pathway at least in part through the oxidative inhibition of protein tyrosine phosphatases [286]. It is also possible that H₂O₂ activates SREBP by inducing ER stress [287], producing the mature forms of the SREBP-1c that translocate to the nucleus. It is important to note the different response to H₂O₂ observed between the two types of cells, HepG2 and COS-7 cells. Such differences may be due to a difference in nutrient metabolic function, which could cause differences in SREBP-1c needs, highlighting the specialized role of this TF.

Table 2Phosphorylated and acetylated TP53 amino acid residues in different cell types exposed to extracellular single doses of H₂O₂.

Cell type	[H ₂ O ₂] (μM)	Exposure time	Domain	Phosphorylation	Kinases	Biological role	References
GM00637	200		N-terminal domain (transactivation)	Ser9, Ser15, Ser20	Ser20 residue requires Plk3 activated by ATM	Induction of p21, suggesting functional activation of p53	[334]
A549	50, 100	30 min	C-terminal domain (regulatory)	Ser15, Ser20, Ser37 or Ser392	ATM-dependent except for Ser37 and Ser392		[368]
HUVEC	100	5 min to 1 h	N-terminal domain (transactivation)	specifically at Ser15	ATM kinase (downstream PDGF receptor)	Increased transcription of p21 ^{CIP/WAF} gene	[333]
NHF	100	1 h	N-terminal domain (transactivation)	Thr81	JNK	P53 stabilization and activation of transcriptional activities	[369]
Human keratinocytes	50, 100	1 h or 16 h	N-terminal domain (transactivation)	Ser15?	AMPK?	Senescence in human keratinocytes by activating p21 ^{CIP1}	[370]
AF5	400	2 h, 6 h, 12 h	N-terminal domain (transactivation)	Ser15	?	Induction of P21, MDM2 and apoptotic proteins	[160]
TIG-3	150	1 h	C-terminal domain (regulatory)	Acetylation Lys373 and 382	Deacetylases Downregulation of SIRT1	Cell senescence; induction of p21	[344]
HUVEC	0, 25, 50, 100	6 days	C-terminal domain (regulatory)	Lys382	Downregulation of SIRT1	Unaltered levels of P53, P21, Bcl-2, and Bax; Up-regulation of p16INK4a; G1 arrest and aging	[371]
Human skin keratinocytes	50, 125, 250	0.5 and 2 h	C-terminal domain (Regulatory)	Lys382	Downregulation of SIRT1 involving JNK activation	Induced cell death	[343]

DNA binding and transactivation

In the chain of events we have been describing transactivation and DNA binding represent the last step in order for a gene to be activated by a TF. In the simplest case, the TF senses directly H₂O₂, changes its conformation, and as such it increases (e.g. OxyR and mammalian HSF1) or decreases (PerR) its affinity towards DNA. Many mammalian TFs contain cysteine residues in their DNA binding domain whose oxidation impairs DNA binding, constituting an additional layer of control in a complex signaling pathway. In addition, H₂O₂ modulation mediated by PTM of the TF can modulate the affinity of the TF either towards DNA (c-JUN, SP-1, CREB, and NRF2), towards a co-activator (HIF-1) or towards a repressor (c-JUN, Yeast Hsf). The state of chromatin adds another layer of control and, actually, H₂O₂ is able to target specific regions of chromatin to selectively activate individual genes, as exemplified for SP1 (Fig. 8A).

Modulation of DNA binding by oxidation or reduction

Oxidation of the transcription factor In bacterial TFs and in the TF responsible for mammalian general stress response, HSF1, the signaling pathway boils down to controlling binding/dissociation from DNA. In these examples the amino acid residues or prosthetic groups that react with H₂O₂ mediate affinity towards DNA.

Cysteine oxidation. In *E. coli*, OxyR functions as a sensor of H₂O₂ and a transducer of oxidative stress due to its capacity to be oxidized directly [61,67]. Different studies have been done in order to understand the mechanism underlying oxidant sensing by OxyR (see review [288]). Two basic models have been proposed: (i) in the first

model, reduced OxyR reacts directly with H₂O₂ to form a sulfenic acid at Cys199, which rapidly forms a disulfide bond with Cys208. The formation of this specific disulfide bond induces structural changes in the regulatory domain leading to altered DNA binding properties and allowing a productive interaction between OxyR and RNA polymerase, which then triggers expression of the target genes. OxyR is rapidly deactivated by slow reduction *via* a glutathione-dependent mechanism [23,289–291]; (ii) in the second model, the “molecular code” hypothesis, modification of Cys199 alone is sufficient to activate OxyR in the absence of disulfide bond formation; several PTMs can occur in Cys199, including sulfonylation (Cys199-SOH), S-nitrosylation (Cys199-SNO), and S-glutathionylation (Cys199-S-S-G) forming a modified OxyR able to activate the transcription of target genes [292]; although all these modifications lead to a transcriptionally active OxyR, they differ in structure, cooperative properties, DNA binding affinity, and promoter activities. By this second mechanism, the authors suggest a redox “molecular code” resulting in a more selective regulatory response in which distinct subsets of OxyR target genes are differentially regulated depending on the particular oxidized form of OxyR involved. Recent data shows that Cys199 nitrosylation is associated with the activation of a sub-set of genes that differ from those upregulated in response to H₂O₂ [293], which supports the “molecular code” mechanism. Probably both mechanisms operate *in vivo* [67].

Mammalian HSF1 can be activated directly by H₂O₂ [242]. The initial step in HSF1 activation is its trimerization. The trimerization of HSF1 increases the affinity of HSF1 for HSEs by several orders of magnitude [294]. Multimerization of monomeric recombinant mouse Hsf1 and Hsf1 DNA binding is induced *in vitro* by heat shock temperatures and by extracellular H₂O₂ (0.1 μM–200 μM) [242]. Ahn and




Fig. 8. Regulation of DNA binding and transactivation of TF by H₂O₂. **(A)** TF affinity to DNA or to transcription activators is regulated by H₂O₂ through different mechanisms involving either conformational alterations induced by PTM or association with activators/inhibitors. In addition, H₂O₂ is able to modulate TF activation by direct TF oxidation inside or outside DNA binding domain. **(B)** PerR DNA affinity is lost by a conformational alteration induced by a Fenton reaction of H₂O₂ with Fe²⁺ forming 2-oxo-histidine. Both Fe²⁺ and Mn²⁺ (black spheres) can bind to the regulatory site but when Mn²⁺ is bound 2-oxo-histidine is not formed. Structural ZnII ions (yellow spheres). **(C)** Ref-1 induces the DNA binding activity of several TFs by reducing cysteine residues of those TFs. Ref-1 is regenerated by thioredoxin that reduces Cys65. **(D)** H₂O₂ promotes c-Jun phosphorylation releasing its repressor complex, which contains HDAC3. This phosphorylation is dependent of the activity of JNK, which is activated by ASK1. ASK1 activation is regulated by H₂O₂ through three alternative mechanisms: GST oxidation releasing ASK1, Trx oxidation releasing ASK1 or ASK1 oxidation by PRX1. Factors colored red are activators of TF-dependent gene expression; factors colored blue are inhibitors of TF-dependent gene expression. Dashed lines indicate activation/inhibition.

Thiele [242] were able to show, using site directed mutagenesis, that *in vivo* activation of Hsf1 by extracellular H₂O₂ (500 μM, 1 h) requires the formation of a disulfide bond between the two cysteine residues (Cys35 and Cys105) present in the Hsf1 DNA-binding domain. Similarly to what happens for murine Hsf1, H₂O₂, diamide and heat stress also induce the formation *in vitro* of an intramolecular disulfide bond between the Cys36 and Cys103 residues in the DNA-binding domain of human HSF1, which leads to HSF1 trimerization and DNA binding [295]. However, for human HSF1 it was found that another disulfide bond involves other cysteine residues (Cys153, Cys373 and Cys378) located in the trimerization and in the transcriptional activation domains, and their oxidation leads to inhibition of HSF1 trimerization and DNA binding [295].

Metal center oxidation. PerR provides a very interesting example in which H₂O₂ sensing is mediated by Fenton chemistry and not cysteine oxidation. Because this is a less common mechanism for H₂O₂ signaling effects, we will discuss in some detail the mechanistic and structural aspects of the modification of this TF by H₂O₂. This TF is a transcription repressor and its oxidation releases it from the promoter

regions of the target genes, thus activating gene expression. The crystallographic structures of PerR from *B. subtilis* [296–298] and *Streptococcus pyogenes* [299] have helped to gain insight into the mechanism of PerR as an H₂O₂ sensor and also of how Fe²⁺ binding to the protein leads to activation of PerR binding to DNA. Just like the other proteins of the Fur family, PerR is a homodimer and each monomer has two functional domains, an N-terminal DNA binding domain that contains a winged helix interaction motif and a C-terminal domain that is involved in the dimerization of the protein [296]. Each monomer also contains two metal-binding sites, a structural binding site with tightly bound Zn²⁺ coordinated by four cysteine residues [300] and a regulatory site, which under physiological conditions can be loosely occupied either by Fe²⁺ (PerR: Zn,Fe) or Mn²⁺ (PerR: Zn,Mn), and that is required for DNA binding [63]. The Zn(Cys)₄ site gives structural stability to the dimer by locking together three β-strands present in the dimerization domain [66,296,300,301]. Insights into the structure of the regulatory site of PerR emerged both from mutational analyses [66] and from structural modeling based on the PerR apoprotein (PerR: Zn) [296]. Both studies suggest that the five ligands for the metal in the regulatory site, all conserved within PerR family members, consist of three histidine residues (His37, His91, and His93) and two aspartate residues (Asp85 and Asp104).

Lee and Helmann [300] estimated a rate constant of ~10⁵ M⁻¹ s⁻¹

for the inactivation of PerR: Zn,Fe by H_2O_2 , which is comparable to that for the oxidation of OxyR by H_2O_2 [291], and showed that PerR senses H_2O_2 at its regulatory site through metal-catalyzed histidine oxidation. The mechanism for PerR oxidation by H_2O_2 would rely on the production of a hydroxyl radical through the Fenton reaction, involving the bound Fe^{2+} , leading to the rapid and direct incorporation of an oxygen atom into one of the histidine ligands (His37 or His91) forming 2-oxo-histidine [66]. The formation of 2-oxo-histidine has also been reported for the inactivation of Cu,Zn-superoxide dismutase [302] and in metal-catalyzed oxidations of histidine-containing enzymes [303]. Oxidation of His37 is the preferred site of 2-oxo-histidine formation and does not lead to changes in the affinity for the metal ion bound to the regulatory site. On the contrary, when His91 is oxidized PerR has a considerably reduced affinity for the regulatory site metal ion [297]. Oxidation of His37 and/or His91 in PerR prevents DNA binding due to a conformational change in PerR and the inactivated oxidized protein dissociates from the DNA, which leads to a derepression of the PerR regulon [66,297] (Fig. 8B). No repair mechanisms are known for 2-oxo-histidine, which has led to the proposal that oxidized PerR is not recycled [66].

The metal content of the regulatory site determines the ability of PerR to sense H_2O_2 [63,69,299]. While the PerR: Zn,Fe protein is highly sensitive to H_2O_2 , peroxide induction of the PerR regulon through oxidation and inactivation of PerR is inefficient with the PerR: Zn,Mn protein. *In vivo* inactivation of PerR: Zn,Mn requires 10 mM extracellular H_2O_2 , a level at least 1000 times higher than that needed for inactivation of PerR: Zn,Fe [300]. Fe^{2+} has an approximate 30-fold higher affinity for the regulatory site than Mn^{2+} and *in vivo* there is a competition between Fe^{2+} and Mn^{2+} for binding to the regulatory site [67]. Thus it is possible that *in vivo* PerR is regulated both by H_2O_2 levels and the relative levels of Fe^{2+} and Mn^{2+} . So, when Fe^{2+} levels are low and Mn^{2+} levels are high the PerR regulon is not derepressed upon exposure to H_2O_2 [304].

Reduction of the transcription factor by Ref1 and thioredoxin There are several TFs that contain cysteine residues that must be reduced in order to bind DNA, although *per se* this is not sufficient for TF activation. To what extent H_2O_2 oxidizes these cysteine residues *in vivo* to inhibit DNA binding is unknown. For example, the DNA-binding activity of SP1, which is strongly decreased by exposing nuclear protein extracts from 4-month-old rat tissues for 1 h with 5 mM–20 mM extracellular H_2O_2 , is restored by the treatment with high dithiothreitol concentrations [305]. Concerning NF- κ B, 1 mM extracellular H_2O_2 , although able to induce NF- κ B translocation to the nucleus in human endothelial cells, does not induce NF- κ B-dependent transactivation of DNA [306].

Cells have a nuclear enzymatic system to keep cysteine residues in TFs reduced, the apurinic/aprimidinic endonuclease/redox effector factor-1 (APE/REF-1) [307]. REF-1 was first recognized by its DNA base excision repair activity and, therefore, it was initially named apurinic/aprimidinic endonuclease [308]. It is the rate limiting enzyme in the mammalian base excision repair pathway [309]. However, in addition to its DNA repair function, REF-1 also displays, in mammals but not in other vertebrates, a distinct redox regulation role as a redox co-activator of different TFs [307,310]. The two biological activities displayed by REF-1 are located in two functionally distinct domains. The redox activity is associated with the Cys65 residue located in N-terminal domain [311], which contains the NLS region, while the enzymatic activity on the abasic sites of DNA is associated with the C-terminal domain [312] (Fig. 8C). REF-1 induces the DNA binding activity of several TFs including AP-1 [307], ATF/CREB family [307], MYB [307], EGR-1 [313], HIF-1 α [314], NF- κ B [315], PEBP2 [316], TP53 [317], HLF [318], NF- κ B [319], and PAX proteins [320] by catalyzing the reduction of cysteine residues of those TFs.

Thioredoxin also plays a role in keeping critical Cys residues in TFs

reduced. In fact, overexpression of thioredoxin reverses the oxidation of the Cys62 residue in p50 NF- κ B subunit [321], and endogenous thioredoxin is responsible for the maintenance of reducing conditions within the nucleus, so that NF- κ B is able to bind to DNA [322]. Furthermore, mutating Cys35Ser in nuclear thioredoxin-1 inhibits NRF2 transcriptional activity and DNA binding [323], which requires reduction of NRF2 Cys506 residue present in the DNA binding domain [298]. Alternatively, thioredoxin may also work as a REF-1 partner because it reduces the oxidized Cys65 of REF-1 [324].

Finally, cysteine oxidation allows discriminating between target sequences in distinct genes. Oxidation of Cys277 in TP53 caused a decrease in TP53 binding to the GADD45, involved in DNA-repair, but not to the p21^(WAF1/CIP1) responding element [325].

Modulation of DNA binding by PTM

TP53 activation is the most representative example of how a complex network of PTMs, namely phosphorylation, acetylation, ADP-ribosylation, ubiquitylation, sumoylation and neddylation [195,327–331] modulate binding to DNA and TF activity. Most of these modifications occur in the N- and C-terminal transactivation and regulatory domains of TP53, respectively. From the vast information on the literature it seems that the ability of TP53 to specifically select a target gene is closely related to the distinct combinations between PTMs of the TP53 at N-terminal and C-terminal domains like phosphorylation and acetylation, which allows cells to cope and respond to diverse cellular stresses [327,330,332]. H_2O_2 modulation of these PTMs further adds a layer of complexity and can contribute for selective gene expression by H_2O_2 .

Phosphorylation of the TF Concerning TP53, upon stresses the majority of the available sites are rapidly phosphorylated, although some sites (e.g. Thr55, Ser376 and Ser378) are constitutively phosphorylated in cells and dephosphorylated in response to stress conditions [327,329]. TP53 can be modified by phosphorylation by a broad range of kinases, including ATM/ATR/DNA-PK, and CHK1/CHK2 [329,333]. Concerning exposure of cells to H_2O_2 it seems that H_2O_2 is able to induce different pathways that lead to diverse TP53 phosphorylation patterns, which is most probably related with used dosages and cell types as summarized in Table 2, but in all the cases leads to transcription of TP53-target genes. In most cases ATM activity is required for these phosphorylations either directly [334,335] or indirectly by other ATM-activated kinases, like PLK3 (Polo-like kinase-3) [335].

Three examples in which phosphorylation of the TF inhibits DNA binding and TF activation are HSF1, SP1 and c-JUN. Concerning HSF1 and SP1 phosphorylation, the kinase involved is JNK [336,337]. For c-JUN, phosphorylation in its C-terminal DNA-binding domain at Thr239 is catalyzed by GSK-3 β , while phosphorylation at Ser243 is catalyzed by an unknown protein kinase [338]. This is relevant since the phosphorylation of c-JUN catalyzed by GSK-3 contributes to the repression of growth factor-inducible genes in quiescent cells [339].

Finally, concerning CREB, this TF is regulated *via* phosphorylation at the Ser133 residue present in the P-box of the activation domain. This PTM is attributed to different kinases, depending upon the cell type or the stimulus [95]. H_2O_2 can either increase or decrease binding, depending on the cell type. For example, H_2O_2 decreases the DNA-binding activity of CREB to the CRE domain in neuronal cells, contrary to what happens in glia cells, where H_2O_2 increases the DNA-binding activity of CREB [340]. In immortalized hippocampal neuronal cells (H19-7 cells), a CREB-dependent decrease of both Bcl-2 protein and mRNA levels follows exposure to extracellular H_2O_2 (150 μM) [341]. These observations are accompanied by a decrease of CREB phosphorylation. When Bcl-2 is overexpressed in H19-7 cells, these cells are more resistant to apoptosis induced by reactive oxygen species. The loss of CREB function, which is essential for neuronal survival, contributes to oxidative stress-induced neuronal dysfunction.

It is predicted that a low level of CREB induced by H_2O_2 may influence the apoptosis responses in cells.

Acetylation of the TF In general, H_2O_2 increases acetylation of TP53 [342] (see Table 2) and NRF2 [343], and in both cases this correlates with binding to DNA and TF activity. As in the case of NRF2 [343], in most of cases where increased levels of acetylated TP53 induced by H_2O_2 were found, a simultaneous decrease in the activity of the deacetylase Sirtuin 1 (SIRT1) also occurs, but the upregulation of target genes can vary in different cell types [344,345]. It should be mentioned that SIRT1 is extremely sensitive to H_2O_2 inhibition, since extracellular concentrations as low as 1 μ M inhibit SIRT1 by oxidizing critical cysteine residues in the SIRT1 active center [241]. In addition, the RNA-binding protein HUR binds to the 3' untranslated region of the mRNA encoding SIRT1, leading to its stabilization and increased levels. H_2O_2 triggers the dissociation of HUR from the HUR-SIRT1 mRNA complex, promoting SIRT1 mRNA decay, reducing SIRT1 abundance, a process that seems to be regulated by Chk2 kinase [167]. Other TFs also targeted by SIRT1 with the concomitant inhibition of their transcriptional activity are c-JUN, c-FOS and HIF-1 α . It is thus tempting to hypothesize that acetylation of these TFs is modulated by H_2O_2 . SIRT1 may function as a termination pathway because REF-1 was found to reduce SIRT1 cysteine residues, stimulating its activity [241].

Other PTMs compete with acetylation for C-terminal TP53 lysine residues like ubiquitination, methylation, sumoylation, and neddylation [327], which suggests a cross-talk between the different modifications at each specific modifiable residue creating the possibility of a wide variety of patterns of distinct cellular responses to different signals. How H_2O_2 affects these PTMs is still unknown but H_2O_2 may regulate either the enzymes involved in their addition or removal. Actually, H_2O_2 has been implicated as a key regulator of the sumoylation–desumoylation equilibrium by affecting the SUMO conjugating enzymes E1 and E2 [201,202] and SUMO proteases [346].

Binding to a co-activator

Binding to a co-activator is a common mechanism of regulation. Regulation at this level by H_2O_2 is illustrated by the hydroxylation of HIF. HIF-1 α and HIF-2 α contain a conserved asparagyl residue that is hydroxylated by factor inhibiting HIF (FIH) enzymes (Asn803 in HIF1 α ; Asn851 in HIF-2 α for human proteins). This hydroxylation inhibits the association with p300, a transcriptional co-activator. It has been shown that FIH is more sensitive to inhibition by H_2O_2 than PHD and actually FIH may represent the preferential target for H_2O_2 -dependent regulation of HIF [347]. In the case of NRF2, mutations at Cys119, Cys235, and Cys506 prevent binding of NRF2 to p300 [348].

Release from inhibitor

c-JUN provides a very interesting example in which H_2O_2 control is done not by promoting the binding of the TF with a co-activator, but instead it promotes the release of the TF from an inhibitor. Induction of c-JUN is mostly dependent on its upstream kinase, JNK. Once activated, JNK translocates to the nucleus [349] where it catalyzes the phosphorylation of c-JUN at Ser63, Ser73, Thr91 and Thr93 [338,350], and regulates its transcriptional capacity by dissociating c-JUN from a repressor complex that contains HDAC3 [351]. After release from this complex, gene activation, which involves interactions with transcriptional co-activators like histone acetyl transferases, CREB, Smad3 and CBP, can occur in the absence of functional JNK [351] (Fig. 8D). H_2O_2 is at the center of these events because JNK is activated by ASK1 [352], and ASK1 is extremely sensitive to H_2O_2 by several mechanisms. ASK1 contains 5–6 cysteine residues that, upon exposure to H_2O_2 , form intermolecular disulfide bonds that mediate ASK1 homo-oligomerization and kinase activity [353]. Thioredoxin-1 reduces these disulfide bonds inhibiting the activation of ASK1. ASK1

is probably not a direct sensor of H_2O_2 , with its oxidation being mediated by Prx1, which acts as the direct sensor of H_2O_2 [29]. An alternative mechanism states that thioredoxin-1 forms an inactive complex with ASK1; upon exposure to H_2O_2 , thioredoxin is oxidized dissociating the complex, allowing for ASK1 activation [354]. Thioredoxin may be recycled by interaction with REF-1 [324]. Similarly, dissociation of the GST (glutathione S-transferase)-ASK1 complex following GSTp oligomerization promoted by H_2O_2 has been proposed to mediate JNK activation [355]. Following oxidation, ASK1 is activated by phosphorylation [353]. Thus, inhibition of serine/threonine protein phosphatase 5 (PP5), an ASK1 phosphatase, has been also proposed as a mechanism for ASK1 activation by H_2O_2 [356]. Besides the ASK1/JNK axis, H_2O_2 causes oxidation and inhibition of JNK-inactivating phosphatases by oxidizing their catalytic cysteine, thus contributing to JNK activation [357,358].

Another example of TF activation by release from an inhibitor is provided by the Hsf1 yeast counterpart of HSF1. Activation of Hsf1 by H_2O_2 in yeast does not involve, apparently, a direct reaction of H_2O_2 with the TF, since it exists as a trimer in the nucleus already bound to DNA. Instead, recent evidence suggests that H_2O_2 may react with the Hsp70 chaperone Ssa1, which interacts with the transactivation domain of human HSF1 repressing it in the absence of stress [148–150,359]. In fact, deletion of SSA1 and SSA2 genes, which code for the two cytoplasmic isoforms of Hsp70 in *S. cerevisiae*, leads to derepression of Hsf1 transcriptional activity [360]. Wang et al. [359] found that activation of Hsf1 in yeast by H_2O_2 probably involves as a direct target the oxidation of two cysteine residues in Ssa1. In this study H_2O_2 oxidation of Ssa1 was mimicked by substituting Cys264 and Cys303 with aspartic acid, which resembles a cysteine sulfinic acid, and this led to a non-functional Ssa1 and to constitutive Hsf1 activation.

Chromatin state

An illustration on how H_2O_2 can affect the state of chromatin locally near the promoter region of specific genes is exemplified by the methylation state of SP1. Recently, SP1 was reported to be specifically methylated *in vivo*, which suppresses its transcriptional activity [361]. In fact, H_2O_2 treatments increased the methylation of SP1 and enhanced the interaction between SP1 and HDAC1 in a dose- and time-dependent manner. The experiments were done with HeLa cells treated with 400 μ M H_2O_2 until 6 h or with 1 mM H_2O_2 for 2 h. The negative role induced by the methylation of SP1 is a result of the direct interaction with the protein lysine methyltransferase Suv39H1 (Su(var) 3–9 homologue 1), which allows SP1 to recruit HDAC1 to the promoter region of its target genes. After SP1 methylation associated with Suv39H1 and HDAC1, the chromatin is remodeled with histone deacetylation and histone methylation, culminating in repression of SP1 target genes expression. The authors suggest that the recruitment of HDAC1 is essential for SP1 methylation; however, the enzyme that provides the methyltransferase activity and the residue(s) that are methylated are still unknown [361].

Selective gene expression of individual genes whose promoter regions are exposed may also be accomplished by H_2O_2 based on the affinity of promoter regions towards the TF. In the case of NF- κ B, simple association–dissociation equilibria helps to understand why genes with lower affinity towards NF- κ B are preferentially modulated by H_2O_2 rather than those with higher affinity [164]. Understanding how H_2O_2 not just regulates a TF, but also how it selectively discriminates between genes responding to the same TF is essential to have a complete view of regulation of gene expression by H_2O_2 .

Conclusions

The complexity of the regulation of TFs by H_2O_2 increases from bacteria to unicellular eukaryotes, and from these to multicellular organisms. One consequence is that while the regulatory mechanisms

in bacteria and yeast are well defined, a lot of uncertainties still remain in mammalian cells. As noted in a recent review, while in lower organisms a few publications were sufficient to establish the regulatory pathways, in mammalian systems thousands of publications have failed to do so [22]. One example that illustrates the impact of this higher regulatory complexity is NRF2. In an excellent work, the cysteine residues of KEAP1 oxidized by H_2O_2 were identified [189]. If NRF2 was to be activated just by a simple oxidation of KEAP1, then the regulatory pathway would be solved. Instead, NRF2 is also regulated by synthesis *de novo*, DNA binding and transactivation and nuclear cytoplasm trafficking. To overcome the problems caused by the higher complexity of mammalian systems, experimental approaches with a more solid quantitative basis are needed. The concept of operating mechanism should be in place. When a new mechanism is discovered, it is imperative to know if it is operating *in vivo* and to assess its relative importance compared with alternative competing pathways. Otherwise the newly discovered mechanism will certainly add new information but its biological meaning is uncertain. Here we deduce a set of simple equations that help to extract kinetic quantitative data from typical experiments. Although, these equations are simplifications and may lack the rigor of a true kinetic approach, they, hopefully, can be easily applied to typical experimental data present in most signaling papers. We exemplified this with a couple of calculations arriving to a conclusion that in HeLa cells KEAP1 reactivity with H_2O_2 is weak, while a highly reactive sensor is probably mediating stimulation of NRF2 synthesis *de novo* by H_2O_2 .

Part of the complexity mentioned above may be artificial because there is a tendency to integrate information coming from different types of cells. One of the consequences of multicellularity is cell specialization and so different signaling mechanisms operate in different cells. For example, H_2O_2 activates SP1 translation only in neurons and not in glia cells. In spite of this cellular specialization, it is tempting to identify a set of players that recurrently appear as intermediates in H_2O_2 signaling pathways and that may function as hubs, namely ASK1/MAPK; GSK3 β , ATM, c-ABL, SIRT1, REF-1, as well as several phosphatases. Many of these molecules are kinases and phosphatases, underlying the well-known fact that H_2O_2 affects the phosphorylation state of signaling pathways. Recent advances with the discovery of the inhibition of Prxs by phosphorylation, show that this is a two-way relationship where signaling by phosphorylation/dephosphorylation affect the levels of H_2O_2 by decreasing its removal. If other thiol proteins have their SH groups reactivity with H_2O_2 also affected by its phosphorylation state, a cross-talk between protein thiol oxidation state and protein phosphorylation state would be established.

Our calculations shown in Table 1 help to understand how H_2O_2 signaling can be specific. Depending on the time of exposure and on H_2O_2 concentration different sensors may be activated. This chemical specificity combined with the localized production of H_2O_2 , where only a handful of molecular sensors are available, yields a high degree of biological specificity. We have proposed that H_2O_2 is not only a generic modulator of TFs and signaling molecules, but has the potential of being a specific regulator of individual genes [115], and the vast array of regulatory mechanisms by which H_2O_2 exerts its effects described here supports such view. To know which genes are activated by H_2O_2 in each type of tissue and how this is affected by the presence of single nucleotide polymorphisms in the promoter regions of genes is essential to have a complete view of the regulation of gene expression by H_2O_2 . Ultimately, such understanding will be essential for redox biology to be an active player in the emerging area of personalized medicine.

Acknowledgments

Supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal (PTDC/BIA-PRO/101624/2008 and PEst-OE/UI0612/2013). A fellowship was given to H.S. (FCT-SFRH/BSAB/1174/2011).

References

- [1] B Chance, The state of catalase in the Respiring bacterial cell, *Science* (New York, N.Y.), 116 (1952) 202–3. <http://dx.doi.org/10.1126/science.116.3008.202>, 17792314.
- [2] MP Czech, JC Lawrence, WS Lynn, Evidence for the involvement of sulfhydryl oxidation in the regulation of fat cell hexose transport by insulin, *Proceedings of the National Academy of Sciences of the United States of America*. 71 (1974) 4173–7. <http://dx.doi.org/10.1073/pnas.71.10.4173>, 4372610.
- [3] AA White, KM Crawford, CS Patt, PJ Lad, Activation of soluble guanylate cyclase from rat lung by incubation or by hydrogen peroxide, *Journal of Biological Chemistry*. 251 (1976) 7304–12.
- [4] TM Burke, MS Wolin, Hydrogen peroxide elicits pulmonary arterial relaxation and guanylate cyclase activation, *American Journal of Physiology*. 252 (1987) H721–H732.
- [5] D Heffetz, Y Zick, H_2O_2 potentiates phosphorylation of novel putative substrates for the insulin receptor kinase in intact Fao cells, *Journal of Biological Chemistry*. 264 (1989) 10126–32.
- [6] RH Burdon, C Rice-Evans, Free radicals and the regulation of mammalian cell proliferation, *Free Radical Research Communications*. 6 (1989) 345–58. <http://dx.doi.org/10.3109/10715768909087918>, 2676744.
- [7] MF Christman, G Storz, BN Ames, OxyR, a positive regulator of hydrogen peroxide-inducible genes in *Escherichia coli* and *Salmonella typhimurium*, is homologous to a family of bacterial regulatory proteins, *Proceedings of the National Academy of Sciences of the United States of America*. 86 (1989) 3484–8. <http://dx.doi.org/10.1073/pnas.86.10.3484>, 2471187.
- [8] FJ Staal, M Roederer, LA Herzenberg, LA Herzenberg, Intracellular thiols regulate activation of nuclear factor kappa B and transcription of human immunodeficiency virus, *Proceedings of the National Academy of Sciences of the United States of America*. 87 (1990) 9943–7. <http://dx.doi.org/10.1073/pnas.87.24.9943>, 2263644.
- [9] R Schreck, P Rieber, PA Baeuerle, Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1, *EMBO Journal*. 10 (1991) 2247–58.
- [10] B Meier, AR Cross, JT Hancock, FJ Kaup, OT Jones, Identification of a superoxide-generating NADPH oxidase system in human fibroblasts, *Biochemical Journal*. 275 (1991) 241–5.
- [11] TP Sztrowski, CF Nathan, Production of large amounts of hydrogen peroxide by human tumor cells, *Cancer Research*. 51 (1991) 794–8.
- [12] M Goodman, RM Bostick, O Kucuk, DP Jones, Clinical trials of antioxidants as cancer prevention agents: past, present, and future, *Free Radical Biology and Medicine*. 51 (2011) 1068–84. <http://dx.doi.org/10.1016/j.freeradbiomed.2011.05.018>, 21683786.
- [13] G Ferrer-Sueta, B Manta, H Botti, R Radi, M Trujillo, A Denicola, Factors affecting protein thiol reactivity and specificity in peroxide reduction, *Chemical Research in Toxicology*. 24 (2011) 434–50. <http://dx.doi.org/10.1021/tx100413v>, 21391663.
- [14] R Brigelius-Flohé, M Maiorino, Glutathione peroxidases, *Biochimica et Biophysica Acta (BBA) General Subjects*. 1830 (2013) 3289–303.
- [15] SG Rhee, HA Woo, IS Kil, SH Bae, Peroxiredoxin functions as a peroxidase and a regulator and sensor of local peroxides, *Journal of Biological Chemistry*. 287 (2012) 4403–10. <http://dx.doi.org/10.1074/jbc.R111.283432>, 22147704.
- [16] K Mahadev, A Zilbering, L Zhu, BJ Goldstein, Insulin-stimulated hydrogen peroxide reversibly inhibits protein-tyrosine phosphatase 1b *in vivo* and enhances the early insulin action cascade, *Journal of Biological Chemistry*. 276 (2001) 21938–42. <http://dx.doi.org/10.1074/jbc.C100109200>, 11297536.
- [17] A Salmeen, JN Andersen, MP Myers, T-C Meng, JA Hinks, NK Tonks, et al, Redox regulation of protein tyrosine phosphatase 1B involves a sulphenylamide intermediate, *Nature*. 423 (2003) 769–73. <http://dx.doi.org/10.1038/nature01680>, 12802338.
- [18] T-C Meng, T Fukada, NK Tonks, Reversible oxidation and inactivation of protein tyrosine phosphatases *in vivo*, *Molecular Cell*. 9 (2002) 387–99. [http://dx.doi.org/10.1016/S1097-2765\(02\)00445-8](http://dx.doi.org/10.1016/S1097-2765(02)00445-8), 11864611.
- [19] JM Denu, KG Tanner, Specific and reversible inactivation of protein tyrosine phosphatases by hydrogen peroxide: evidence for a sulfenic acid intermediate and implications for redox regulation, *Biochemistry (Moscow)*. 37 (1998) 5633–42.
- [20] CC Winterbourn, MB Hampton, Thiol chemistry and specificity in redox signaling, *Free Radical Biology and Medicine*. 45 (2008) 549–61. <http://dx.doi.org/10.1016/j.freeradbiomed.2008.05.004>, 18544350.
- [21] HJ Forman, M Maiorino, F Ursini, Signaling functions of reactive oxygen species, *Biochemistry (Moscow)*. 49 (2010) 835–42.
- [22] R Brigelius-Flohé, L Flohé, Basic principles and emerging concepts in the redox control of transcription factors, *Antioxidants and Redox Signalling*. 15 (2011) 2335–81. <http://dx.doi.org/10.1089/ars.2010.3534>, 21194351.
- [23] M Zheng, Activation of the OxyR transcription factor by reversible disulfide bond formation, *Science* (New York, N.Y.). 279 (1998) 1718–22.
- [24] S Iwakami, H Misu, T Takeda, M Sugimori, S Matsugo, S Kaneko, et al, Concentration-dependent dual effects of hydrogen peroxide on insulin signal transduction in H4IIEC hepatocytes, *PLoS One*. 6 (2011) e27401. <http://dx.doi.org/10.1371/journal.pone.0027401>, 22102892.
- [25] AC Matias, N Pedroso, N Teodoro, HS Marinho, F Antunes, JM Nogueira, et al, Down-regulation of fatty acid synthase increases the resistance of *Saccharomyces cerevisiae* cells to H_2O_2 , *Free Radical Biology and Medicine*. 43 (2007) 1458–65. <http://dx.doi.org/10.1016/j.freeradbiomed.2007.08.003>, 17936191.
- [26] AC Matias, HS Marinho, L Cyrne, E Herrero, F Antunes, Biphasic modulation of

- fatty acid synthase by hydrogen peroxide in *Saccharomyces cerevisiae*, Archives of Biochemistry and Biophysics. 515 (2011) 107–11. <http://dx.doi.org/10.1016/j.abb.2011.08.009>, 21907179.
- [27] R Kelley, T Ideker, Genome-wide fitness and expression profiling implicate Mga2 in adaptation to hydrogen peroxide, PLoS Genetics. 5 (2009) 1–13.
- [28] P Jaakkola, DR Mole, YM Tian, MJ Wilson, J Gielbert, SJ Gaskell, et al, Targeting of HIF- α to the von Hippel–Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation, Science (New York, N.Y.). 292 (2001) 468–72.
- [29] RM Jarvis, SM Hughes, EC Ledgerwood, Peroxiredoxin 1 functions as a signal peroxidase to receive, transduce, and transmit peroxide signals in mammalian cells, Free Radical Biology and Medicine. 53 (2012) 1522–30. <http://dx.doi.org/10.1016/j.freeradbiomed.2012.08.001>, 22902630.
- [30] WG Kaelin Jr, PJ Ratcliffe, Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway, Molecular Cell. 30 (2008) 393–402. <http://dx.doi.org/10.1016/j.molcel.2008.04.009>, 18498744.
- [31] YS Bae, SW Kang, MS Seo, IC Baines, E Tekle, PB Chock, et al, Epidermal growth factor (EGF)-induced generation of hydrogen peroxide. Role in EGF receptor-mediated tyrosine phosphorylation, Journal of Biological Chemistry. 272 (1997) 217–21. <http://dx.doi.org/10.1074/jbc.272.1.217>, 8995250.
- [32] CE Paulsen, TH Truong, FJ Garcia, A Homann, V Gupta, SE Leonard, et al, Peroxide-dependent sulfenylation of the EGFR catalytic site enhances kinase activity, Nature Chemical Biology. 8 (2012) 57–64.
- [33] SR Lee, KS Kwon, SR Kim, SG Rhee, Reversible inactivation of protein-tyrosine phosphatase 1B in A431 cells stimulated with epidermal growth factor, Journal of Biological Chemistry. 273 (1998) 15366–72. <http://dx.doi.org/10.1074/jbc.273.25.15366>, 9624118.
- [34] JN LaButti, G Chowdhury, TJ Reilly, KS Gates, Redox regulation of protein tyrosine phosphatase 1B (PTP1B) by peroxymonophosphate (=O3POOH), Journal of the American Chemical Society. 129 (2007) 5320–1. <http://dx.doi.org/10.1021/ja070194j>, 17411049.
- [35] DF Trindade, G Cerchiaro, O Augusto, A role for peroxymonocarbonate in the stimulation of biotiniol peroxidation by the bicarbonate/carbon dioxide pair, Chemical Research in Toxicology. 19 (2006) 1475–82. <http://dx.doi.org/10.1021/tx060146x>, 17112235.
- [36] H Zhou, H Singh, ZD Parsons, SM Lewis, S Bhattacharya, DR Seiner, et al, The biological buffer bicarbonate/CO₂ potentiates H₂O₂-mediated inactivation of protein tyrosine phosphatases, Journal of the American Chemical Society. 133 (2011) 15803–5. <http://dx.doi.org/10.1021/ja2077137>, 21913686.
- [37] F Antunes, E Cadenas, Cellular titration of apoptosis with steady-state concentrations of H₂O₂. Sub-micromolar levels of H₂O₂ induce apoptosis through Fenton chemistry independent of cellular thiol state, Free Radical Biology and Medicine. 30 (2001) 1008–18. [http://dx.doi.org/10.1016/S0891-5849\(01\)00493-2](http://dx.doi.org/10.1016/S0891-5849(01)00493-2), 11316581.
- [38] F Antunes, E Cadenas, Estimation of H₂O₂ gradients across biomembranes, FEBS Letters. 475 (2000) 121–6. [http://dx.doi.org/10.1016/S0014-5793\(00\)01638-0](http://dx.doi.org/10.1016/S0014-5793(00)01638-0), 10858501.
- [39] NM Mishina, PA Tyurin-Kuzmin, KN Markvicheva, AV Vorotnikov, VA Tkachuk, V Laketa, et al, Does cellular hydrogen peroxide diffuse or act locally? Antioxidants and Redox Signalling. 14 (2011) 1–7. <http://dx.doi.org/10.1089/ars.2010.3539>.
- [40] K Chen, MT Kirber, H Xiao, Y Yang, JF Keaney, Regulation of ROS signal transduction by NADPH oxidase 4 localization, Journal of Cell Biology. 181 (2008) 1129–39. <http://dx.doi.org/10.1083/jcb.200709049>, 18573911.
- [41] LC Seaver, JA Imlay, Hydrogen peroxide fluxes and compartmentalization inside growing *Escherichia coli*, Journal of Bacteriology. 183 (2001) 7182–9. <http://dx.doi.org/10.1128/JB.183.24.7182-7189.2001>, 11717277.
- [42] V Oliveira-Marques, T Silva, F Cunha, G Covas, HS Marinho, F Antunes, et al, A quantitative study of the cell-type specific modulation of c-Rel by hydrogen peroxide and TNF- α , Redox Biology. 1 (2013) 347–52. <http://dx.doi.org/10.1016/j.redox.2013.05.004>, 24024170.
- [43] HS Marinho, L Cyrne, E Cadenas, F Antunes, The cellular steady-state of H₂O₂: latency concepts and gradients, Methods in Enzymology. 527 (2013) 3–19. <http://dx.doi.org/10.1016/B978-0-12-405882-8.00001-5>, 23830623.
- [44] MR Branco, HS Marinho, Cyrne L, F Antunes, Decrease of H₂O₂ plasma membrane permeability during adaptation to H₂O₂ in *Saccharomyces cerevisiae*, Journal of Biological Chemistry. 279 (2004) 6501–6.
- [45] V Folmer, N Pedroso, AC Matias, SC Lopes, F Antunes, L Cyrne, et al, H₂O₂ induces rapid biophysical and permeability changes in the plasma membrane of *Saccharomyces cerevisiae*, Biochimica et Biophysica Acta Biomembranes. 1778 (2008) 1141–7.
- [46] N Pedroso, AC Matias, L Cyrne, F Antunes, C Borges, R Malhó, et al, Modulation of plasma membrane lipid profile and microdomains by H₂O₂ in *Saccharomyces cerevisiae*, Free Radical Biology and Medicine. 46 (2009) 289–98. <http://dx.doi.org/10.1016/j.freeradbiomed.2008.10.039>, 19027845.
- [47] M Gavella, V Garaj-Vrhovac, V Lipovac, M Antica, G Gajski, N Car, Ganglioside GT1b protects human spermatozoa from hydrogen peroxide-induced DNA and membrane damage, International Journal of Andrology. 33 (2010) 536–44.
- [48] GP Bienert, ALB Møller, KA Kristiansen, A Schulz, IM Møller, JK Schjoerring, et al, Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes, Journal of Biological Chemistry. 282 (2007) 1183–92. <http://dx.doi.org/10.1074/jbc.M603761200>, 17105724.
- [49] EW Miller, BC Dickinson, CJ Chang, Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling, Proceedings of the National Academy of Sciences of the United States of America. 107 (2010) 15681–6. <http://dx.doi.org/10.1073/pnas.1005776107>, 20724658.
- [50] HA Woo, SH Yim, DH Shin, D Kang, D-Y Yu, SG Rhee, Inactivation of peroxiredoxin I by phosphorylation allows localized H₂O₂ accumulation for cell signaling, Cell. 140 (2010) 517–28. <http://dx.doi.org/10.1016/j.cell.2010.01.009>, 20178744.
- [51] TH Truong, KS Carroll, Redox regulation of Epidermal Growth Factor Receptor signaling through cysteine oxidation, Biochemistry (Moscow). 51 (2012) 9954–65. <http://dx.doi.org/10.1021/bi301441e>.
- [52] MB Toledano, A-G Planson, A Delaunay-Moisan, Reining in H₂O₂ for safe signaling, Cell. 140 (2010) 454–6. <http://dx.doi.org/10.1016/j.cell.2010.02.003>, 20178737.
- [53] SJ Rawat, CL Creasy, JR Peterson, J Chernoff, The tumor suppressor Mst1 promotes changes in the cellular redox state by phosphorylation and inactivation of peroxiredoxin-1 protein, Journal of Biological Chemistry. 288 (2013) 8762–71. <http://dx.doi.org/10.1074/jbc.M112.414524>, 23386615.
- [54] N Watanabe, JW Zmijewski, W Takabe, M Umezue-Goto, Le Goffe C, A Sekine, et al, Activation of mitogen-activated protein kinases by lysophosphatidylcholine-induced mitochondrial reactive oxygen species generation in endothelial cells, American Journal of Pathology. 168 (2006) 1737–48. <http://dx.doi.org/10.2353/ajpath.2006.050648>, 16651638.
- [55] VE Kagan, VA Tyurin, J Jiang, YY Tyurina, VB Ritov, AA Amoscato, et al, Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors, Nature Chemical Biology. 1 (2005) 223–32. <http://dx.doi.org/10.1038/nchembio727>, 16408039.
- [56] A Higdon, AR Diers, JY Oh, A Landar, VM Darley-Usmar, Cell signalling by reactive lipid species: new concepts and molecular mechanisms, Biochemical Journal. 442 (2012) 453–64. <http://dx.doi.org/10.1042/BJ20111752>, 22364280.
- [57] JA Imlay, Cellular defenses against superoxide and hydrogen peroxide, Annual Review of Biochemistry. 77 (2008) 755–76. <http://dx.doi.org/10.1146/annurev.biochem.77.061606.161055>, 18173371.
- [58] K Tao, K Makino, S Yonei, A Nakata, H Shinagawa, Molecular cloning and nucleotide sequencing of oxyR, the positive regulatory gene of a regulon for an adaptive response to oxidative stress in *Escherichia coli*: homologues between OxyR protein and a family of bacterial activator proteins, Molecular and General Genetics. 218 (1989) 371–6.
- [59] MA Schell, Molecular biology of the LysR family of transcriptional regulators, Annual Review of Microbiology. 47 (1993) 597–626. <http://dx.doi.org/10.1146/annurev.mi.47.100193.003121>, 8257110.
- [60] MB Toledano, I Kullik, F Trinh, PT Baird, TD Schneider, G Storz, Redox-dependent shift of OxyR-DNA contacts along an extended DNA-binding site: a mechanism for differential promoter selection, Cell. 78 (1994) 897–909. [http://dx.doi.org/10.1016/S0092-8674\(94\)90702-1](http://dx.doi.org/10.1016/S0092-8674(94)90702-1), 8087856.
- [61] G Storz, LA Tartaglia, BN Ames, Transcriptional regulator of oxidative stress-inducible genes: direct activation by oxidation, Science (New York, N.Y.). 248 (1990) 189–94.
- [62] N Bsat, A Herbig, L Casillas-Martinez, P Setlow, JD Helmann, *Bacillus subtilis* contains multiple Fur homologues: identification of the iron uptake (Fur) and peroxide regulon (PerR) repressors, Molecular Microbiology. 29 (1998) 189–98. <http://dx.doi.org/10.1046/j.1365-2958.1998.00921.x>, 9701813.
- [63] AF Herbig, JD Helmann, Roles of metal ions and hydrogen peroxide in modulating the interaction of the *Bacillus subtilis* PerR peroxide regulon repressor with operator DNA, Molecular Microbiology. 41 (2001) 849–59.
- [64] JD Helmann, MF Wu, A Gaballa, PA Kobel, MM Morshedi, P Fawcett, et al, The global transcriptional response of *Bacillus subtilis* to peroxide stress is coordinated by three transcription factors, Journal of Bacteriology. 185 (2003) 243–53. <http://dx.doi.org/10.1128/JB.185.1.243-253.2003>, 12486061.
- [65] JW Lee, JD Helmann, Functional specialization within the Fur family of metal-oregulators, Biometals : An International Journal on the Role of Metal Ions in Biology, Biochemistry, and Medicine. 20 (2007) 485–99.
- [66] J-W Lee, JD Helmann, The PerR transcription factor senses H₂O₂ by metal-catalysed histidine oxidation, Nature. 440 (2006) 363–7. <http://dx.doi.org/10.1038/nature04537>, 16541078.
- [67] JM Dubbs, S Mongkolsuk, Peroxide-sensing transcriptional regulators in bacteria, Journal of Bacteriology. 194 (2012) 5495–503. <http://dx.doi.org/10.1128/JB.00304-12>, 22797754.
- [68] L Chen, L Keramati, JD Helmann, Coordinate regulation of *Bacillus subtilis* peroxide stress genes by hydrogen peroxide and metal ions, Proceedings of the National Academy of Sciences of the United States of America. 92 (1995) 8190–4. <http://dx.doi.org/10.1073/pnas.92.18.8190>, 7667267.
- [69] M Fungthong, AF Herbig, N Bsat, JD Helmann, Regulation of the *Bacillus subtilis* fur and perR genes by PerR: not all members of the PerR regulon are peroxide inducible, Journal of Bacteriology. 184 (2002) 3276–86. <http://dx.doi.org/10.1128/JB.184.12.3276-3286.2002>, 12029044.
- [70] A Gaballa, JD Helmann, A peroxide-induced zinc uptake system plays an important role in protection against oxidative stress in *Bacillus subtilis*, Molecular Microbiology. 45 (2002) 997–1005. <http://dx.doi.org/10.1046/j.1365-2958.2002.03068.x>, 12180919.
- [71] M Rinnerthaler, S Büttner, P Laun, G Heeren, TK Felder, H Klinger, et al, Yno1p/Aim14p, a NADPH-oxidase ortholog, controls extramitochondrial reactive oxygen species generation, apoptosis, and actin cable formation in yeast, Proceedings of the National Academy of Sciences of the United States of America. 109 (2012) 8658–63. <http://dx.doi.org/10.1073/pnas.1201629109>, 22586098.
- [72] K Block, Y Gorin, HE Abboud, Subcellular localization of Nox4 and regulation in diabetes, Proceedings of the National Academy of Sciences of the United States of America. 106 (2009) 14385–90. <http://dx.doi.org/10.1073/pnas.0906805106>, 19706525.
- [73] L Fernandes, C Rodrigues-Pousada, Struhl K, Yap, a novel family of eight bZIP

- proteins in *Saccharomyces cerevisiae* with distinct biological functions, *Molecular and Cellular Biology*. 17 (1997) 6982–93. .
- [74] WS Moye-Rowley, KD Harshman, CS Parker, Yeast YAP1 encodes a novel form of the jun family of transcriptional activator proteins, *Genes and Development*. 3 (1989) 283–92. <http://dx.doi.org/10.1101/gad.3.3.283>, 2542125.
 - [75] S Kuge, Regulation of yAP-1 nuclear localization in response to oxidative stress, *EMBO Journal*. 16 (1997) 1710–20. .
 - [76] A Delaunay, A-D Isnard, MB Toledano, H₂O₂ sensing through oxidation of the Yap1 transcription factor, *EMBO Journal*. 19 (2000) 5157–66. .
 - [77] K Pluta, O Lefebvre, NC Martin, WJ Smagowicz, DR Stanford, SR Ellis, et al, Maf1p, a negative effector of RNA polymerase III in *Saccharomyces cerevisiae*, *Molecular and Cellular Biology*. 21 (2001) 5031–40. <http://dx.doi.org/10.1128/MCB.21.15.5031-5040.2001>, 11438659.
 - [78] M Boguta, Maf1, a general negative regulator of RNA polymerase III in yeast, *Biochimica et Biophysica Acta*. 1829 (2013) 376–84. .
 - [79] D Graczyk, J Debski, G Muszyńska, M Bretner, O Lefebvre, M Boguta, Casein kinase II-mediated phosphorylation of general repressor Maf1 triggers RNA polymerase III activation, *Proceedings of the National Academy of Sciences of the United States of America*. 108 (2011) 4926–31. <http://dx.doi.org/10.1073/pnas.1010010108>, 21383183.
 - [80] A Vannini, R Ringel, AG Kusser, O Berninghausen, GA Kassavetis, P Cramer, Molecular basis of RNA polymerase III transcription repression by Maf1, *Cell*. 143 (2010) 59–70. <http://dx.doi.org/10.1016/j.cell.2010.09.002>, 20887893.
 - [81] M Cieślą, M Boguta, Regulation of RNA polymerase III transcription by Maf1 protein, *Acta Biochimica Polonica*. 55 (2008) 215–25. .
 - [82] M Akerfelt, RI Morimoto, L Sistonen, Heat shock factors: integrators of cell stress, development and lifespan, *Nature Reviews. Molecular Cell Biology*. 11 (2010) 545–55. .
 - [83] PK Sorger, MJ Lewis, HR Pelham, Heat shock factor is regulated differently in yeast and HeLa cells, *Nature*. 329 (1987) 81–4. <http://dx.doi.org/10.1038/329081a0>, 3306402.
 - [84] BK Jakobsen, HR Pelham, Constitutive binding of yeast heat shock factor to DNA in vivo, *Molecular and Cellular Biology*. 8 (1988) 5040–2. .
 - [85] A Yamamoto, J Ueda, N Yamamoto, N Hashikawa, H Sakurai, Role of heat shock transcription factor in *Saccharomyces cerevisiae* oxidative stress response, *Eukaryotic Cell*. 6 (2007) 1373–9. <http://dx.doi.org/10.1128/EC.00098-07>, 17586717.
 - [86] A Yamamoto, Y Mizukami, H Sakurai, Identification of a novel class of target genes and a novel type of binding sequence of heat shock transcription factor in *Saccharomyces cerevisiae*, *Journal of Biological Chemistry*. 280 (2005) 11911–19. <http://dx.doi.org/10.1074/jbc.M411256200>, 15647283.
 - [87] MT Martínez-Pastor, G Marchler, C Schüller, A Marchler-Bauer, H Ruis, F Estruch, The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE), *EMBO Journal*. 15 (1996) 2227–35. .
 - [88] A Sebé-Pedrós, A Mendoza, BF Lang, BM Degnan, I Ruiz-Trillo, Unexpected repertoire of metazoan transcription factors in the unicellular holozoan *Cap-saspora owczarzaki*, *Molecular Biology and Evolution*. 28 (2010) 1241–54.
 - [89] M Karin, E Shaulian, AP-1: linking hydrogen peroxide and oxidative stress to the control of cell proliferation and death, *IUBMB Life*. 52 (2001) 17–24. .
 - [90] H Motohashi, O'Connor T, F Katsuoka, JD Engel, M Yamamoto, Integration and diversity of the regulatory network composed of Maf and CNC families of transcription factors, *Gene*. 294 (2002) 1–12. [http://dx.doi.org/10.1016/S0378-1119\(02\)00788-6](http://dx.doi.org/10.1016/S0378-1119(02)00788-6), 12234662.
 - [91] Kappelmann, M, Bosserhoff, A, Kuphal, AP-1/c-Jun transcription factors: regulation and function in malignant melanoma, *European Journal of Cell Biology*. (2013).
 - [92] M Kobayashi, L Li, N Iwamoto, Y Nakajima-Takagi, H Kaneko, Y Nakayama, et al, The antioxidant defense system Keap1-Nrf2 comprises a multiple sensing mechanism for responding to a wide range of chemical compounds, *Molecular and Cellular Biology*. 29 (2009) 493–502. <http://dx.doi.org/10.1128/MCB.01080-08>, 19001094.
 - [93] R Venugopal, AK Jaiswal, Nrf2 and Nrf1 in association with Jun proteins regulate antioxidant response element-mediated expression and coordinated induction of genes encoding detoxifying enzymes, *Oncogene*. 17 (1998) 3145–56. <http://dx.doi.org/10.1038/sj.onc.1202237>, 9872330.
 - [94] AJ Shaywitz, ME Greenberg, CREB: A stimulus-induced transcription factor activated by a diverse array of extracellular signals, *Annual Review of Biochemistry*. 68 (1999) 821–61. <http://dx.doi.org/10.1146/annurev.biochem.68.1.821>, 10872467.
 - [95] M Johannessen, MP Delghandi, U Moens, What turns CREB on? *Cellular Signalling*. 16 (2004) 1211–27. <http://dx.doi.org/10.1016/j.cellsig.2004.05.001>, 15337521.
 - [96] De Cesare D, GM Fimia, P Sassone-Corsi, Signaling routes to CREM and CREB: plasticity in transcriptional activation, *Trends in Biochemical Sciences*. 24 (1999) 281–5. [http://dx.doi.org/10.1016/S0968-0004\(99\)01414-0](http://dx.doi.org/10.1016/S0968-0004(99)01414-0), 10390618.
 - [97] M Johannessen, MP Delghandi, A Rykx, M Dragset, JR Vandenheede, J Van Lint, et al, Protein kinase D induces transcription through direct phosphorylation of the cAMP-response Element-binding Protein, *Journal of Biological Chemistry*. 282 (2007) 14777–87. <http://dx.doi.org/10.1074/jbc.M610669200>, 17389598.
 - [98] AJ Levine, M Oren, The first 30 years of p53: growing ever more complex, *Nature Reviews. Cancer*. 9 (2009) 749–58. <http://dx.doi.org/10.1038/nrc2723>, 19776744.
 - [99] KH Vousden, X Lu, Live or let die: the cell's response to p53, *Nature Reviews. Cancer*. 2 (2002) 594–604. <http://dx.doi.org/10.1038/nrc864>, 12154352.
 - [100] DP Lane, Cancer. p53, guardian of the genome, *Nature*. 358 (1992) 15–16. <http://dx.doi.org/10.1038/358015a0>, 1614522.
 - [101] M Oren, J Bartek, The sunny side of p53, *Cell*. 128 (2007) 826–8. <http://dx.doi.org/10.1016/j.cell.2007.02.027>, 17350568.
 - [102] M Hollstein, P Hainaut, Massively regulated genes: the example of TP53, *Journal of Pathology*. 220 (2010) 164–73. .
 - [103] D Sakamuro, P Sabbatini, E White, GC Prendergast, The polyproline region of p53 is required to activate apoptosis but not growth arrest, *Oncogene*. 15 (1997) 887–98. <http://dx.doi.org/10.1038/sj.onc.1201263>, 9285684.
 - [104] J Bargonetti, JJ Manfredi, X Chen, DR Marshak, C Prives, A proteolytic fragment from the central region of p53 has marked sequence-specific DNA-binding activity when generated from wild-type but not from oncogenic mutant p53 protein, *Genes and Development*. 7 (1993) 2565–74. <http://dx.doi.org/10.1101/gad.7.12b.2565>, 8276239.
 - [105] NP Pavletich, KA Chambers, CO Pabo, The DNA-binding domain of p53 contains the four conserved regions and the major mutation hot spots, *Genes and Development*. 7 (1993) 2556–64. <http://dx.doi.org/10.1101/gad.7.12b.2556>, 8276238.
 - [106] SE Kern, KW Kinzler, A Bruskin, D Jarosz, P Friedman, C Prives, et al, Identification of p53 as a sequence-specific DNA-binding protein, *Science (New York, N.Y.)*. 252 (1991) 1708–11. .
 - [107] WS El-Deiry, SE Kern, JA Pietenpol, KW Kinzler, B Vogelstein, Definition of a consensus binding site for p53, *Nature Genetics*. 1 (1992) 45–9. <http://dx.doi.org/10.1038/ng0492-45>, 1301998.
 - [108] JM Stommel, A leucine-rich nuclear export signal in the p53 tetramerization domain: regulation of subcellular localization and p53 activity by NES masking, *EMBO Journal*. 18 (1999) 1660–72.
 - [109] M Reed, B Woelker, P Wang, Y Wang, ME Anderson, P Tegtmeier, The C-terminal domain of p53 recognizes DNA damaged by ionizing radiation, *Proceedings of the National Academy of Sciences*. 92 (1995) 9455–9. <http://dx.doi.org/10.1073/pnas.92.21.9455>.
 - [110] R Kopan, MXG Ilagan, The canonical Notch signaling pathway: unfolding the activation mechanism, *Cell*. 137 (2009) 216–33. <http://dx.doi.org/10.1016/j.cell.2009.03.045>, 19379690.
 - [111] G Struhl, I Greenwald, Presenilin is required for activity and nuclear access of Notch in *Drosophila*, *Nature*. 398 (1999) 522–5. <http://dx.doi.org/10.1038/19091>, 10206646.
 - [112] ME Fortini, Introduction—Notch in development and disease, *Seminars in Cell and Developmental Biology*. 23 (2012) 419–20. <http://dx.doi.org/10.1016/j.semcdb.2012.03.001>, 22425963.
 - [113] S Ghosh, MJ May, EB Kopp, NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses, *Annual Review of Immunology*. 16 (1998) 225–60. .
 - [114] NX Li, M Karin, Is NF-kappaB the sensor of oxidative stress? *FASEB Journal*. 13 (1999) 1137–43.
 - [115] V Oliveira-Marques, HS Marinho, L Cyrne, F Antunes, Role of hydrogen peroxide in NF-kappaB activation: from inducer to modulator, *Antioxidants and Redox Signalling*. 11 (2009) 2223–43. <http://dx.doi.org/10.1089/ars.2009.2601>, 19496701.
 - [116] De Oliveira-Marques V, L Cyrne, H Marinho, F Antunes, A quantitative study of NF-kappaB activation by H₂O₂: relevance in inflammation and synergy with TNF-alpha, *Journal of Immunology*. 178 (2007) 3893–902.
 - [117] WS Dynan, R Tjian, The promoter-specific transcription factor Sp1 binds to upstream sequences in the SV40 early promoter, *Cell*. 35 (1983) 79–87. [http://dx.doi.org/10.1016/0092-8674\(83\)90210-6](http://dx.doi.org/10.1016/0092-8674(83)90210-6), 6313230.
 - [118] MR Briggs, JT Kadonaga, SP Bell, R Tjian, Purification and biochemical characterization of the promoter-specific transcription factor, Sp1, *Science (New York, N.Y.)*. 234 (1986) 47–52. .
 - [119] JT Kadonaga, AJ Courey, J Ladika, R Tjian, Distinct regions of Sp1 modulate DNA binding and transcriptional activation, *Science (New York, N.Y.)*. 242 (1988) 1566–70. .
 - [120] H Ryu, J Lee, K Zaman, J Kubilis, RJ Ferrante, BD Ross, et al, Sp1 and Sp3 are oxidative stress-inducible, antideath transcription factors in cortical neurons, *Journal of Neuroscience: The Official Journal of the Society for Neuroscience*. 23 (2003) 3597–606. .
 - [121] SH Yeh, WB Yang, PW Gean, CY Hsu, JT Tseng, TP Su, et al, Translational and transcriptional control of Sp1 against ischaemia through a hydrogen peroxide-activated internal ribosomal entry site pathway, *Nucleic Acids Research*. 39 (2011) 5412–23. <http://dx.doi.org/10.1093/nar/gkr161>, 21441538.
 - [122] FS Santiago, H Ishii, S Shafi, R Khurana, P Kanellakis, R Bhindi, et al, Yin Yang-1 inhibits vascular smooth muscle cell growth and intimal thickening by repressing p21WAF1/Cip1 transcription and p21WAF1/Cip1-Cdk4-cyclin D1 assembly, *Circulation Research*. 101 (2007) 146–55. <http://dx.doi.org/10.1161/CIRCRESAHA.106.145235>, 17556661.
 - [123] AW Dunah, Sp1 and TAFII130 transcriptional activity disrupted in early Huntington's disease, *Science*. 296 (2002) 2238–43.
 - [124] Y-W Liu, H-P Tseng, L-C Chen, B-K Chen, W-C Chang, Functional cooperation of simian virus 40 promoter factor 1 and CCAAT/enhancer-binding protein beta and delta in lipopolysaccharide-induced gene activation of IL-10 in mouse macrophages, *Journal of Immunology*. 171 (2003) 821–8. .
 - [125] V Fridmacher, B Kaltschmidt, B Goudeau, D Ndiaye, FM Rossi, J Pfeiffer, et al, Forebrain-specific neuronal inhibition of nuclear factor-kappaB activity leads to loss of neuroprotection, *Journal of Neuroscience*. 23 (2003) 9403–8. .
 - [126] M Meissner, PPARalpha activators inhibit vascular endothelial growth factor receptor-2 expression by repressing Sp1-dependent DNA binding and trans-activation, *Circulation Research*. 94 (2004) 324–32. <http://dx.doi.org/10.1161/01.RES.0000113781.08139.81>, 14684628.

- [127] S Chu, TJ Ferro, Sp1: regulation of gene expression by phosphorylation, *Gene*. 348 (2005) 1–11. <http://dx.doi.org/10.1016/j.gene.2005.01.013>, 15777659.
- [128] NY Tan, LM Khachigian, Sp1 phosphorylation and its regulation of gene transcription, *Molecular and Cellular Biology*. 29 (2009) 2483–8. <http://dx.doi.org/10.1128/MCB.01828-08>, 19273606.
- [129] JA Forsythe, BH Jiang, NV Iyer, F Agani, SW Leung, RD Koos, et al, Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1, *Molecular and Cellular Biology*. 16 (1996) 4604–13.
- [130] GL Semenza, GL Wang, A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation, *Molecular and Cellular Biology*. 12 (1992) 5447–54.
- [131] TN Seagroves, HE Ryan, H Lu, BG Wouters, M Knapp, P Thibault, et al, Transcription factor HIF-1 is a necessary mediator of the pasteur effect in mammalian cells, *Molecular and Cellular Biology*. 21 (2001) 3436–44. <http://dx.doi.org/10.1128/MCB.21.10.3436-3444.2001>, 11313469.
- [132] A Weidemann, RS Johnson, Biology of HIF-1 α , *Cell Death and Differentiation*. 15 (2008) 621–7. <http://dx.doi.org/10.1038/cdd.2008.12>, 18259201.
- [133] M Ivan, K Kondo, H Yang, W Kim, J Valiando, M Ohh, et al, HIF α targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing, *Science*. 292 (2001) 464–8.
- [134] JK Brunelle, EL Bell, NM Quesada, K Vercauteren, V Tiranti, M Zeviani, et al, Oxygen sensing requires mitochondrial ROS but not oxidative phosphorylation, *Cell Metabolism*. 1 (2005) 409–14. <http://dx.doi.org/10.1016/j.cmet.2005.05.002>, 16054090.
- [135] S Kaewpila, S Venkataraman, GR Buettner, LW Oberley, Manganese superoxide dismutase modulates hypoxia-inducible factor-1 α induction via superoxide, *Cancer Research*. 68 (2008) 2781–8. <http://dx.doi.org/10.1158/0008-5472.CAN-07-2635>, 18413745.
- [136] H Shimano, Sterol regulatory element-binding proteins (SREBPs): transcriptional regulators of lipid synthetic genes, *Progress in Lipid Research*. 40 (2001) 439–52. [http://dx.doi.org/10.1016/S0163-7827\(01\)00010-8](http://dx.doi.org/10.1016/S0163-7827(01)00010-8), 11591434.
- [137] MS Brown, JL Goldstein, A proteolytic pathway that controls the cholesterol content of membranes, cells, and blood, *Proceedings of the National Academy of Sciences*. 96 (1999) 11041–8. <http://dx.doi.org/10.1073/pnas.96.20.11041>.
- [138] JD Horton, JL Goldstein, MS Brown, SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver, *Journal of Clinical Investigation*. 109 (2002) 1125–31. <http://dx.doi.org/10.1172/JCI200215593>, 11994399.
- [139] M Sekiya, A Hiraishi, M Touyama, K Sakamoto, Oxidative stress induced lipid accumulation via SREBP1c activation in HepG2 cells, *Biochemical and Biophysical Research Communications*. 375 (2008) 602–7. <http://dx.doi.org/10.1016/j.bbrc.2008.08.068>, 18727921.
- [140] A Nothmann, Topology of SREBP cleavage-activating protein, a polytopic membrane protein with a sterol-sensing domain, *Journal of Biological Chemistry*. 273 (1998) 17243–50. <http://dx.doi.org/10.1074/jbc.273.27.17243>, 9642295.
- [141] X Hua, C Yokoyama, J Wu, MR Briggs, MS Brown, JL Goldstein, et al, SREBP-2, a second basic-helix-loop-helix-leucine zipper protein that stimulates transcription by binding to the sterol regulatory element, *Proceedings of the National Academy of Sciences*. 90 (1993) 11603–7. <http://dx.doi.org/10.1073/pnas.90.24.11603>.
- [142] PJ Espenshade, SREBPs: sterol-regulated transcription factors, *Journal of Cell Science*. 119 (2006) 973–6. <http://dx.doi.org/10.1242/jcs.02866>, 16525117.
- [143] T Yang, PJ Espenshade, ME Wright, D Yabe, Y Gong, R Aebersold, et al, Crucial in cholesterol homeostasis: sterols promote binding of SCAP to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER, *Cell*. 110 (2002) 489–500. [http://dx.doi.org/10.1016/S0092-8674\(02\)00872-3](http://dx.doi.org/10.1016/S0092-8674(02)00872-3), 12202038.
- [144] M Matsuda, SREBP cleavage-activating protein (SCAP) is required for increased lipid synthesis in liver induced by cholesterol deprivation and insulin elevation, *Genes and Development*. 15 (2001) 1206–16. <http://dx.doi.org/10.1101/gad.891301>, 11358865.
- [145] RB Rawson, Failure to cleave sterol regulatory element-binding proteins (SREBPs) causes cholesterol auxotrophy in Chinese hamster ovary cells with genetic absence of SREBP cleavage-activating protein, *Journal of Biological Chemistry*. 274 (1999) 28549–56. <http://dx.doi.org/10.1074/jbc.274.40.28549>, 10497220.
- [146] MS Brown, JL Goldstein, The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor, *Cell*. 89 (1997) 331–40. [http://dx.doi.org/10.1016/S0092-8674\(00\)80213-5](http://dx.doi.org/10.1016/S0092-8674(00)80213-5), 9150132.
- [147] J Anckar, L Sistonen, Regulation of HSF1 function in the heat stress response: Implications in aging and disease, *Annual Review of Biochemistry*. 80 (2011) 1089–115. <http://dx.doi.org/10.1146/annurev-biochem-060809-095203>, 21417720.
- [148] Y Shi, DD Mosser, RI Morimoto, Molecular chaperones as HSF1-specific transcriptional repressors, *Genes and Development*. 12 (1998) 654–66. <http://dx.doi.org/10.1101/gad.12.5.654>, 9499401.
- [149] K Abravaya, MP Myers, SP Murphy, RI Morimoto, The human heat shock protein hsp70 interacts with HSF, the transcription factor that regulates heat shock gene expression, *Genes and Development*. 6 (1992) 1153–64. <http://dx.doi.org/10.1101/gad.6.7.1153>, 1628823.
- [150] R Baler, WJ Welch, R Voellmy, Heat shock gene regulation by nascent polypeptides and denatured proteins: hsp70 as a potential autoregulatory factor, *Journal of Cell Biology*. 117 (1992) 1151–9. <http://dx.doi.org/10.1083/jcb.117.6.1151>, 1607379.
- [151] DS Latchman, Transcription factors: an overview, *International Journal of Biochemistry and Cell Biology*. 29 (1997) 1305–12. [http://dx.doi.org/10.1016/S1357-2725\(97\)00085-X](http://dx.doi.org/10.1016/S1357-2725(97)00085-X), 9570129.
- [152] I Schreck, M Al-Rawi, J-M Mingot, C Scholl, ME Diefenbacher, O'Donnell P, et al, c-Jun localizes to the nucleus independent of its phosphorylation by and interaction with JNK and vice versa promotes nuclear accumulation of JNK, *Biochemical and Biophysical Research Communications*. 407 (2011) 735–40. <http://dx.doi.org/10.1016/j.bbrc.2011.03.092>, 21439937.
- [153] K Chida, S Nagamori, T Kuroki, Nuclear translocation of Fos is stimulated by interaction with Jun through the leucine zipper, *Cellular and Molecular Life Sciences*. 55 (1999) 297–302. <http://dx.doi.org/10.1007/s00180050291>, 10188588.
- [154] A Matsuzawa, H Ichijo, Redox control of cell fate by MAP kinase: physiological roles of ASK1–MAP kinase pathway in stress signaling, *Biochimica et Biophysica Acta*. 1780 (2008) 1325–36.
- [155] P Angel, K Hattori, T Smeal, M Karin, *Cell*. 55 (1988), 875–885.
- [156] AJ Whitmarsh, RJ Davis, Transcription factor AP-1 regulation by mitogen-activated protein kinase signal transduction pathways, *Journal of Molecular Medicine* (Berlin, Germany). 74 (1996) 589–607. <http://dx.doi.org/10.1007/s001090050063>, 8912180.
- [157] D Uberti, E Yavin, S Gil, K-R Ayasola, N Goldfinger, V Rotter, Hydrogen peroxide induces nuclear translocation of p53 and apoptosis in cells of oligodendroglia origin, *Molecular Brain Research*. 65 (1999) 167–75. [http://dx.doi.org/10.1016/S0169-328X\(98\)00339-8](http://dx.doi.org/10.1016/S0169-328X(98)00339-8), 10064887.
- [158] F Gugliesi, Up-regulation of the interferon-inducible IFI16 gene by oxidative stress triggers p53 transcriptional activity in endothelial cells, *Journal of Leukocyte Biology*. 77 (2005) 820–9. <http://dx.doi.org/10.1189/jlb.0904507>, 15728246.
- [159] De Sarno P, Muscarinic Receptor activation protects cells from apoptotic effects of DNA damage, oxidative stress, and mitochondrial Inhibition, *Journal of Biological Chemistry*. 278 (2003) 11086–93. <http://dx.doi.org/10.1074/jbc.M212157200>, 12538580.
- [160] C McNeill-Blue, BA Wetmore, JF Sanchez, WJ Freed, BA Merrick, Apoptosis mediated by p53 in rat neural AF5 cells following treatment with hydrogen peroxide and staurosporine, *Brain Research*. 1112 (2006) 1–15. <http://dx.doi.org/10.1016/j.brainres.2006.07.024>, 16901471.
- [161] R Saldaña-Meyer, F Recillas-Targa, Transcriptional and epigenetic regulation of the p53 tumor suppressor gene, *Epigenetics*. 6 (2011) 1068–77. <http://dx.doi.org/10.4161/epi.6.9.16683>, 21814038.
- [162] H-C Kirch, S Flaswinkel, H Rumpf, D Brockmann, H Esche, Expression of human p53 requires synergistic activation of transcription from the p53 promoter by AP-1, NF- κ B and Myc/Max, *Oncogene*. 18 (1999) 2728–38. <http://dx.doi.org/10.1038/sj.onc.1202626>, 10348347.
- [163] H Wu, G Lozano, NF- κ B activation of p53. A potential mechanism for suppressing cell growth in response to stress, *Journal of Biological Chemistry*. 269 (1994) 20667–74.
- [164] V Oliveira-Marques, HS Marinho, L Cyrne, F Antunes, Modulation of NF- κ B-dependent gene expression by H₂O₂: a major role for a simple chemical process in a complex biological response, *Antioxidants and Redox Signaling*. 11 (2009) 2043–53. <http://dx.doi.org/10.1089/ars.2008.2279>.
- [165] EL Pagé, GA Robitaille, J Pouyssegur, DE Richard, Induction of hypoxia-inducible factor-1 α by transcriptional and translational mechanisms, *Journal of Biological Chemistry*. 277 (2002) 48403–9. <http://dx.doi.org/10.1074/jbc.M209114200>, 12379645.
- [166] S Bonello, C Zähringer, RS BelAiba, T Djordjevic, J Hess, C Michiels, et al, Reactive oxygen species activate the HIF-1 α promoter via a functional NF- κ B site, *Arteriosclerosis, Thrombosis and Vascular Biology*. 27 (2007) 755–61. <http://dx.doi.org/10.1161/01.ATV.0000258979.92828.bc>.
- [167] K Abdelmohsen, R Pullmann, A Lal, HH Kim, S Galban, X Yang, et al, Phosphorylation of HuR by Chk2 regulates SIRT1 expression, *Molecular Cell*. 25 (2007) 543–57. <http://dx.doi.org/10.1016/j.molcel.2007.01.011>, 17317627.
- [168] H Mochizuki, CJ Murphy, JD Brandt, Y Kiuchi, P Russell, Altered stability of mRNAs associated with glaucoma progression in human trabecular meshwork cells following oxidative stress, *Investigative Ophthalmology and Visual Science*. 53 (2012) 1734–41. <http://dx.doi.org/10.1167/jovs.12-7938>, 22395891.
- [169] M Holcik, N Sonenberg, Translational control in stress and apoptosis, *Nature Reviews Molecular Cell Biology*. 6 (2005) 318–27. <http://dx.doi.org/10.1038/nrm1618>, 15803138.
- [170] Q Zhou, L-Z Liu, B Fu, X Hu, X Shi, J Fang, et al, Reactive oxygen species regulate insulin-induced VEGF and HIF-1 α expression through the activation of p70S6K1 in human prostate cancer cells, *Carcinogenesis*. 28 (2007) 28–37. <http://dx.doi.org/10.1093/carcin/bgl085>, 16774940.
- [171] SE Purdom-Dickinson, EV Sheveleva, H Sun, QM Chen, Translational control of Nrf2 protein in activation of antioxidant response by oxidants, *Molecular Pharmacology*. 72 (2007) 1074–81. <http://dx.doi.org/10.1124/mol.107.035360>, 17652445.
- [172] G Covas, HS Marinho, L Cyrne, F Antunes, Activation of Nrf2 by H₂O₂: de novo synthesis versus nuclear translocation, *Methods in Enzymology*. 528 (2013) 157–71. <http://dx.doi.org/10.1016/B978-0-12-405881-1.00009-4>, 23849864.
- [173] W Li, N Thakor, EY Xu, Y Huang, C Chen, R Yu, et al, An internal ribosomal entry site mediates redox-sensitive translation of Nrf2, *Nucleic Acids Research*. 38 (2010) 778–88. <http://dx.doi.org/10.1093/nar/gkp1048>, 19934254.
- [174] M Stoneley, AE Willis, Cellular internal ribosome entry segments: structures, trans-acting factors and regulation of gene expression, *Oncogene*. 23 (2004) 3200–7. <http://dx.doi.org/10.1038/sj.onc.1207551>, 15094769.
- [175] J Zhang, TN Dinh, K Kappeler, G Tsapralis, QM Chen, La autoantigen mediates oxidant induced de novo Nrf2 protein translation, *Molecular and Cellular Proteomics*. 11(M11) (2012) 015032.

- [176] RV Intine, SA Tenenbaum, AL Sakulich, JD Keene, RJ Maraia, Differential phosphorylation and subcellular localization of la RNPs associated with precursor tRNAs and translation-related mRNAs, *Molecular Cell*. 12 (2003) 1301–7. [http://dx.doi.org/10.1016/S1097-2765\(03\)00429-5](http://dx.doi.org/10.1016/S1097-2765(03)00429-5), 14636586.
- [177] F Brenet, ND Socci, N Sonenberg, EC Holland, Akt phosphorylation of la regulates specific mRNA translation in glial progenitors, *Oncogene*. 28 (2009) 128–39. <http://dx.doi.org/10.1038/ncr.2008.376>, 18836485.
- [178] CH Broekhuis, G Neubauer, der Heijden A van, M Mann, CG Proud, WJ van Venrooij, et al, Detailed analysis of the phosphorylation of the human la (SS-B) autoantigen. (De)phosphorylation does not affect its subcellular distribution, *Biochemistry (Moscow)*. 39 (2000) 3023–33. <http://dx.doi.org/10.1021/bi992308c>.
- [179] PW Vesely, PB Staber, G Hoefler, L Kenner, Translational regulation mechanisms of AP-1 proteins, *Mutation Research*. 682 (2009) 7–12. <http://dx.doi.org/10.1016/j.mmr.2009.01.001>, 19167516.
- [180] M Narasimhan, D Patel, D Vedpathak, M Rathinam, G Henderson, L Mahimainathan, Identification of novel microRNAs in post-transcriptional control of Nrf2 expression and redox homeostasis in neuronal, SH-SY5Y cells, *PLOS One*. 7 (2012) e51111. <http://dx.doi.org/10.1371/journal.pone.0051111>, 23236440.
- [181] O Perez-Leal, CA Barrero, S Merali, Translational control of Nrf2 within the open reading frame, *Biochemical and Biophysical Research Communications*. 437 (2013) 134–9. <http://dx.doi.org/10.1016/j.bbrc.2013.06.052>, 23806685.
- [182] PS Ray, R Grover, S Das, Two internal ribosome entry sites mediate the translation of p53 isoforms, *EMBO Reports*. 7 (2006) 404–10.
- [183] W Zhou, GM Edelman, VP Mauro, Transcript leader regions of two *Saccharomyces cerevisiae* mRNAs contain internal ribosome entry sites that function in living cells, *Proceedings of the National Academy of Sciences*. 98 (2001) 1531–6. <http://dx.doi.org/10.1073/pnas.98.4.1531>.
- [184] MH Glickman, A Ciechanover, The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction, *Physiological Reviews*. 82 (2002) 373–428.
- [185] F Kriegenburg, EG Poulsen, A Koch, E Krüger, R Hartmann-Petersen, Redox control of the ubiquitin-proteasome system: from molecular mechanisms to functional significance, *Antioxidants and Redox Signalling*. 15 (2011) 2265–99. <http://dx.doi.org/10.1089/ars.2010.3590>, 21314436.
- [186] A Kobayashi, M-I Kang, H Okawa, M Ohtsui, Y Zenke, T Chiba, et al, Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2, *Molecular and Cellular Biology*. 24 (2004) 7130–9. <http://dx.doi.org/10.1128/MCB.24.16.7130-7139.2004>, 15282312.
- [187] DD Zhang, S-C Lo, JV Cross, DJ Templeton, M Hannink, Keap1 is a Redox-regulated substrate adaptor protein for a Cul3-dependent ubiquitin ligase complex, *Molecular and Cellular Biology*. 24 (2004) 10941–53. <http://dx.doi.org/10.1128/MCB.24.24.10941-10953.2004>, 15572695.
- [188] L Baird, AT Dinkova-Kostova, Diffusion dynamics of the Keap1-Cullin3 interaction in single live cells, *Biochemical and Biophysical Research Communications*. 433 (2013) 58–65. <http://dx.doi.org/10.1016/j.bbrc.2013.02.065>, 23454126.
- [189] S Fourquet, R Guerois, D Biard, MB Toledano, Activation of NRF2 by nitrosative agents and H₂O₂ involves KEAP1 disulfide formation, *Journal of Biological Chemistry*. 285 (2010) 8463–71. <http://dx.doi.org/10.1074/jbc.M109.051714>, 20061377.
- [190] CL Brooks, W Gu, p53 ubiquitination: Mdm2 and beyond, *Molecular Cell*. 21 (2006) 307–15. <http://dx.doi.org/10.1016/j.molcel.2006.01.020>, 16455486.
- [191] D Michael, M Oren, The p53-Mdm2 module and the ubiquitin system, *Seminars in Cancer Biology*. 13 (2003) 49–58. [http://dx.doi.org/10.1016/S1044-579X\(02\)00099-8](http://dx.doi.org/10.1016/S1044-579X(02)00099-8), 12507556.
- [192] JD Oliner, JA Pietenpol, S Thiagalingam, J Gyuris, KW Kinzler, B Vogelstein, Oncoprotein MDM2 conceals the activation domain of tumour suppressor p53, *Nature*. 362 (1993) 857–60. <http://dx.doi.org/10.1038/362857a0>, 8479525.
- [193] Y Haupt, R Maya, A Kazaz, M Oren, Mdm2 promotes the rapid degradation of p53, *Nature*. 387 (1997) 296–9. <http://dx.doi.org/10.1038/387296a0>, 9153395.
- [194] R Honda, H Tanaka, H Yasuda, Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53, *FEBS Letters*. 420 (1997) 25–7. [http://dx.doi.org/10.1016/S0014-5793\(97\)01480-4](http://dx.doi.org/10.1016/S0014-5793(97)01480-4), 9450543.
- [195] MHG Kubbutat, SN Jones, KH Voudsen, Regulation of p53 stability by Mdm2, *Nature*. 387 (1997) 299–303. <http://dx.doi.org/10.1038/387299a0>, 9153396.
- [196] H Zhan, T Suzuki, K Aizawa, K Miyagawa, R Nagai, Ataxia telangiectasia mutated (ATM)-mediated DNA damage response in oxidative stress-induced vascular endothelial cell senescence, *Journal of Biological Chemistry*. 285 (2010) 29662–70. <http://dx.doi.org/10.1074/jbc.M110.125138>, 20639198.
- [197] X Sun, P Majumder, H Shioya, F Wu, S Kumar, R Weichselbaum, et al, Activation of the cytoplasmic c-Abl tyrosine kinase by reactive oxygen species, *Journal of Biological Chemistry*. 275 (2000) 17237–40. <http://dx.doi.org/10.1074/jbc.C000099200>, 10770918.
- [198] R Maya, M Balass, S-T Kim, D Shkedy, J-FM Leal, O Shifman, et al, ATM-dependent phosphorylation of Mdm2 on serine 395: Role in p53 activation by DNA damage, *Genes and Development*. 15 (2001) 1067–77. <http://dx.doi.org/10.1101/gad.886901>, 11331603.
- [199] Z Goldberg, Tyrosine phosphorylation of Mdm2 by c-Abl: implications for p53 regulation, *EMBO Journal*. 21 (2002) 3715–27.
- [200] T Buschmann, SY Fuchs, CG Lee, ZQ Pan, Z Ronai, SUMO-1 modification of Mdm2 prevents its self-ubiquitination and increases Mdm2 ability to ubiquitinate p53, *Cell*. 101 (2000) 753–62. [http://dx.doi.org/10.1016/S0092-8674\(00\)80887-9](http://dx.doi.org/10.1016/S0092-8674(00)80887-9), 10892746.
- [201] G Bossis, F Melchior, Regulation of SUMOylation by reversible oxidation of SUMO conjugating enzymes, *Molecular Cell*. 21 (2006) 349–57. <http://dx.doi.org/10.1016/j.molcel.2005.12.019>, 16455490.
- [202] De la Vega I, I Grishina, R Moreno, M Krüger, T Braun, ML Schmitz, A Redox-regulated SUMO/acetylation switch of HIPK2 controls the survival threshold to oxidative stress, *Molecular Cell*. 46 (2012) 472–83. <http://dx.doi.org/10.1016/j.molcel.2012.03.003>, 22503103.
- [203] X Wang, J Wang, X Jiang, MdmX protein is essential for Mdm2 protein-mediated p53 polyubiquitination, *Journal of Biological Chemistry*. 286 (2011) 23725–34. <http://dx.doi.org/10.1074/jbc.M110.213868>, 21572037.
- [204] K Okamoto, Y Taya, H Nakagama, Mdmx enhances p53 ubiquitination by altering the substrate preference of the Mdm2 ubiquitin ligase, *FEBS Letters*. 583 (2009) 2710–14. <http://dx.doi.org/10.1016/j.febslet.2009.07.021>, 19619542.
- [205] V Zuckerman, K Lenos, GM Popowicz, I Silberman, T Grossman, J-C Marine, et al, c-Abl phosphorylates Mdmx and regulates its interaction with p53, *Journal of Biological Chemistry*. 284 (2009) 4031–9. <http://dx.doi.org/10.1074/jbc.M809211200>, 19075013.
- [206] V Lopez-Pajares, MM Kim, Z-M Yuan, Phosphorylation of MDMX mediated by Akt leads to stabilization and Induces 14-3-3 Binding, *Journal of Biological Chemistry*. 283 (2008) 13707–13. <http://dx.doi.org/10.1074/jbc.M710030200>, 18356162.
- [207] L Chen, DM Gilkes, Y Pan, WS Lane, J Chen, ATM and Chk2-dependent phosphorylation of MDMX contribute to p53 activation after DNA damage, *EMBO Journal*. 24 (2005) 3411–22.
- [208] H Zhao, F Tragano, AP Albino, Z Darzynkiewicz, Oxidative stress induces cell cycle-dependent Mre11 recruitment, ATM and Chk2 activation and histone H2AX phosphorylation, *Cell Cycle*. 7 (2008) 1490–5. <http://dx.doi.org/10.4161/cc.7.10.5963>, 18418078.
- [209] S Bedri, SM Cizek, I Rastarhuyeva, JR Stone, Regulation of protein kinase CK1α/β by dephosphorylation in response to hydrogen peroxide, *Archives of Biochemistry and Biophysics*. 466 (2007) 242–9. <http://dx.doi.org/10.1016/j.jabb.2007.06.010>, 17626781.
- [210] M Salazar, AI Rojo, D Velasco, RM de Sagarra, A Cuadrado, Glycogen synthase kinase-3β inhibits the xenobiotic and antioxidant cell response by direct phosphorylation and nuclear exclusion of the transcription factor Nrf2, *Journal of Biological Chemistry*. 281 (2006) 14841–51. <http://dx.doi.org/10.1074/jbc.M513737200>, 16551619.
- [211] P Rada, AI Rojo, S Chowdhry, M McMahon, JD Hayes, A Cuadrado, SCF/β-TCP promotes glycogen synthase kinase 3-dependent degradation of the Nrf2 transcription factor in a Keap1-independent manner, *Molecular and Cellular Biology*. 31 (2011) 1121–33. <http://dx.doi.org/10.1128/MCB.01204-10>, 21245377.
- [212] DAE Cross, DR Alessi, P Cohen, M Andjelkovich, BA Hemmings, Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B, *Nature*. 378 (1995) 785–9. <http://dx.doi.org/10.1038/378785a0>, 8524413.
- [213] M Shaw, P Cohen, DR Alessi, The activation of protein kinase B by H₂O₂ or heat shock is mediated by phosphoinositide 3-kinase and not by mitogen-activated protein kinase-activated protein kinase-2, *Biochemical Journal*. 336 (1998) 241–6.
- [214] AS Blair, E Hajdich, GJ Litherland, HS Hundal, Regulation of glucose transport and glycogen synthesis in L6 muscle cells during oxidative stress. Evidence for cross-talk between the insulin and SAPK2/p38 mitogen-activated protein kinase signaling pathways, *Journal of Biological Chemistry*. 274 (1999) 36293–9. <http://dx.doi.org/10.1074/jbc.274.51.36293>, 10593919.
- [215] AI Rojo, MR Sagarra, A Cuadrado, GSK-3β down-regulates the transcription factor Nrf2 after oxidant damage: relevance to exposure of neuronal cells to oxidative stress, *Journal of Neurochemistry*. 105 (2008) 192–202. <http://dx.doi.org/10.1111/j.1471-4159.2007.05124.x>, 18005231.
- [216] O Kaidanovich-Beilin, JR Woodgett, GSK-3: Functional insights from cell biology and animal models, *Frontiers in Molecular Neuroscience*. 4 (2011).
- [217] S Chowdhry, Y Zhang, M McMahon, C Sutherland, A Cuadrado, JD Hayes, Nrf2 is controlled by two distinct β-TrCP recognition motifs in its Neh6 domain, one of which can be modulated by GSK-3 activity, *Oncogene*. 32 (2013) 3765–81. <http://dx.doi.org/10.1038/ncr.2012.388>, 22964642.
- [218] M Ushio-Fukai, RW Alexander, M Akers, KK Griendling, p38 mitogen-activated protein kinase is a critical component of the redox-sensitive signaling pathways activated by angiotensin II. Role in vascular smooth muscle cell hypertrophy, *Journal of Biological Chemistry*. 273 (1998) 15022–9. <http://dx.doi.org/10.1074/jbc.273.24.15022>, 9614110.
- [219] R Aikawa, I Komuro, T Yamazaki, Y Zou, S Kudoh, M Tanaka, et al, Oxidative stress activates extracellular signal-regulated kinases through Src and Ras in cultured cardiac myocytes of neonatal rats, *Journal of Clinical Investigation*. 100 (1997) 1813–21. <http://dx.doi.org/10.1172/JCI119709>, 9312182.
- [220] H Konishi, M Tanaka, Y Takemura, H Matsuzaki, Y Ono, U Kikkawa, et al, Activation of protein kinase C by tyrosine phosphorylation in response to H₂O₂, *Proceedings of the National Academy of Sciences*. 94 (1997) 11233–7. <http://dx.doi.org/10.1073/pnas.94.21.11233>.
- [221] AK Jain, AK Jaiswal, GSK-3β acts upstream of Fyn kinase in regulation of nuclear export and degradation of NF-E2 related factor 2, *Journal of Biological Chemistry*. 282 (2007) 16502–10. <http://dx.doi.org/10.1074/jbc.M611336200>, 17403689.
- [222] Y Feng, Y Xia, G Yu, X Shu, H Ge, K Zeng, et al, Cleavage of GSK-3β by calpain counteracts the inhibitory effect of Ser9 phosphorylation on GSK-3β activity induced by H₂O₂, *Journal of Neurochemistry*. 126 (2013) 234–42. <http://dx.doi.org/10.1111/jnc.12285>, 23646926.
- [223] SB Cullinan, D Zhang, M Hannink, E Arvisais, RJ Kaufman, JA Diehl, Nrf2 is a direct PERK substrate and effector of PERK-dependent cell survival, *Molecular and Cellular Biology*. 23 (2003) 7198–209. <http://dx.doi.org/10.1128/MCB.23.20.7198-7209.2003>, 14517290.
- [224] P Palapati, DA Averill-Bates, Activation of ER stress and apoptosis by hydrogen

- peroxide in HeLa cells: protective role of mild heat preconditioning at 40 °C, *Biochimica et Biophysica Acta Molecular Cell Research*. 1813 (2011) 1987–99. <http://dx.doi.org/10.1016/j.bbamcr.2011.07.021>.
- [225] SK Niture, AK Jain, AK Jaiswal, Antioxidant-induced modification of Nrf2 cysteine 151 and PKC-delta-mediated phosphorylation of Nrf2 serine 40 are both required for stabilization and nuclear translocation of Nrf2 and increased drug resistance, *Journal of Cell Science*. 122 (2009) 4452–64. <http://dx.doi.org/10.1242/jcs.058537>, 19920073.
- [226] AM Musti, M Treier, D Bohmann, Reduced ubiquitin-dependent degradation of c-Jun after phosphorylation by MAP kinases, *Science (New York, N.Y.)*. 275 (1997) 400–2.
- [227] N Ozgen, J Guo, Z Gertsberg, P Danilo, MR Rosen, SF Steinberg, Reactive oxygen species decrease cAMP response element binding protein expression in cardiomyocytes via a protein kinase D1-dependent mechanism that does not require Ser133 phosphorylation, *Molecular Pharmacology*. 76 (2009) 896–902. <http://dx.doi.org/10.1124/mol.109.056473>, 19620255.
- [228] B Gao, S-M Lee, D Fang, The tyrosine kinase c-Abl protects c-Jun from ubiquitination-mediated degradation in T cells, *Journal of Biological Chemistry*. 281 (2006) 29711–18. <http://dx.doi.org/10.1074/jbc.M604596200>, 16901904.
- [229] M Xie, K Sabapathy, Tyrosine 170 is dispensable for c-Jun turnover, *Cellular Signalling*. 22 (2010) 330–7. <http://dx.doi.org/10.1016/j.cellsig.2009.09.040>, 19818398.
- [230] JH Marxsen, P Stengel, K Doege, P Heikkinen, T Jokilehto, T Wagner, et al, Hypoxia-inducible factor-1 (HIF-1) promotes its degradation by induction of HIF- α -prolyl-4-hydroxylases, *Biochemical Journal*. 381 (2004) 761–7. <http://dx.doi.org/10.1042/BJ20040620>, 15104534.
- [231] Y Pan, KD Mansfield, CC Bertozzi, V Rudenko, DA Chan, AJ Giaccia, et al, Multiple factors affecting cellular redox status and energy metabolism modulate hypoxia-inducible factor prolyl hydroxylase activity in vivo and in vitro, *Molecular and Cellular Biology*. 27 (2007) 912–25. <http://dx.doi.org/10.1128/MCB.01223-06>, 17101781.
- [232] D Gerald, E Berra, YM Frapart, DA Chan, AJ Giaccia, D Mansuy, et al, D reduces tumor angiogenesis by protecting cells from oxidative stress, *Cell*. 118 (2004) 781–94. <http://dx.doi.org/10.1016/j.cell.2004.08.025>, 15369676.
- [233] P Goyal, N Weissmann, F Grimminger, C Hegel, L Bader, F Rose, et al, Upregulation of NAD(P)H oxidase 1 in hypoxia activates hypoxia-inducible factor 1 via increase in reactive oxygen species, *Free Radical Biology and Medicine*. 36 (2004) 1279–88. <http://dx.doi.org/10.1016/j.freeradbiomed.2004.02.071>, 15110393.
- [234] RD Guzy, B Hoyos, E Robin, H Chen, L Liu, KD Mansfield, et al, Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing, *Cell Metabolism*. 1 (2005) 401–8. <http://dx.doi.org/10.1016/j.cmet.2005.05.001>, 16054089.
- [235] KD Mansfield, RD Guzy, Y Pan, RM Young, TP Cash, PT Schumacker, et al, Mitochondrial dysfunction resulting from loss of cytochrome c impairs cellular oxygen sensing and hypoxic HIF- α activation, *Cell Metabolism*. 1 (2005) 393–9. <http://dx.doi.org/10.1016/j.cmet.2005.05.003>, 16054088.
- [236] S Naranjo-Suarez, BA Carlson, PA Tsuji, M-H Yoo, VN Gladyshev, DL Hatfield, HIF-independent regulation of thioredoxin reductase 1 contributes to the high levels of reactive oxygen species induced by hypoxia, *PLOS One*. 7 (2012) e30470. <http://dx.doi.org/10.1371/journal.pone.0030470>, 22348009.
- [237] J Zhou, B Brüne, Cytokines and hormones in the regulation of hypoxia inducible factor-1 α (HIF-1 α), *Cardiovascular and Hematological Agents in Medicinal Chemistry*. 4 (2006) 189–97.
- [238] DE Richard, E Berra, J Pouyssegur, Nonhypoxic pathway mediates the induction of hypoxia-inducible factor 1 α in vascular smooth muscle cells, *Journal of Biological Chemistry*. 275 (2000) 26765–71.
- [239] JJ Haddad, SC Land, A non-hypoxic, ROS-sensitive pathway mediates TNF- α -dependent regulation of HIF-1 α , *FEBS Letters*. 505 (2001) 269–74. [http://dx.doi.org/10.1016/S0014-5793\(01\)02833-2](http://dx.doi.org/10.1016/S0014-5793(01)02833-2).
- [240] D Zhang, T Suganuma, JL Workman, Acetylation regulates Jun protein turnover in *Drosophila*, *Biochimica et Biophysica Acta – Gene Regulatory Mechanisms*. 1829 (2013) 1218–24.
- [241] S-B Jung, C-S Kim, Y-R Kim, A Naqvi, T Yamamori, S Kumar, et al, Redox factor-1 activates endothelial SIRTUIN1 through reduction of conserved cysteine sulphydryls in its deacetylase Domain, *PLOS One*. 8 (2013) e65415.
- [242] SG Ahn, DJ Thiele, Redox regulation of mammalian heat shock factor 1 is essential for Hsp gene activation and protection from stress, *Genes and Development*. 17 (2003) 516–28. <http://dx.doi.org/10.1101/gad.1044503>, 12600944.
- [243] S Boissnard, G Lagniel, C Garmendia-Torres, M Molin, E Boy-Marcotte, M Jacquet, et al, H₂O₂ activates the nuclear localization of Msn2 and Maf1 through thioredoxins in *Saccharomyces cerevisiae*, *Eukaryotic Cell*. 8 (2009) 1429–38. <http://dx.doi.org/10.1128/EC.00106-09>, 19581440.
- [244] J Lee, RD Moir, IM Willis, Regulation of RNA polymerase III transcription involves SCH9-dependent and SCH9-independent branches of the target of rapamycin (TOR) pathway, *Journal of Biological Chemistry*. 284 (2009) 12604–8. <http://dx.doi.org/10.1074/jbc.C900020200>, 19299514.
- [245] YV Budovskaya, JS Stephan, SJ Deminoff, PK Herman, An evolutionary proteomics approach identifies substrates of the cAMP-dependent protein kinase, *Proceedings of the National Academy of Sciences of the United States of America*. 102 (2005) 13933–8. <http://dx.doi.org/10.1073/pnas.0501046102>, 16172400.
- [246] RD Moir, J Lee, RA Haeusler, N Desai, DR Engelke, IM Willis, Protein kinase A regulates RNA polymerase III transcription through the nuclear localization of Maf1, *Proceedings of the National Academy of Sciences of the United States of America*. 103 (2006) 15044–9. <http://dx.doi.org/10.1073/pnas.0607129103>, 17005718.
- [247] Y Wei, XF Zheng, TORC1 association with rDNA chromatin as a mechanism to co-regulate Pol I and Pol III, *Cell Cycle*. 8 (2009) 3802–3. <http://dx.doi.org/10.4161/cc.8.23.10039>, 19934660.
- [248] D Oficjalska-Pham, O Harismendy, WJ Smagowicz, Gonzalez de Peredo A, M Boguta, A Sentenac, et al, General repression of RNA polymerase III transcription is triggered by protein phosphatase type 2A-mediated dephosphorylation of Maf1, *Molecular Cell*. 22 (2006) 623–32. <http://dx.doi.org/10.1016/j.molcel.2006.04.008>, 16762835.
- [249] De Wever V, W Reiter, A Ballarini, G Ammerer, C Brocard, A dual role for PP1 in shaping the Msn2-dependent transcriptional response to glucose starvation, *EMBO Journal*. 24 (2005) 4115–23.
- [250] W Reiter, E Klopff, De Wever V, D Anrather, A Petryshyn, A Roetzer, et al, Yeast protein phosphatase 2A-Cdc55 regulates the transcriptional response to hyperosmolarity stress by regulating Msn2 and Msn4 chromatin recruitment, *Molecular and Cellular Biology*. 33 (2013) 1057–72. <http://dx.doi.org/10.1128/MCB.00834-12>, 23275436.
- [251] K Gulshan, Oxidant-specific folding of Yap1p regulates both transcriptional activation and nuclear localization, *Journal of Biological Chemistry*. 280 (2005) 40524–33. <http://dx.doi.org/10.1074/jbc.M504716200>, 16219769.
- [252] S Kuge, T Toda, N Iizuka, A Nomoto, Crm1 (Xpo1) dependent nuclear export of the budding yeast transcription factor yAP-1 is sensitive to oxidative stress, *Genes to Cells: Devoted to Molecular and Cellular Mechanisms*. 3 (1998) 521–32. <http://dx.doi.org/10.1046/j.1365-2443.1998.00209.x>, 9797454.
- [253] T Isoyama, Nuclear import of the Yeast AP-1-like transcription factor Yap1p is mediated by transport receptor Pse1p, and this import step is not affected by oxidative stress, *Journal of Biological Chemistry*. 276 (2001) 21863–9. <http://dx.doi.org/10.1074/jbc.M009528200>, 11274141.
- [254] S Okazaki, T Tachibana, A Naganuma, N Mano, S Kuge, Multistep disulfide bond formation in Yap1 is required for sensing and transduction of H₂O₂ stress signal, *Molecular Cell*. 27 (2007) 675–88. <http://dx.doi.org/10.1016/j.molcel.2007.06.035>, 17707237.
- [255] C Yan, Crm1p mediates regulated nuclear export of a yeast AP-1-like transcription factor, *EMBO Journal*. 17 (1998) 7416–29.
- [256] A Delaunay, D Pflieger, MB Barrault, J Vinh, MB Toledano, A thiol peroxidase is an H₂O₂ receptor and redox-transducer in gene activation, *Cell*. 111 (2002) 471–81. [http://dx.doi.org/10.1016/S0092-8674\(02\)01048-6](http://dx.doi.org/10.1016/S0092-8674(02)01048-6), 12437921.
- [257] D'Autrèaux B, MB Toledano, ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis, *Nature Reviews Molecular Cell Biology*. 8 (2007) 813–24. <http://dx.doi.org/10.1038/nrm2256>, 17848967.
- [258] EA Veal, Ybp1 is Required for the hydrogen peroxide-induced oxidation of the Yap1 transcription factor, *Journal of Biological Chemistry*. 278 (2003) 30896–904. <http://dx.doi.org/10.1074/jbc.M303542200>, 12743123.
- [259] K Gulshan, SA Rovinsky, WS Moye-Rowley, YBP1 and its homologue YBP2/YBH1 influence oxidative-stress tolerance by nonidentical mechanisms in *Saccharomyces cerevisiae*, *Eukaryotic Cell*. 3 (2004) 318–30. <http://dx.doi.org/10.1128/EC.3.2.318-330.2004>, 15075262.
- [260] W Li, S-W Yu, A-NT Kong, Nrf2 possesses a redox-sensitive nuclear exporting signal in the Neh5 transactivation domain, *Journal of Biological Chemistry*. 281 (2006) 27251–63. <http://dx.doi.org/10.1074/jbc.M602746200>, 16790425.
- [261] W Li, MR Jain, C Chen, X Yue, V Hebbbar, R Zhou, et al, Nrf2 possesses a redox-insensitive nuclear export signal overlapping with the leucine zipper motif, *Journal of Biological Chemistry*. 280 (2005) 28430–8. <http://dx.doi.org/10.1074/jbc.M410601200>, 15917227.
- [262] AK Jain, AK Jaiswal, Phosphorylation of tyrosine 568 controls nuclear export of Nrf2, *Journal of Biological Chemistry*. 281 (2006) 12132–42. <http://dx.doi.org/10.1074/jbc.M511198200>, 16513647.
- [263] G Gloire, S Legrand-Poels, J Piette, NF-kappaB activation by reactive oxygen species: fifteen years later, *Biochemical Pharmacology*. 72 (2006) 1493–505. <http://dx.doi.org/10.1016/j.bcp.2006.04.011>, 16723122.
- [264] JR Woo, SJ Kim, W Jeong, YH Cho, SC Lee, YJ Chung, et al, Structural basis of cellular redox regulation by human TRP14, *Journal of Biological Chemistry*. 279 (2004) 48120–5. <http://dx.doi.org/10.1074/jbc.M407079200>, 15355959.
- [265] Y Jung, H Kim, SH Min, SG Rhee, W Jeong, Dynein light chain LC8 negatively regulates NF-kappaB through the redox-dependent interaction with IkappaBalpha, *Journal of Biological Chemistry*. 283 (2008) 23863–71. <http://dx.doi.org/10.1074/jbc.M803072200>, 18579519.
- [266] D-F Lee, H-P Kuo, M Liu, C-K Chou, W Xia, Y Du, et al, KEAP1 E3 ligase-mediated down-regulation of NF-kB signaling by targeting IKK, *Molecular Cell*. 36 (2009) 131–40. <http://dx.doi.org/10.1016/j.molcel.2009.07.025>, 19818716.
- [267] AK Jain, DA Bloom, AK Jaiswal, Nuclear import and export signals in control of Nrf2, *Journal of Biological Chemistry*. 280 (2005) 29158–68. <http://dx.doi.org/10.1074/jbc.M502083200>, 15901726.
- [268] M Theodore, Y Kawai, J Yang, Y Kleshchenko, SP Reddy, F Villalta, et al, Multiple nuclear localization signals function in the nuclear import of the transcription factor Nrf2, *Journal of Biological Chemistry*. 283 (2008) 8984–94. <http://dx.doi.org/10.1074/jbc.M709040200>, 18238777.
- [269] Y Watai, A Kobayashi, H Nagase, M Mizukami, J McEvoy, JD Singer, et al, Subcellular localization and cytoplasmic complex status of endogenous Keap1, *Genes to Cells: Devoted to Molecular and Cellular Mechanisms*. 12 (2007) 1163–78. <http://dx.doi.org/10.1111/j.1365-2443.2007.01118.x>, 17903176.
- [270] M Velichkova, T Hasson, Keap1 regulates the oxidation-sensitive shuttling of Nrf2 into and out of the nucleus via a Crm1-dependent nuclear export mechanism, *Molecular and Cellular Biology*. 25 (2005) 4501–13. <http://dx.doi.org/10.1128/MCB.25.11.4501-4513.2005>, 15899855.
- [271] Z Sun, S Zhang, JY Chan, DD Zhang, Keap1 controls postinduction repression of

- the Nrf2-mediated antioxidant response by escorting nuclear export of Nrf2, *Molecular and Cellular Biology*. 27 (2007) 6334–49. <http://dx.doi.org/10.1128/MCB.00630-07.17636022>.
- [272] BJ Buckley, S Li, AR Whorton, Keap1 modification and nuclear accumulation in response to S-nitrosocysteine, *Free Radical Biology and Medicine*. 44 (2008) 692–8. <http://dx.doi.org/10.1016/j.freeradbiomed.2007.10.055>, 18062931.
- [273] JW Kaspar, SK Niture, AK Jaiswal, Antioxidant-induced Nrf2 (Keap1) tyrosine 85 phosphorylation controls the nuclear export and degradation of the Irf2-Cul3-Rbx1 complex to allow normal Nrf2 activation and repression, *Journal of Cell Science*. 125 (2012) 1027–38. <http://dx.doi.org/10.1242/jcs.097295>, 22448038.
- [274] T Sugano, M Nitta, H Ohmori, M Yamaizumi, Nuclear accumulation of p53 in normal human fibroblasts is induced by various cellular stresses which evoke the heat shock response, independently of the cell cycle, *Japanese Journal of Cancer Research: Gann*. 86 (1995) 415–18. <http://dx.doi.org/10.1111/j.1349-7006.1995.tb03072.x>, 7790313.
- [275] O'Brate A, P Giannakakou, The importance of p53 location: nuclear or cytoplasmic zip code? Drug Resistance Updates: Reviews and Commentaries in Antimicrobial and Anticancer Chemotherapy. 6 (2003) 313–22. <http://dx.doi.org/10.1016/j.drug.2003.10.004>, 14744495.
- [276] D Xirodimas, MK Saville, C Edling, DP Lane, S Lain, Different effects of p14ARF on the levels of ubiquitinated p53 and Mdm2 in vivo, *Oncogene*. 20 (2001) 4972–83. <http://dx.doi.org/10.1038/sj.onc.1204656>, 11526482.
- [277] JD Weber, LJ Taylor, MF Roussel, CJ Sherr, D Bar-Sagi, Nucleolar arf sequesters Mdm2 and activates p53, *Nature Cell Biology*. 1 (1999) 20–6. <http://dx.doi.org/10.1038/8991>, 10559859.
- [278] LD Mayo, DB Donner, A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus, *Proceedings of the National Academy of Sciences*. 98 (2001) 11598–603. <http://dx.doi.org/10.1073/pnas.181181198>.
- [279] Y Ogawara, Akt enhances Mdm2-mediated ubiquitination and Degradation of p53, *Journal of Biological Chemistry*. 277 (2002) 21843–50. <http://dx.doi.org/10.1074/jbc.M109745200>, 11923280.
- [280] V Stambolic, D MacPherson, D Sas, Y Lin, B Snow, Y Jang, et al, Regulation of PTEN transcription by p53, *Molecular Cell*. 8 (2001) 317–25. [http://dx.doi.org/10.1016/S1097-2765\(01\)00323-9](http://dx.doi.org/10.1016/S1097-2765(01)00323-9), 11545734.
- [281] A Béghin, E-L Matera, S Brunet-Manquat, C Dumontet, Expression of Arl2 is associated with p53 localization and chemosensitivity in a breast cancer cell line, *Cell Cycle*. 7 (2008) 3074–82. <http://dx.doi.org/10.4161/cc.7.19.6777>, 18818514.
- [282] N Coant, Ben Mkaddem S, E Pedruzzi, C Guichard, X Tréton, R Ducroc, et al, NADPH oxidase 1 modulates WNT and NOTCH1 signaling to control the fate of proliferative progenitor cells in the colon, *Molecular and Cellular Biology*. 30 (2010) 2636–50. <http://dx.doi.org/10.1128/MCB.01194-09>, 20351171.
- [283] J-H Zhu, C-L Chen, S Flavahan, J Harr, B Su, NA Flavahan, Cyclic stretch stimulates vascular smooth muscle cell injury by redox-dependent activation of Notch3, *American Journal of Physiology – Heart and Circulatory Physiology*. 300 (2011) H1770–H1780. <http://dx.doi.org/10.1152/ajpheart.00535.2010>, 21169401.
- [284] Z Zhang, P Oliver, JR Lancaster Jr, PO Schwarzenberger, MS Joshi, J Cork, et al, Reactive oxygen species mediate tumor necrosis factor alpha-converting, enzyme-dependent ectodomain shedding induced by phorbol myristate acetate, *F.A.S.E.B. Journal*. 15 (2001) 303–5.
- [285] Y Wang, AH Herrera, Y Li, KK Belani, B Walcheck, Regulation of mature ADAM17 by redox agents for L-selectin shedding, *Journal of Immunology*. 182 (2009) 2449–57. <http://dx.doi.org/10.4049/jimmunol.0802770>, 19201900.
- [286] BJ Goldstein, K Mahadev, X Wu, L Zhu, H Motoshima, Role of insulin-induced reactive oxygen species in the insulin signaling Pathway, *Antioxidants and Redox Signalling*. 7 (2005) 1021–31. <http://dx.doi.org/10.1089/ars.2005.7.1021>, 15998257.
- [287] KD Tardif, G Waris, A Siddiqui, Hepatitis C virus, ER stress, and oxidative stress, *Trends in Microbiology*. 13 (2005) 159–63. <http://dx.doi.org/10.1016/j.tim.2005.02.004>, 15817385.
- [288] MSB Paget, MJ Buttner, Thiol-based regulatory switches, *Annual Review of Genetics*. 37 (2003) 91–121. <http://dx.doi.org/10.1146/annurev.genet.37.110801.142538>, 14616057.
- [289] F Aslund, M Zheng, J Beckwith, G Storz, Regulation of the OxyR transcription factor by hydrogen peroxide and the cellular thiol-disulfide status, *Proceedings of the National Academy of Sciences of the United States of America*. 96 (1999) 6161–5. <http://dx.doi.org/10.1073/pnas.96.11.6161>, 10339558.
- [290] K Tao, In vivo oxidation-reduction kinetics of OxyR, the transcriptional activator for an oxidative stress-inducible regulon in *Escherichia coli*, *FEBS Letters*. 457 (1999) 90–2. [http://dx.doi.org/10.1016/S0014-5793\(99\)01013-3](http://dx.doi.org/10.1016/S0014-5793(99)01013-3).
- [291] C Lee, SM Lee, P Mukhopadhyay, SJ Kim, SC Lee, W-S Ahn, et al, Redox regulation of OxyR requires specific disulfide bond formation involving a rapid kinetic reaction path, *Nature Structural and Molecular Biology*. 11 (2004) 1179–85. <http://dx.doi.org/10.1038/nsmb856>, 15543158.
- [292] SO Kim, K Merchant, R Nudelmann, WF Beyer, T Keng, J DeAngelo, et al, OxyR: a molecular code for redox-related signaling, *Cell*. 109 (2002) 383–96. [http://dx.doi.org/10.1016/S0092-8674\(02\)00723-7](http://dx.doi.org/10.1016/S0092-8674(02)00723-7), 12015987.
- [293] D Seth, A Hausladen, Y-J Wang, JS Stamler, Endogenous protein S-nitrosylation in *E. coli*: regulation by OxyR, *Science*. 336 (2012) 470–3.
- [294] H Xiao, O Perisic, JT Lis, Cooperative binding of *Drosophila* heat shock factor to arrays of a conserved 5 bp unit, *Cell*. 64 (1991) 585–93. [http://dx.doi.org/10.1016/0092-8674\(91\)90242-Q](http://dx.doi.org/10.1016/0092-8674(91)90242-Q), 1899357.
- [295] M Lu, HE Kim, CR Li, S Kim, IJ Kwak, YJ Lee, et al, Two distinct disulfide bonds formed in human heat shock transcription factor 1 act in opposition to regulate its DNA binding activity, *Biochemistry (Moscow)*. 47 (2008) 6007–15.
- [296] DA Traoré, El Ghazouani A, S Ilango, J Dupuy, L Jacquamet, JL Ferrer, et al, Crystal structure of the apo-PerR-Zn protein from *Bacillus subtilis*, *Molecular Microbiology*. 61 (2006) 1211–19. <http://dx.doi.org/10.1111/j.1365-2958.2006.05313.x>, 16925555.
- [297] DA Traoré, L Jacquamet, F Borel, JL Ferrer, D Lascoux, JL Ravanat, et al, Structural and functional characterization of 2-oxo-histidine in oxidized PerR protein, *Nature Chemical Biology*. 5 (2009) 53–9. <http://dx.doi.org/10.1038/nchembio.133>, 19079268.
- [298] D Bloom, S Dhakshinamoorthy, AK Jaiswal, Site-directed mutagenesis of cysteine to serine in the DNA binding region of Nrf2 decreases its capacity to up-regulate antioxidant response element-mediated expression and antioxidant induction of NAD(P)H:quinone oxidoreductase1 gene, *Oncogene*. 21 (2002) 2191–200. <http://dx.doi.org/10.1038/sj.onc.1205288>, 11948402.
- [299] N Makthal, S Rastegari, M Sanson, Z Ma, RJ Olsen, JD Helmann, et al, Crystal structure of peroxide stress regulator from *Streptococcus pyogenes* provides functional insights into the mechanism of oxidative stress sensing, *Journal of Biological Chemistry*. 288 (2013) 18311–24. <http://dx.doi.org/10.1074/jbc.M113.456590>, 23645680.
- [300] JW Lee, JD Helmann, Biochemical characterization of the structural Zn²⁺ site in the *Bacillus subtilis* peroxide sensor PerR, *Journal of Biological Chemistry*. 281 (2006) 23567–78. <http://dx.doi.org/10.1074/jbc.M603968200>, 16766519.
- [301] L Jacquamet, DA Traoré, JL Ferrer, O Proux, D Testemale, JL Hazemann, et al, Structural characterization of the active form of PerR: Insights into the metal-induced activation of PerR and Fur proteins for DNA binding, *Molecular Microbiology*. 73 (2009) 20–31. <http://dx.doi.org/10.1111/j.1365-2958.2009.06753.x>, 19508285.
- [302] K Uchida, S Kawakishi, Identification of oxidized histidine generated at the active site of Cu,Zn-superoxide dismutase exposed to H₂O₂. Selective generation of 2-oxo-histidine at the histidine 118, *Journal of Biological Chemistry*. 269 (1994) 2405–10.
- [303] C Schöneich, Mechanisms of metal-catalyzed oxidation of histidine to 2-oxo-histidine in peptides and proteins, *Journal of Pharmaceutical and Biomedical Analysis*. 21 (2000) 1093–7. [http://dx.doi.org/10.1016/S0731-7085\(99\)00182-X](http://dx.doi.org/10.1016/S0731-7085(99)00182-X), 10708394.
- [304] MJ Faulkner, JD Helmann, Peroxide stress elicits adaptive changes in bacterial metal ion homeostasis, *Antioxidants and Redox Signalling*. 15 (2011) 175–89. <http://dx.doi.org/10.1089/ars.2010.3682>, 20977351.
- [305] R Ammendola, M Mesuraca, T Russo, F Cimino, The DNA-binding efficiency of Sp1 is affected by redox changes, *European Journal of Biochemistry*. 225 (1994) 483–9. <http://dx.doi.org/10.1111/j.1432-1033.1994.t01-1-00483.x>, 7925470.
- [306] L Jornt, H Petersen, AF Junod, Modulation of the DNA binding activity of transcription factors CREB, NF kappaB and HSF by H₂O₂ and TNF alpha. Differences between in vivo and in vitro effects, *F.E.B.S. Letters*. 416 (1997) 381–6. [http://dx.doi.org/10.1016/S0014-5793\(97\)01244-1](http://dx.doi.org/10.1016/S0014-5793(97)01244-1).
- [307] S Xanthoudakis, G Miao, F Wang, YC Pan, T Curran, Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme, *EMBO Journal*. 11 (1992) 3323–35.
- [308] B Dimple, T Herman, DS Chen, Cloning and expression of APE, the cDNA encoding the major human apurinic endonuclease: definition of a family of DNA repair enzymes, *Proceedings of the National Academy of Sciences of the United States of America*. 88 (1991) 11450–4. <http://dx.doi.org/10.1073/pnas.88.24.11450>, 1722334.
- [309] JS Sung, B Dimple, Roles of base excision repair subpathways in correcting oxidized abasic sites in DNA, *FEBS Journal*. 273 (2006) 1620–9.
- [310] G Tell, F Quadrioglio, C Tiribelli, MR Kelley, The many functions of APE1/Ref-1: not only a DNA repair enzyme, *Antioxidants and Redox Signalling*. 11 (2009) 601–20. <http://dx.doi.org/10.1089/ars.2008.2194>.
- [311] M Luo, S Delaplane, A Jiang, A Reed, Y He, M Fishel, et al, Role of the multi-functional DNA repair and redox signaling protein Ape1/ref-1 in cancer and endothelial cells: small-molecule inhibition of the redox function of Ape1, *Antioxidants and Redox Signalling*. 10 (2008) 1853–67. <http://dx.doi.org/10.1089/ars.2008.2120>, 18627350.
- [312] CD Mol, T Izumi, S Mitra, JA Tainer, DNA-bound structures and mutants reveal abasic DNA binding by APE1 and DNA repair coordination [corrected], *Nature*. 403 (2000) 451–6. <http://dx.doi.org/10.1038/35000249>, 10667800.
- [313] RP Huang, ED Adamson, Characterization of the DNA-binding properties of the early growth response-1 (Egr-1) transcription factor: Evidence for modulation by a redox mechanism, *DNA and Cell Biology*. 12 (1993) 265–73. <http://dx.doi.org/10.1089/dna.1993.12.265>, 8466649.
- [314] LE Huang, Z Arany, DM Livingston, HF Bunn, Activation of hypoxia-inducible transcription factor depends primarily upon redox-sensitive stabilization of its alpha subunit, *Journal of Biological Chemistry*. 271 (1996) 32253–9. <http://dx.doi.org/10.1074/jbc.271.50.32253>, 8943284.
- [315] H Nakshatri, P Bhat-Nakshatri, RA Currie, Subunit association and DNA binding activity of the heterotrimeric transcription factor NF-Y is regulated by cellular redox, *Journal of Biological Chemistry*. 271 (1996) 28784–91. <http://dx.doi.org/10.1074/jbc.271.46.28784>, 8910521.
- [316] Y Akamatsu, T Ohno, K Hirota, H Kagoshima, Y Yodoi, K Shigesada, Redox regulation of the DNA binding activity in transcription factor PEBP2. The roles of two conserved cysteine residues, *Journal of Biological Chemistry*. 272 (1997) 14497–500. <http://dx.doi.org/10.1074/jbc.272.23.14497>, 9169404.
- [317] C Gaidon, NC Moorthy, C Prives, Ref-1 regulates the transactivation and proapoptotic functions of p53 in vivo, *EMBO Journal*. 18 (1999) 5609–21.

- [318] M Ema, K Hirota, J Mimura, H Abe, J Yodoi, K Sogawa, et al, Molecular mechanisms of transcription activation by HLF and HIF1 alpha in response to hypoxia: their stabilization and redox signal-induced interaction with CBP/p300, *EMBO Journal*, 18 (1999) 1905–14. .
- [319] T Nishi, N Shimizu, M Hiramoto, I Sato, Y Yamaguchi, M Hasegawa, et al, Spatial redox regulation of a critical cysteine residue of NF-kappa B in vivo, *Journal of Biological Chemistry*, 277 (2002) 44548–56. <http://dx.doi.org/10.1074/jbc.M202970200>, 12213807.
- [320] X Cao, F Kambe, S Ohmori, H Seo, Oxidoreductive modification of two cysteine residues in paired domain by ref-1 regulates DNA-binding activity of pax-8, *Biochemical and Biophysical Research Communications*, 297 (2002) 288–93. [http://dx.doi.org/10.1016/S0006-291X\(02\)02196-4](http://dx.doi.org/10.1016/S0006-291X(02)02196-4), 12237116.
- [321] JR Matthews, N Wakasugi, JL Virelizier, J Yodoi, RT Hay, Thioredoxin regulates the DNA binding activity of NF-kappa B by reduction of a disulphide bond involving cysteine 62, *Nucleic Acids Research*, 20 (1992) 3821–30. <http://dx.doi.org/10.1093/nar/20.15.3821>, 1508666.
- [322] T Hayashi, Y Ueno, T Okamoto, Oxidoreductive regulation of nuclear factor kappa B. Involvement of a cellular reducing catalyst thioredoxin, *Journal of Biological Chemistry*, 268 (1993) 11380–8. .
- [323] JM Hansen, WH Watson, DP Jones, Compartmentation of Nrf-2 redox control: regulation of cytoplasmic activation by glutathione and DNA binding by thioredoxin-1, *Toxicological Sciences: An Official Journal of the Society of Toxicology*, 82 (2004) 308–17. <http://dx.doi.org/10.1093/toxsci/kfh231>, 15282410.
- [324] K Hirota, M Matsui, S Iwata, A Nishiyama, K Mori, J Yodoi, AP-1 transcriptional activity is regulated by a direct association between thioredoxin and ref-1, *Proceedings of the National Academy of Sciences of the United States of America*, 94 (1997) 3633–8. <http://dx.doi.org/10.1073/pnas.94.8.3633>, 9108029.
- [325] J Buzek, L Latonen, S Kurki, K Peltonen, M Laiho, Redox state of tumor suppressor p53 regulates its sequence-specific DNA binding in DNA-damaged cells by cysteine 277, *Nucleic Acids Research*, 30 (2002) 2340–8. <http://dx.doi.org/10.1093/nar/30.11.2340>, 12034820.
- [326] C Dai, W Gu, p53 post-translational modification: deregulated in tumorigenesis, *Trends in Molecular Medicine*, 16 (2010) 528–36. <http://dx.doi.org/10.1016/j.molmed.2010.09.002>, 20932800.
- [327] J-P Kruse, W Gu, MSL2 Promotes Mdm2-independent cytoplasmic localization of p53, *Journal of Biological Chemistry*, 284 (2009) 3250–63. .
- [328] AM Bode, Z Dong, Post-translational modification of p53 in tumorigenesis, *Nature Reviews Cancer*, 4 (2004) 793–805. <http://dx.doi.org/10.1038/nrc1455>, 15510160.
- [329] E Appella, CW Anderson, Post-translational modifications and activation of p53 by genotoxic stresses: p53 post-translational modifications, *European Journal of Biochemistry*, 268 (2001) 2764–72. <http://dx.doi.org/10.1046/j.1432-1327.2001.02225.x>, 11358490.
- [330] MS Colman, CA Afshari, JC Barrett, Regulation of p53 stability and activity in response to genotoxic stress, *Mutation Research*, 462 (2000) 179–88. [http://dx.doi.org/10.1016/S1383-5742\(00\)00035-1](http://dx.doi.org/10.1016/S1383-5742(00)00035-1), 10767629.
- [331] Y Xu, Regulation of p53 responses by post-translational modifications, *Cell Death and Differentiation*, 10 (2003) 400–3. <http://dx.doi.org/10.1038/sj.cdd.4401182>, 12719715.
- [332] J-P Kruse, W Gu, Modes of p53 Regulation, *Cell*, 137 (2009) 609–22. <http://dx.doi.org/10.1016/j.cell.2009.04.050>, 19450511.
- [333] K Chen, Activation of p53 by oxidative stress involves platelet-derived growth factor-beta receptor-mediated ataxia telangiectasia mutated (ATM) kinase activation, *Journal of Biological Chemistry*, 278 (2003) 39527–33. <http://dx.doi.org/10.1074/jbc.M304423200>, 12890678.
- [334] S Xie, Q Wang, H Wu, J Cogswell, L Lu, M Jhanwar-Uniyal, et al, Reactive oxygen species-induced phosphorylation of p53 on serine 20 is mediated in part by polo-like kinase-3, *Journal of Biological Chemistry*, 276 (2001) 36194–9. <http://dx.doi.org/10.1074/jbc.M104157200>, 11447225.
- [335] R Dai, W Frejtag, B He, Y Zhang, NF Mivechi, C-Jun NH2-terminal kinase targeting and phosphorylation of heat shock factor-1 suppress its transcriptional activity, *Journal of Biological Chemistry*, 275 (2000) 18210–18. <http://dx.doi.org/10.1074/jbc.M000958200>, 10747973.
- [336] S Chu, TJ Ferro, Identification of a hydrogen peroxide-induced PP1-JNK1-Sp1 signaling pathway for gene regulation, *American Journal of Physiology – Lung Cellular and Molecular Physiology*, 291 (2006) L983–L992. <http://dx.doi.org/10.1152/ajplung.00454.2005>, 16815888.
- [337] S Morton, RJ Davis, A McLaren, P Cohen, A reinvestigation of the multisite phosphorylation of the transcription factor c-Jun, *EMBO Journal*, 22 (2003) 3876–86.
- [338] JW Tullai, S Tacheva, LJ Owens, JR Graham, GM Cooper, AP-1 is a component of the transcriptional network regulated by GSK-3 in quiescent cells, *PLoS One*, 6 (2011) e20150. <http://dx.doi.org/10.1371/journal.pone.0020150>, 21647439.
- [339] E Iwata, M Asanuma, S Nishibayashi, Y Kondo, N Ogawa, Different effects of oxidative stress on activation of transcription factors in primary cultured rat neuronal and glial cells, *Molecular Brain Research*, 50 (1997) 213–20. [http://dx.doi.org/10.1016/S0169-328X\(97\)00190-3](http://dx.doi.org/10.1016/S0169-328X(97)00190-3), 9406937.
- [340] S Pugazhenthai, A Nesterova, P Jambal, K Audestirk, M Kern, L Cabell, et al, Oxidative stress-mediated down-regulation of bcl-2 promoter in hippocampal neurons: down-regulation of CREB by oxidative stress, *Journal of Neurochemistry*, 84 (2003) 982–96. <http://dx.doi.org/10.1046/j.1471-4159.2003.01606.x>, 12603823.
- [341] CL Brooks, W Gu, The impact of acetylation and deacetylation on the p53 pathway, *Protein and Cell*, 2 (2011) 456–62. <http://dx.doi.org/10.1007/s13238-011-1063-9>, 21748595.
- [342] Y Kawai, L Garduño, M Theodore, J Yang, IJ Arinze, Acetylation–deacetylation of the transcription factor Nrf2 (nuclear factor erythroid 2-related factor 2) regulates its transcriptional activity and nucleocytoplasmic localization, *Journal of Biological Chemistry*, 286 (2011) 7629–40. <http://dx.doi.org/10.1074/jbc.M110.208173>, 21196497.
- [343] C Cao, S Lu, R Kivlin, B Wallin, E Card, A Bagdasarian, et al, SIRT1 confers protection against UVB- and H₂O₂-induced cell death via modulation of p53 and JNK in cultured skin keratinocytes, *Journal of Cellular and Molecular Medicine*, 13 (2009) 3632–43. <http://dx.doi.org/10.1111/j.1582-4934.2008.00453.x>, 18681908.
- [344] A Furukawa, S Tada-Oikawa, S Kawanishi, S Oikawa, H₂O₂ accelerates cellular senescence by accumulation of acetylated p53 via decrease in the function of SIRT1 by NAD⁺ depletion, *Cellular Physiology and Biochemistry*, 20 (2007) 45–54. .
- [345] Z Xu, LSM Lam, LH Lam, SF Chau, TB Ng, SWN Au, Molecular basis of the redox regulation of SUMO proteases: a protective mechanism of intermolecular disulfide linkage against irreversible sulphydryl oxidation, *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology*, 22 (2008) 127–37. .
- [346] N Masson, RS Singleton, R Sekirnik, DC Trudgian, LJ Ambrose, MX Miranda, et al, The FIH hydroxylase is a cellular peroxide sensor that modulates HIF transcriptional activity, *EMBO Reports*, 13 (2012) 251–7. .
- [347] X He, Q Ma, NRF2 cysteine residues are critical for oxidant/electrophile-sensing, Kelch-like ECH-associated protein-1-dependent ubiquitination–proteasomal degradation, and transcription activation, *Molecular Pharmacology*, 76 (2009) 1265–78. <http://dx.doi.org/10.1124/mol.109.058453>, 19786557.
- [348] M Cavigelli, F Dolfi, FX Claret, M Karin, Induction of c-fos expression through JNK-mediated TCF/Elk-1 phosphorylation, *EMBO Journal*, 14 (1995) 5957–64. .
- [349] B Dérjard, M Hibi, IH Wu, T Barrett, B Su, T Deng, et al, JNK1: A protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain, *Cell*, 76 (1994) 1025–37. [http://dx.doi.org/10.1016/0092-8674\(94\)90380-8](http://dx.doi.org/10.1016/0092-8674(94)90380-8), 8137421.
- [350] C Weiss, S Schneider, EF Wagner, X Zhang, E Seto, D Bohmann, JNK phosphorylation relieves HDAC3-dependent suppression of the transcriptional activity of c-Jun, *EMBO Journal*, 22 (2003) 3686–95. .
- [351] H Ichijo, E Nishida, K Irie, P ten Dijke, M Saitoh, T Moriguchi, et al, Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways, *Science*, 275 (1997) 90–4.
- [352] PJ Nadeau, SJ Charette, MB Toledano, J Landry, Disulfide bond-mediated multimerization of ASK1 and its reduction by thioredoxin-1 regulate H₂O₂-induced c-Jun NH2-terminal kinase activation and apoptosis, *Molecular Biology of the Cell*, 18 (2007) 3903–13. <http://dx.doi.org/10.1091/mbc.E07-05-0491>, 17652454.
- [353] M Saitoh, H Nishitoh, M Fujii, K Takeda, K Tobiume, Y Sawada, et al, Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1, *EMBO Journal*, 17 (1998) 2596–606. .
- [354] V Adler, Z Yin, SY Fuchs, M Benezra, L Rosario, KD Tew, et al, Regulation of JNK signaling by GSTp, *EMBO Journal*, 18 (1999) 1321–34. .
- [355] K Morita, M Saitoh, K Tobiume, H Matsuura, S Enomoto, H Nishitoh, et al, Negative feedback regulation of ASK1 by protein phosphatase 5 (PP5) in response to oxidative stress, *EMBO Journal*, 20 (2001) 6028–36. .
- [356] H Kamata, S-I Honda, S Maeda, L Chang, H Hirata, M Karin, Reactive oxygen species promote TNFalpha-induced death and sustained JNK activation by inhibiting MAP kinase phosphatases, *Cell*, 120 (2005) 649–61. <http://dx.doi.org/10.1016/j.cell.2004.12.041>, 15766528.
- [357] TD Foley, JJ Armstrong, BR Kupchak, Identification and H₂O₂ sensitivity of the major constitutive MAPK phosphatase from rat brain, *Biochemical and Biophysical Research Communications*, 315 (2004) 568–74. <http://dx.doi.org/10.1016/j.bbrc.2004.01.096>, 14975738.
- [358] Y Wang, PA Gibney, JD West, KA Morano, The yeast Hsp70 Ssa1 is a sensor for activation of the heat shock response by thiol-reactive compounds, *Molecular Biology of the Cell*, 23 (2012) 3290–8. <http://dx.doi.org/10.1091/mbc.E12-06-0447>, 22809627.
- [359] EA Craig, K Jacobsen, Mutations of the heat inducible 70 kilodalton genes of yeast confer temperature sensitive growth, *Cell*, 38 (1984) 841–9. [http://dx.doi.org/10.1016/0092-8674\(84\)90279-4](http://dx.doi.org/10.1016/0092-8674(84)90279-4), 6386178.
- [360] J-Y Chuang, W-C Chang, J-J Hung, Hydrogen peroxide induces Sp1 methylation and thereby suppresses cyclin B1 via recruitment of Suv39H1 and HDAC1 in cancer cells, *Free Radical Biology and Medicine*, 51 (2011) 2309–18. <http://dx.doi.org/10.1016/j.freeradbiomed.2011.10.001>, 22036763.
- [361] CC Winterbourn, D Metodiewa, Reactivity of biologically important thiol compounds with superoxide and hydrogen peroxide, *Free Radical Biology and Medicine*, 27 (1999) 322–8. [http://dx.doi.org/10.1016/S0891-5849\(99\)00051-9](http://dx.doi.org/10.1016/S0891-5849(99)00051-9), 10468205.
- [362] C Little, O'Brien PJ, Mechanism of peroxide-inactivation of the sulphydryl enzyme glyceraldehyde-3-phosphate dehydrogenase, *European Journal of Biochemistry*, 10 (1969) 533–8. <http://dx.doi.org/10.1111/j.1432-1033.1969.tb00721.x>, 5348077.
- [363] R Goldman, DA Stoyanovsky, BW Day, VE Kagan, Reduction of phenoxyl radicals by thioredoxin results in selective oxidation of its SH-groups to disulfides. An antioxidant function of thioredoxin, *Biochemistry (Moscow)*, 34 (1995) 4765–72.
- [364] J Sohn, J Rudolph, Catalytic and chemical competence of regulation of cdc25 phosphatase by oxidation/reduction, *Biochemistry (Moscow)*, 42 (2003) 10060–70.

- [365] AV Peskin, FM Low, LN Paton, GJ Maghzal, MB Hampton, CC Winterbourn, The high reactivity of peroxiredoxin 2 with H₂O₂ is not reflected in its reaction with other oxidants and thiol reagents, *Journal of Biological Chemistry*. 282 (2007) 11885–92. <http://dx.doi.org/10.1074/jbc.M700339200>, 17329258.
- [366] M Trujillo, A Clippe, B Manta, G Ferrer-Sueta, A Smeets, J-P Declercq, et al, Pre-steady state kinetic characterization of human peroxiredoxin 5: taking advantage of Trp84 fluorescence increase upon oxidation, *Archives of Biochemistry and Biophysics*. 467 (2007) 95–106. <http://dx.doi.org/10.1016/j.abb.2007.08.008>, 17892856.
- [367] B Chance, DS Greenstein, FJW Roughton, The mechanism of catalase action. I. Steady-state analysis, *Archives of Biochemistry and Biophysics*. 37 (1952) 301–39. [http://dx.doi.org/10.1016/0003-9861\(52\)90194-X](http://dx.doi.org/10.1016/0003-9861(52)90194-X), 14953443.
- [368] KC Das, R Dashnamoorthy, Hyperoxia activates the ATR-Chk1 pathway and phosphorylates p53 at multiple sites, *American Journal of Physiology: Lung Cellular and Molecular Physiology*. 286 (2004) L87–L97, .
- [369] T Buschmann, O Potapova, A Bar-Shira, VN Ivanov, SY Fuchs, S Henderson, et al, Jun NH2-terminal kinase phosphorylation of p53 on Thr-81 is Important for p53 Stabilization and transcriptional activities in response to stress, *Molecular and Cellular Biology*. 21 (2001) 2743–54. <http://dx.doi.org/10.1128/MCB.21.8.2743-2754.2001>, 11283254.
- [370] Y Ido, A Duranton, F Lan, JM Cacicedo, TC Chen, L Breton, et al, Acute activation of AMP-activated protein kinase prevents H₂O₂-induced premature senescence in primary human keratinocytes, *PLOS One*. 7 (2012) e35092. <http://dx.doi.org/10.1371/journal.pone.0035092>, 22514710.
- [371] YJ Lin, YZ Zhen, J Wei, B Liu, ZY Yu, G Hu, Effects of Rhein lysinate on H₂O₂-induced cellular senescence of human umbilical vascular endothelial cells, *Acta Pharmacologica Sinica*. 32 (2011) 1246–52. <http://dx.doi.org/10.1038/aps.2011.101>, 21909125.
- [372] Hyun Ae Woo, Woojin Jeong, Tong-Shin Chang, Kwang Joo Park, Sung Jun Park, Jeong Soo Yang, et al, Reduction of cysteine sulfinic acid by sulfiredoxin is specific to 2-cys peroxiredoxins. *The Journal of biological chemistry*. 280(5) (2005) 3125–8, .