

# Transcription Factors That Defend Bacteria Against Reactive Oxygen Species

James A. Imlay

Department of Microbiology, University of Illinois, Urbana, Illinois 61801;  
email: jimlay@illinois.edu

Annu. Rev. Microbiol. 2015. 69:93–108

First published online as a Review in Advance on  
June 11, 2015

The *Annual Review of Microbiology* is online at  
micro.annualreviews.org

This article's doi:  
10.1146/annurev-micro-091014-104322

Copyright © 2015 by Annual Reviews.  
All rights reserved

## Keywords

hydrogen peroxide, superoxide, OxyR, PerR, SoxR, reactive oxygen species

## Abstract

Bacteria live in a toxic world in which their competitors excrete hydrogen peroxide or superoxide-generating redox-cycling compounds. They protect themselves by activating regulons controlled by the OxyR, PerR, and SoxR transcription factors. OxyR and PerR sense peroxide when it oxidizes key thiolate or iron moieties, respectively; they then induce overlapping sets of proteins that defend their vulnerable metalloenzymes. An additional role for OxyR in detecting electrophilic compounds is possible. In some nonenteric bacteria, SoxR appears to control the synthesis and export of redox-cycling compounds, whereas in the enteric bacteria it defends the cell against the same agents. When these compounds oxidize its iron-sulfur cluster, SoxR induces proteins that exclude, excrete, or modify them. It also induces enzymes that defend the cell against the superoxide that such compounds make. Recent work has brought new insight into the biochemistry and physiology of these responses, and comparative studies have clarified their evolutionary histories.

## Contents

INTRODUCTION .....	94
DETECTING HYDROGEN PEROXIDE: OxyR AND PerR .....	94
The Threat of H <sub>2</sub> O <sub>2</sub> .....	94
Defensive Tactics: The OxyR Regulon .....	95
How Does OxyR Sense H <sub>2</sub> O <sub>2</sub> ? .....	96
Has OxyR Evolved to Respond to Stresses Other Than H <sub>2</sub> O <sub>2</sub> ? .....	97
PerR of <i>Bacillus subtilis</i> : An Alternative to OxyR .....	98
SUPEROXIDE STRESS: RETHINKING SoxR .....	99
The Threat of Redox-Cycling Compounds .....	99
The Sensing Mechanism of SoxR .....	99
What Is the Inducer That Oxidizes SoxR? .....	100
A More Expansive View: SoxR Beyond <i>E. Coli</i> .....	102
A FINAL WORD ABOUT OTHER ANTIOXIDANT REGULONS .....	102
PROSPECTS .....	103

## INTRODUCTION


The most harmful effects of oxygen are mediated by its partially reduced forms—hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and superoxide (O<sub>2</sub><sup>−</sup>). These species continuously arise inside cells through the autoxidation of redox enzymes, and so aerobes maintain high titers of scavenging enzymes that hold these oxidants below the threshold of toxicity. But by the 1980s it was realized that a variety of natural circumstances can elevate these species above that threshold and that microbes universally maintain inducible defensive systems to counteract such events. Several such systems have now been dissected, and the goal of this review is to map out what is known and what key problems remain. The focus will be on the three transcription factors that have been most intensively examined: OxyR and PerR, which respond to H<sub>2</sub>O<sub>2</sub> stress, and SoxR, which defrays the toxic effects of O<sub>2</sub><sup>−</sup>.

## DETECTING HYDROGEN PEROXIDE: OxyR AND PerR

### The Threat of H<sub>2</sub>O<sub>2</sub>

H<sub>2</sub>O<sub>2</sub> arises in natural habitats through reactions between sulfur and oxygen at oxic/anoxic interfaces, the photochemical reduction of oxygen by chromophores, and the redox cycling of pigments. More ominously, plants, animals, and certain bacteria excrete H<sub>2</sub>O<sub>2</sub> to poison local microbes (5, 27, 79, 98). H<sub>2</sub>O<sub>2</sub> is an ideal weapon. Because it is small and uncharged, it passively crosses membranes at rates similar to that of water, so it cannot be excluded by the targeted cell (97, 115). Once inside it disrupts multiple aspects of iron metabolism, thereby attacking a feature of life that is virtually universal.

The ways in which H<sub>2</sub>O<sub>2</sub> damages cells have been identified in the model bacterium *Escherichia coli*. Exogenous H<sub>2</sub>O<sub>2</sub> damages DNA, so it is mutagenic. This effect arises from Fenton chemistry, in which H<sub>2</sub>O<sub>2</sub> reacts with the intracellular pool of unincorporated iron (42):




The hydroxyl radical ( $\text{HO}\cdot$ ) reacts at diffusion-limited rates with most biomolecules. Some of the iron pool is loosely associated with DNA (94), and Fenton reactions on its surface produce the lesions that cause mutagenesis.

However, the most marked effect of  $\text{H}_2\text{O}_2$  stress is to inactivate two families of iron-containing enzymes: nonredox mononuclear enzymes and [4Fe-4S]-containing dehydratases (2, 46, 103, 104). Both classes of enzymes employ a solvent-exposed reduced iron atom to bind and activate substrates, and in both cases incoming  $\text{H}_2\text{O}_2$  can contact and oxidize the iron. The metal dissociates, and activity is lost. Such enzymes sit within the TCA cycle, the pentose phosphate pathway, and key biosynthetic pathways, so growth stops. As little as  $0.5\ \mu\text{M}$   $\text{H}_2\text{O}_2$  is sufficient to poison cells in this way, so the titers of scavenging enzymes are calibrated to keep the steady-state  $\text{H}_2\text{O}_2$  concentration below this level. This goal becomes much harder when bacteria enter a habitat containing extracellular  $\text{H}_2\text{O}_2$ . The  $\text{H}_2\text{O}_2$  quickly flows across membranes and into the cytoplasm. For that circumstance virtually all bacteria maintain inducible defensive regulons governed by either the OxyR or the PerR transcription factors.

## Defensive Tactics: The OxyR Regulon

The OxyR protein is more widely distributed. It features a sensory cysteine residue that is directly oxidized by  $\text{H}_2\text{O}_2$ , thereby generating a disulfide bond that alters its binding to DNA (**Figure 1a**) (4, 11). In some bacteria OxyR acts as a repressor, and its oxidation inactivates this function and stimulates gene expression (39, 40, 72, 73, 105). More commonly, OxyR acts as an activator protein that recruits RNA polymerase. In *E. coli* the oxidized protein stimulates the transcription of about two dozen genes that are well suited to address the problems that  $\text{H}_2\text{O}_2$  causes (**Table 1**) (120). To drive down the  $\text{H}_2\text{O}_2$  level, OxyR strongly induces both an NADH peroxidase (AhpCF) and a catalase (KatG) (44). A mini-ferritin, Dps, is synthesized to sequester unincorporated iron (10, 31). This has the effect of substantially suppressing the rate of DNA damage, especially in collaboration with the induction of Fur and Yaa proteins (71, 111, 119). The MntH manganese importer is induced, apparently to enable Mn—which does not react with  $\text{H}_2\text{O}_2$ —to displace Fe as the catalytic cofactor of mononuclear enzymes (3, 50, 103). Induction of the Suf iron-sulfur assembly system enables the repair of damaged iron-sulfur clusters. Although the cell normally uses an Isc system for this purpose, Isc does not work well when iron levels decline, a situation that ensues from the



**Figure 1**

Activation of OxyR. (a) Oxidation of a sensory cysteine by  $\text{H}_2\text{O}_2$  perturbs global conformation. In *E. coli* the disulfide-bonded form of the protein stabilizes the transcription complex. (b) The acute  $\text{H}_2\text{O}_2$  sensitivity of OxyR may depend on a shift in hydrogen bonds from the sensory cysteine thiolate to the incoming  $\text{H}_2\text{O}_2$ . This idea derives from models of peroxiredoxin behavior (18, 106).

**Table 1** The *Escherichia coli* OxyR regulon<sup>a</sup>

Gene	Activity	Impact
<i>katG</i>	Catalase	Scavenge H <sub>2</sub> O <sub>2</sub>
<i>ahpCF</i>	NADH peroxidase	
<i>dps</i>	Mini-ferritin	Minimize free iron
<i>fur</i>	Iron-import repressor	
<i>yaaA</i>	Unknown biochemistry	
<i>mntH</i>	Manganese importer	Activates mononuclear enzymes
<i>sufA-E</i>	Iron-sulfur assembly	Activates Fe/S enzymes
<i>hemF</i>	Coproporphyrinogen III oxidase	Sustain heme synthesis
<i>hemH</i>	Ferrochelatase	
<i>gor</i>	Glutathione reductase	Maintain thiol
<i>trxC</i>	Thioredoxin	
<i>grxA</i>	Glutaredoxin	
<i>dsbG</i>	Protein sulfenate reductase	

<sup>a</sup>Several genes of unknown function are not included. The PerR regulon of *Bacillus subtilis* includes catalase, mini-ferritin (MrgA), Fur, and heme biosynthesis genes (7). In that bacterium, Mn importers and Suf proteins are constitutively synthesized, whereas disulfide-reducing systems are controlled by an independent regulator, Spx, that does not respond to H<sub>2</sub>O<sub>2</sub>.

action of Dps (47, 85). Similarly, the iron-dependent enzymes in the heme biosynthetic pathway are replaced (HemF) or induced (HemH) to sustain this process as intracellular iron becomes scarce (77).

Three regulon members typically reduce disulfide bonds: glutathione reductase, glutaredoxin 1, and thioredoxin 2. The glutaredoxin assists in deactivating OxyR once H<sub>2</sub>O<sub>2</sub> has dissipated (4), but the roles of the others are less obvious. Deletion of these genes does not cause obvious sensitivity to H<sub>2</sub>O<sub>2</sub>, which inspires other ideas about their purpose in this regulon (see below).

## How Does OxyR Sense H<sub>2</sub>O<sub>2</sub>?

Recent studies have analyzed both the specificity and the sensitivity of OxyR. *E. coli* OxyR features a hyperreactive thiol (Cys199) that is quickly oxidized by H<sub>2</sub>O<sub>2</sub> to a sulfenic acid (–SOH). The adducted residue moves from the hydrophobic cleft in which it is normally buried, and it swings into the proximity of Cys208, with which it condenses to form a disulfide bond (11). This shift locks the domain into a conformation that activates the protein as a transcription factor. The protein functions as a dimer, and it displays some cooperativity that may avoid activation when H<sub>2</sub>O<sub>2</sub> is scant (61). The fact that OxyR uses cysteine oxidation to detect H<sub>2</sub>O<sub>2</sub> enables it to work like a rheostat, given that disulfide formation is among the few amino acid oxidations that are reversible.

The surprising aspect of this chemistry is that cysteine per se reacts very sluggishly with H<sub>2</sub>O<sub>2</sub> (2 M<sup>–1</sup> s<sup>–1</sup>) (116). With such a rate constant, the half-time for oxidation by 0.5 μM H<sub>2</sub>O<sub>2</sub>—an intracellular concentration that suffices to block *E. coli* growth (103)—is approximately one week. In sharp contrast, the sensing cysteine residue on OxyR exhibits a rate constant of 10<sup>5</sup> M<sup>–1</sup> s<sup>–1</sup> and detects micromolar H<sub>2</sub>O<sub>2</sub> within seconds (4). What is the source of this kinetic improvement?

The hyperreactive behavior of OxyR C199 has not been deconstructed in detail, but it resembles that of the catalytic cysteine residues on peroxiredoxins, the ubiquitous peroxidases (including *E. coli* AhpC) that degrade H<sub>2</sub>O<sub>2</sub> through cycles of cysteine oxidation and reduction

(18, 106). As with OxyR, their reaction involves a nucleophilic attack of a cysteine thiolate on  $\text{H}_2\text{O}_2$  (**Figure 1a**), with cleavage of the dioxygen bond. A nearby cationic residue sets the stage by ensuring deprotonation of the cysteine (83)—but this should provide only an order-of-magnitude improvement in reactivity. A plausible explanation for the remaining stimulation is depicted in **Figure 1b**. The peroxidatic cysteine residue of peroxiredoxins participates in hydrogen-bond networks that shield its charge within a largely hydrophobic cleft (91). Entry of  $\text{H}_2\text{O}_2$  may shift these bonds toward the  $\text{H}_2\text{O}_2$  itself, freeing the thiolate. Several aspects of this are conducive to catalysis. Release of the cysteine residue generates a nucleophile whose potency is enhanced in a low-dielectric environment. The hydrogen bonds that shift toward  $\text{H}_2\text{O}_2$  likely polarize its dioxygen bond, making it more conducive to attack, and one of these residues probably protonates the hydroxide leaving group to pull the reaction forward. The cysteine residue of OxyR is invariably flanked by arginine and histidine residues that may play the same role.


A hydrogen-bond network configured to fit  $\text{H}_2\text{O}_2$  should also enhance the rate constant by disfavoring water in the active site, given that  $\text{H}_2\text{O}$  is too small a molecule to bridge a hydrogen-bond network tailored for  $\text{H}_2\text{O}_2$  (93). The upshot is that thiol-based peroxidases achieve a level of reactivity ( $10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) that is appropriate for their physiological roles, and similar features may allow OxyR to do so as well. A new structure of the *Pseudomonas aeruginosa* OxyR protein suggests that  $\text{H}_2\text{O}_2$  binding might engage a hydrogen-bond network that ultimately links the deprotonation of cysteine with the protonation of the product hydroxide group (48).

In *E. coli* OxyR quickly responds to the submicromolar levels of  $\text{H}_2\text{O}_2$  that threaten to disrupt metabolism. The facultative bacterium *Vibrio vulnificus* takes it a step further, employing two OxyR proteins that are calibrated to sense distinct levels of  $\text{H}_2\text{O}_2$ : The more sensitive OxyR is activated by the low levels of  $\text{H}_2\text{O}_2$  that are endogenously generated upon aeration, whereas the less sensitive one responds to more-severe influxes of  $\text{H}_2\text{O}_2$  from the external environment (53).

## Has OxyR Evolved to Respond to Stresses Other Than $\text{H}_2\text{O}_2$ ?

Because OxyR responds to adduction of its sensory thiol, it can also be activated by electrophiles (118). Experimenters have commonly used the synthetic chemical diamide as a proxy for such compounds. An interesting question is whether OxyR has an authentic role in sensing stressors other than  $\text{H}_2\text{O}_2$ . Seth et al. (99) reported that OxyR becomes nitrosylated during anaerobic nitrate reduction, presumably by nitric oxide that escapes from the system. Indeed, an *oxyR* mutant exhibited poorer growth in this circumstance, suggesting that some elements of the regulon help the cell to tolerate this stress.

Many bacteria contain transcription factors other than OxyR that appear to be dedicated to responding to electrophiles, whose attack on thiols can indirectly generate disulfide bonds. Both the Spx system of *Bacillus subtilis* and the SigR/RsrA system of *Streptomyces coelicolor* induce a mixture of glutaredoxins, thioredoxins, chaperones, and proteases, which collectively avert protein aggregation from the denaturing effects of disulfide formation and thiol adduction (49, 52, 86, 96, 121). These bacteria additionally contain OxyR or PerR systems that respond to  $\text{H}_2\text{O}_2$ —but in these bacteria, the latter systems do not include disulfide-reducing genes that are part of the *E. coli* OxyR regulon. Thus, it seems possible that *E. coli* is exceptional in multitasking OxyR with responding to both stresses:  $\text{H}_2\text{O}_2$ , which primarily threatens iron enzymes, and electrophiles, which react with cysteine residues much more avidly than does  $\text{H}_2\text{O}_2$ .



**Figure 2**

Structure of PerR in its (a) holoenzyme and (b) demetallated forms. Structures were derived from the Protein Data Bank (3F8N and 2FE3; <http://www.rcsb.org/pdb>) (45, 107). Orange spheres represent Zn, and purple spheres represent Mn.

### PerR of *Bacillus subtilis*: An Alternative to OxyR

Gram-positive bacteria often employ a different sensor of  $\text{H}_2\text{O}_2$ : PerR. This dimeric protein belongs to the Fur family of metal-binding transcription factors, which have evolved in their various forms to sense iron (Fur proper), manganese (Mur), zinc (Zur), nickel (Nur), or  $\text{H}_2\text{O}_2$  (PerR) (19, 38). Each subunit contains a structural site that irreversibly binds zinc, plus at least one regulatory site that reversibly binds the activating metal (**Figure 2a**). Only the metal-bound forms of the proteins are competent to bind DNA, and they generally work as repressors. In *B. subtilis* PerR, the regulatory metal is ferrous iron, and it is affixed in a distorted square pyramid with four coplanar ligands and a single axial ligand (45). When  $\text{H}_2\text{O}_2$  levels rise,  $\text{H}_2\text{O}_2$  accesses the sixth coordination site and oxidizes the iron atom. The ferryl or hydroxyl radical that is thereby formed oxidizes the histidine ligands of the iron atom to 2-oxohistidine (64). The modified residues can no longer bind the metal, and the resultant apoprotein undergoes a global conformational change that eradicates its ability to bind DNA (**Figure 2b**). Because holo-PerR acts as a repressor, the effect is to derepress regulon members whenever  $\text{H}_2\text{O}_2$  levels rise.

Strikingly, *B. subtilis* PerR reacts with  $\text{H}_2\text{O}_2$  at essentially the same rate constant as does *E. coli* OxyR ( $10^5 \text{ M}^{-1} \text{ s}^{-1}$ ) (63). Other mononuclear iron proteins, such as *E. coli* peptide deformylase, have similar inactivation rate constants (2), indicating that this value is largely dictated by the fundamental rate of Fenton reactions. However, *B. subtilis* also employs a true Fur protein whose iron cofactor seems to be resistant to oxidation by  $\text{H}_2\text{O}_2$ . Inspection of this discrepancy elicited evidence that in the Fur protein a coordinating glutamate residue probably acts as a bidentate ligand to the iron atom, thereby occluding its sixth coordination site (89). The analogous aspartate ligand of PerR does not. Given that  $\text{H}_2\text{O}_2$  must directly contact iron in order to oxidize it (28),  $\text{H}_2\text{O}_2$  oxidizes PerR but not Fur.

The PerR regulon of *B. subtilis* includes many of the same genes that OxyR controls in other bacteria: genes encoding a catalase, a Dps homolog (MrgA), and Fur (7) (**Table 1**). Yet PerR offers several intriguing departures from the OxyR mechanism of  $\text{H}_2\text{O}_2$  sensing. First, given that there is no known mechanism of oxo-histidine reduction, the reaction is presumably irreversible, perhaps making the system more of an on/off switch. This arrangement is a rarity: With few exceptions (80), bacterial regulators are reversible, and their activities are set by the dynamic equilibrium between activating and deactivating events. It is not clear whether the irreversibility of PerR deactivation provides any advantage.

Second, like other mononuclear proteins, PerR can bind manganese in place of iron, both in vitro and in vivo (9). Given that Mn is not oxidized by  $\text{H}_2\text{O}_2$ , the upshot is that PerR does not respond to  $\text{H}_2\text{O}_2$  whenever the cytoplasm is rich in manganese. As manganese helps to defend



cells against  $\text{H}_2\text{O}_2$ , this arrangement suggests that a primary goal of the PerR and OxyR responses may be to raise the Mn/Fe ratio (117); if that ratio is already high, no induction is needed.

Finally, PerR does not control the synthesis of disulfide-reducing redoxins, as their synthesis is governed by the Spx response regulator (121), which is not responsive to  $\text{H}_2\text{O}_2$ . PerR appears to be focused exclusively on the detection and prevention of Fenton chemistry.

## SUPEROXIDE STRESS: RETHINKING SoxR


### The Threat of Redox-Cycling Compounds

Because  $\text{O}_2^-$  is a charged molecule, it cannot penetrate bacteria from outside (57, 74); therefore, any cytoplasmic  $\text{O}_2^-$  stress is due to  $\text{O}_2^-$  that is generated internally. Endogenous superoxide ( $\text{O}_2^-$ ) that is formed by enzyme autoxidation is kept very scarce ( $\sim 10^{-10}$  M) by the action of superoxide dismutases (SODs) (41). *E. coli* mutants that lack cytoplasmic SODs exhibit growth defects (8) that have been traced back to deficiencies in the same enzymes that  $\text{H}_2\text{O}_2$  inactivates: [4Fe-4S] dehydratases and mononuclear iron enzymes (20, 24, 35, 60, 103). These mutants are unable to synthesize branched-chain or aromatic amino acids, and they cannot catabolize TCA-cycle substrates.

The same phenotypes arise even in wild-type *E. coli* if it encounters viologens (e.g., paraquat), quinones, or phenazines (37). These cyclic organic chemicals can penetrate cells, where they directly abstract electrons from redox enzymes and transfer them to oxygen. Toxic doses of  $\text{O}_2^-$  are thereby produced. Indeed, quinones and phenazines are released by various bacteria and plants, often to poison microflora (43, 87, 109). In 1977 it was discovered that *E. coli* responds to this stress by inducing higher titers of SOD (36). This adaptation is mediated by two proteins dubbed SoxR and SoxS (superoxide) (32, 108). SoxR is a [2Fe-2S]-containing transcription factor that senses the stress; it then induces transcription of *soxS*. SoxS is a second transcription factor that then activates scores of defensive genes scattered around the chromosome (6). Several of these encode proteins that help suppress the toxicity of  $\text{O}_2^-$ : superoxide dismutase; aconitase A and fumarase C, which are superoxide-resistant isozymes of vulnerable [4Fe-4S] enzymes (68, 110); and YggX, which facilitates cluster-repair processes (30, 34, 90). Like OxyR, the SoxRS system is not activated during normal aerobic growth and does not control the basal level of defensive systems; instead, it becomes active only when forcing conditions emerge.

### The Sensing Mechanism of SoxR

SoxR is a homodimer that binds one [2Fe-2S]<sup>+</sup> cluster per monomer (Figure 3); a second domain includes a DNA-binding HTH motif (114). Electron paramagnetic resonance studies demonstrate that when cells are exposed to redox-cycling compounds, the clusters are quickly oxidized to the [2Fe-2S]<sup>2+</sup> state (15, 26). Upon removal of the oxidant the clusters revert within minutes to their reduced forms. The *E. coli soxS* gene features unusually long 20-bp spacing between its -10 and -35 promoter motifs, and the current model is that cluster oxidation contorts the SoxR dimer and thereby alters the conformation of the bound DNA, moving the promoter motifs to the same side of the helix so that RNA polymerase can productively bind. Physical studies have confirmed that changes in SoxR redox state distort bound DNA (23). Genetic studies indicate that the SoxR reduction to the deactivated form is catalyzed by Rsx/Rse proteins (56) that likely employ NADPH as a source of electrons (55). Thus these data collectively suggest that transcription of *soxS* depends on the dynamic equilibrium between oxidation and reduction of the SoxR iron-sulfur clusters.



**Figure 3**

SoxR in association with DNA. (a) Oxidation of the cluster (arrow) allows the protein to twist the DNA, improving the  $-10$  to  $-35$  spacing. (b) The cluster is exposed on the protein surface so that it is accessible to diverse oxidants. The structure was derived from the Protein Data Bank (2ZHG; <http://www.rcsb.org/pdb>) (114).

When SoxR returns to its reduced form, the transcription of the full regulon quickly ceases owing to the rapid proteolysis of extant SoxS protein (33).


### What Is the Inducer That Oxidizes SoxR?

The apparent parallel with  $\text{H}_2\text{O}_2$  detection by OxyR suggested the obvious possibility that SoxR is activated when  $\text{O}_2^-$  directly oxidizes its clusters. The clusters are located near the protein surface and appear accessible to a small molecule like  $\text{O}_2^-$  (Figure 3) (114). Unfortunately, two efforts to directly detect cluster oxidation by  $\text{O}_2^-$  in vitro led to contradictory results (21, 34), suggesting that one of these complicated experiments was compromised in some way. The in vivo data, however, are clearer and somewhat surprising. In  $\text{SOD}^-$  mutants—which suffer debilitating levels of  $\text{O}_2^-$ —the *soxS* gene is induced only 2- to 4-fold (25, 34). This activation is consistent with the idea that  $\text{O}_2^-$  can oxidize the clusters, but it is far less than the  $\sim 40$ -fold induction that occurs when wild-type cells are treated with redox-cycling compounds. Similarly modest effects were observed for members of the regulon (34, 67, 69). This induction is too minor to be effectual: The SoxRS regulon provided no protection to  $\text{SOD}^-$  mutants unless *soxS* was forcibly induced from a heterologous promoter (34). Thus, high  $\text{O}_2^-$  concentrations are not sufficient to adequately activate SoxR.

Further, overproduction of SOD does not diminish the ability of redox agents to activate SoxR in wild-type cells (25, 29, 34, 68, 76), leading workers to conclude that these drugs oxidize SoxR independently of  $\text{O}_2^-$ . Indeed, exposure to redox compounds can activate SoxR even in anoxic cells, from which  $\text{O}_2^-$  is absolutely absent (14, 34, 58).

An alternative explanation for SoxR oxidation by redox agents is that they deplete NADPH pools and thereby disrupt the ability of Rse/Rsx to keep SoxR in its reduced form. Indeed, some genes within the *E. coli* SoxR regulon provide mechanisms to restore NADPH pools (59, 81). However, measurements indicate that the shift in NADPH redox status [to  $-0.314$  V (58)] is not enough to explain the near-total oxidation of SoxR [ $E_o' = -0.293$  V (54)], and mutations that impair cellular NADPH formation do not activate SoxR (34). Thus, although NADPH depletion might augment the inducing effects of redox agents, by itself it is probably not sufficient to turn on the system.





**Figure 4**

Toxic actions of a redox-cycling agent and the defensive tactics that minimize its accumulation. (a) Redox-cycling compounds such as plumbagin (PB) catalyze electron transfer from flavoproteins to oxygen, directly damage [4Fe-4S]-dependent dehydratases, and activate SoxR by oxidation. Components of the SoxR system exclude the agent by ① altering the lipopolysaccharide coat (via *waaYZ*) (62), ② inhibiting porin synthesis (*micF*) (12), ③ exporting the compound (*tolC*, *acrAB*) (75), and ④ modifying it (*nfsA*, *ygfZ*) (66, 70, 95). (b) Relationship of *E. coli* SoxRS to nonenteric SoxR and *E. coli* MarA. In nonenteric bacteria SoxR detects endogenous redox-active compounds like pyocyanin and induces exporters that excrete it. The SoxRS regulon could plausibly have been created by lateral transfer of *soxR* and duplication of *marA*, as SoxS and MarA exhibit 50% identity. The SoxRS regulon consists of many genes plus several genes that specifically defend against oxidizing compounds. Only representative genes are shown; for a full list, see EcoCyc (<http://ecocyc.org>).

Therefore, it seems likely that the redox compounds themselves directly oxidize the clusters of SoxR, just as they do the metal and flavin moieties of typical redox enzymes. This reaction has been demonstrated *in vitro*, with rate constants that accommodate the rates of SoxR oxidation that are observed *in vivo* (34). The position of the iron atom near the SoxR protein surface ensures that it can be oxidized by diverse univalent oxidants without regard for their structures.

The constituency of the regulon supports the idea that the SoxRS response specifically arose to defend bacteria against redox-cycling compounds. A major element of the SoxRS response in *E. coli* is the induction of proteins that collectively block the entry and accumulation of redox-active compounds within the cell: lipopolysaccharide-modifying enzymes that diminish the ability of such compounds to penetrate the envelope, broad-specificity export systems that actively pump them out of the cytoplasm, and derivatizing enzymes that deactivate them through covalent modification (1, 62, 66, 70, 75, 95) (Figure 4a). Interestingly, SoxR is an important defense against redox-active agents even under anoxic conditions, indicating that these agents can exert O<sub>2</sub><sup>-</sup>-independent mechanisms of toxicity. Their primary action under these conditions is unclear, but such compounds can form adducts to proteins by acting as Michael acceptors, they can deactivate [4Fe-4S] dehydratases by directly oxidizing the clusters, and they can deplete cellular NADPH pools by oxidizing NADPH-reducible enzymes and then dumping the electrons into the anaerobic respiratory chain (34).

Finally, it is worth noting that although H<sub>2</sub>O<sub>2</sub> is an ineffective activator of SoxR (34, 120), nitric oxide (NO) can activate it by forming dinitrosyl iron complexes with the atoms of the iron-sulfur cluster (22, 84). The significance of this effect is uncertain, because dedicated NO scavenging systems are controlled by independent regulatory proteins (NsrR and NorR) that are much more responsive to NO. In one study (92), NO doses that activated the aerobic and anaerobic NO

scavenging systems by 30- and 180-fold only induced *soxS* 5-fold. Furthermore, this slight SoxS induction had no apparent effect on SoxS-controlled genes, leading the authors to suggest that this degree of SoxS induction was inadequate for activation of the full regulon. That outcome dovetails with the failure of 3-fold SoxS induction to provide any protection to SOD<sup>-</sup> cells (34).

### A More Expansive View: SoxR Beyond *E. coli*

Studies of SoxR and SoxS in other bacteria have prompted a substantial reappraisal of their biological role. Dietrich et al. (13) found homologs of SoxR among  $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ -proteobacteria and among actinobacteria, but not elsewhere. Strikingly, only enterics employ SoxS; in all the other bacteria, SoxR directly controls expression of all regulon members. Just as striking is the fact that in these nonenterics the SoxR regulon is quite different from that of *E. coli*: It typically comprises only a handful of genes of unclear function (82), and these generally do not include *sodA* or other genes that might confer resistance to O<sub>2</sub><sup>-</sup> (13, 14, 51, 82, 101). The implication is that the physiological purpose of SoxR in nonenterics is quite different from its purpose in *E. coli*.

Evidence regarding this purpose has come from studies of *P. aeruginosa* and *Streptomyces coelicolor*. Both bacteria synthesize and excrete redox-cycling compounds: pyocyanin for *P. aeruginosa* and actinorhodin for *S. coelicolor*. Different roles have been suggested for these compounds, including toxifying competitors, signaling, assisting electron transfer to distant respiratory substrates, and solubilizing iron (14, 112, 113). The pumps that excrete these compounds, and perhaps the enzymes that complete their synthesis, are induced in both bacteria when SoxR is oxidized by either the redox agent itself or a precursor to it (13, 101) (**Figure 4b**). The SoxR protein can also respond to exogenous synthetic redox compounds such as paraquat, but this is presumably adventitious, as SoxR activation does not enhance resistance to such compounds (88). The implication is that these bacteria use SoxR to control the export of endogenous redox agents into the environment.

Whether or not it is to the advantage of the producing organism, these agents can penetrate and toxify other microbes. It is possible that the latter laterally acquired SoxR as a sensor that could perceive these substances and induce defenses. In *E. coli* the arriving *soxR* gene may have piggybacked onto a preexisting system that responds to other toxic chemicals (**Figure 4b**). SoxS is a paralog of the MarA and Rob transcription factors, which respond to and provide protection against exogenous nonredox agents (salicylate and bile salts, for example). The three proteins are similar enough (~50% identity) that they recognize common regulatory sites, so their regulons overlap extensively (78). The simple model, then, is that *E. coli* acquired SoxR and, after duplication of MarA or Rob, fashioned SoxS to connect SoxR to a preexisting regulon that serves to exclude, modify, or export hazardous compounds. SoxR provided the means to activate this system in response to redox-active compounds. Notably, the reduction potential of *E. coli* SoxR is lower than those of *P. aeruginosa* and *S. coelicolor* SoxR proteins, and accordingly it responds to a larger range of redox agents (100, 102). Extant antioxidant defenses, such as *sodA* and *fumC*, may then have been added to the regulon. The latter genes are significantly more responsive to SoxS than to MarA (78).

### A FINAL WORD ABOUT OTHER ANTIOXIDANT REGULONS

Workers have identified distinct bacterial transcription factors that are activated by singlet oxygen [e.g., RpoE/ChrR (65)], organic hydroperoxides [OhrR (17)], disulfide stress [Spx (121) and SigR/RsrA (52)], or hypochlorous acid [HypT (16)]. Unfortunately, space restrictions preclude a discussion of these systems here. Implicit in this diversity is the fact that “oxidative stress” is an

overbroad term that subsumes a panoply of oxidants that (a) vary in their chemical properties, (b) attack very different targets, and (c) are sensed and defended against in completely different ways. For example, none of those systems is activated by physiological doses of  $\text{H}_2\text{O}_2$  or  $\text{O}_2^-$ , and they provide very little crossover resistance against them.

## PROSPECTS

Because life evolved in an anoxic world, the iron-intensive biochemistry that was inherited by contemporary organisms exhibits some incompatibility with the presence of oxygen. This vulnerability is problematic for bacteria that experience fluctuations in local oxygen tension, and it is exploited by competitors that would stymie their growth. A key goal of current work on these regulons is to identify the natural circumstances in which they are activated. As this review emphasizes, important clues can be deduced from the biochemistry of the sensor proteins themselves and from the membership of the regulons that they control. The emergence of studies in nonenteric bacteria promises to provide clarity by offering new perspectives on these familiar regulators.

## DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

Work in the author's lab is supported by grants GM49640 and GM101012 from the National Institutes of Health.

## LITERATURE CITED

1. Aiba H, Matsuyama S, Mizuno T, Mizushima S. 1987. Function of *micF* as an antisense RNA in osmoregulatory expression of the *ompF* gene in *Escherichia coli*. *J. Bacteriol.* 169:3007–12
2. Anjem A, Imlay JA. 2012. Mononuclear iron enzymes are primary targets of hydrogen peroxide stress. *J. Biol. Chem.* 287:15544–56
3. Anjem A, Varghese S, Imlay JA. 2009. Manganese import is a key element of the OxyR response to hydrogen peroxide in *Escherichia coli*. *Mol. Microbiol.* 72:844–58
4. Aslund F, Zheng M, Beckwith J, Storz G. 1999. Regulation of the OxyR transcription factor by hydrogen peroxide and the cellular thiol-disulfide status. *PNAS* 96:6161–65
5. Bedard K, Lardy B, Krause K-H. 2007. NOX family NADPH oxidases: not just in mammals. *Biochemie* 89:1107–12
6. Blanchard JL, Wholely WY, Conlon EM, Pomposiello PJ. 2007. Rapid changes in gene expression dynamics in response to superoxide reveal SoxRS-dependent and -independent transcriptional networks. *PLOS ONE* 2:e1186
7. Bsat N, Herbig A, Casillas-Martinez L, Setlow P, Helmann JD. 1998. *Bacillus subtilis* contains multiple Fur homologues: identification of the iron uptake (Fur) and peroxide regulon (PerR) repressors. *Mol. Microbiol.* 29:189–98
8. Carlouz A, Touati D. 1986. Isolation of superoxide dismutase mutants in *Escherichia coli*: Is superoxide dismutase necessary for aerobic life? *EMBO J.* 5:623–30
9. Chen L, Keramati L, Helmann JD. 1995. Coordinate regulation of *Bacillus subtilis* peroxide stress genes by hydrogen peroxide and metal ions. *PNAS* 92:8190–94
10. Chiancone E, Ceci P. 2010. The multifaceted capacity of Dps proteins to combat bacterial stress conditions: detoxification of iron and hydrogen peroxide and DNA binding. *Biochim. Biophys. Acta* 1800:798–805

11. Choi H, Kim S, Mukhopadhyay P, Cho S, Woo J, et al. 2001. Structural basis of the redox switch in the OxyR transcription factor. *Cell* 105:103–13
12. Chou JH, Greenberg JT, Demple B. 1993. Posttranscriptional repression of *Escherichia coli* ompF protein in response to redox stress. Positive control of the micF antisense RNA by the soxRS locus. *J. Bacteriol.* 175:1026–31
13. Dietrich LE, Teal TK, Price-Whelan A, Newman DK. 2008. Redox-active antibiotics control gene expression and community behavior in divergent bacteria. *Science* 321:1203–6
14. Dietrich LEP, Price-Whelan A, Petersen A, Whiteley M, Newman DK. 2006. The phenazine pyocyanin is a terminal signalling factor in the quorum sensing network of *Pseudomonas aeruginosa*. *Mol. Microbiol.* 61:1308–21
15. Ding H, Demple B. 1997. In vivo kinetics of a redox-regulated transcriptional switch. *PNAS* 94:8445–49
16. Drazic A, Miura H, Peschek J, Le Y, Bach NC, et al. 2013. Methionine oxidation activates a transcription factor in response to oxidative stress. *PNAS* 110:9493–98
17. Dubbs JM, Mongkolsuk S. 2012. Peroxide-sensing transcriptional regulators in bacteria. *J. Bacteriol.* 194:5495–503
18. Ferrer-Sueta G, Manta B, Botti H, Radi R, Trujillo M, Denicola A. 2011. Factors affecting protein thiol reactivity and specificity in peroxide reduction. *Chem. Res. Toxicol.* 24:434–50
19. Fillat MF. 2014. The FUR (ferric uptake regulator) superfamily: diversity and versatility of key transcriptional regulators. *Arch. Biochem. Biophys.* 546:41–52
20. Flint DH, Tuminello JF, Emptage MH. 1993. The inactivation of Fe-S cluster containing hydro-lyases by superoxide. *J. Biol. Chem.* 268:22369–76
21. Fujikawa M, Kobayashi K, Kozawa T. 2012. Direct oxidation of the [2Fe-2S] cluster in SoxR protein by superoxide: distinct differential sensitivity to superoxide-mediated signal transduction. *J. Biol. Chem.* 287:35702–8
22. Fujikawa M, Kobayashi K, Kozawa T. 2014. Mechanistic studies on formation of the dinitrosyl iron complex of the [2Fe-2S] cluster of SoxR protein. *J. Biochem.* 156:163–72
23. Fujikawa M, Kobayashi K, Kozawa T. 2015. Redox-dependent DNA distortion in a SoxR protein-promoter complex studied using fluorescent probes. *J. Biochem.* 157:389–97
24. Gardner PR, Fridovich I. 1991. Superoxide sensitivity of the *Escherichia coli* aconitase. *J. Biol. Chem.* 266:19328–33
25. Gaudu P, Dubrac S, Touati D. 2000. Activation of SoxR by overproduction of desulfoferrodoxin: multiple ways to induce the soxRS regulon. *J. Bacteriol.* 182:1761–63
26. Gaudu P, Moon N, Weiss B. 1997. Regulation of the soxRS oxidative stress regulon. *J. Biol. Chem.* 272:5082–86
27. Glass GA, DeLisle DM, DeTogni P, Gabig TG, Magee BH, et al. 1986. The respiratory burst oxidase of human neutrophils: further studies of the purified enzyme. *J. Biol. Chem.* 261:13247–51
28. Goldstein S, Meyerstein D, Czapski G. 1993. The Fenton reagents. *Free Rad. Biol. Med.* 15:435–45
29. Gort AS, Imlay JA. 1998. Balance between endogenous superoxide stress and antioxidant defenses. *J. Bacteriol.* 180:1402–10
30. Gralnick JA, Downs DM. 2003. The YggX protein of *Salmonella enterica* is involved in Fe(II) trafficking and minimizes the DNA damage caused by hydroxyl radicals: Residue CYS-7 is essential for YggX function. *J. Biol. Chem.* 278:20708–15
31. Grant RA, Filman DJ, Finkel SE, Kolter R, Hogle JM. 1998. The crystal structure of Dps, a ferritin homolog that binds and protects DNA. *Nat. Struct. Biol.* 5:294–303
32. Greenberg JT, Monach P, Chou JH, Joseph PD, Demple B. 1990. Positive control of a global antioxidant defense regulon activated by superoxide-generating agents in *Escherichia coli*. *PNAS* 87:6181–85
33. Griffith KL, Shah IM, Wolf RE Jr. 2004. Proteolytic degradation of *Escherichia coli* transcription activators Sox and MarA as the mechanism for reversing the induction of the superoxide (SoxRS) and multiple antibiotic resistance (Mar) regulons. *Mol. Microbiol.* 51:1801–16
34. Gu M, Imlay JA. 2011. The SoxRS response of *Escherichia coli* is directly activated by redox-cycling drugs rather than by superoxide. *Mol. Microbiol.* 79:1136–50
35. Gu M, Imlay JA. 2013. Superoxide poisons mononuclear iron enzymes by causing mismetallation. *Mol. Microbiol.* 89:123–34

36. Hassan HM, Fridovich I. 1977. Regulation of the synthesis of superoxide dismutase in *Escherichia coli*: induction by methyl viologen. *J. Biol. Chem.* 252:7667–72
37. Hassan HM, Fridovich I. 1978. Superoxide radical and the oxygen enhancement of the toxicity of paraquat in *Escherichia coli*. *J. Biol. Chem.* 253:8143–48
38. Helmann JD. 2014. Specificity of metal sensing: iron and manganese homeostasis in *Bacillus subtilis*. *J. Biol. Chem.* 289:28112–20
39. Heo YJ, Chung IY, Cho WJ, Lee BY, Kim JH, et al. 2010. The major catalase gene (*katA*) of *Pseudomonas aeruginosa* PA14 is under both positive and negative control of the global transactivator OxyR in response to hydrogen peroxide. *J. Bacteriol.* 192:381–90
40. Ieva R, Roncarati D, Metruccio MM, Seib KL, Scarlato V, Delany I. 2008. OxyR tightly regulates catalase expression in *Neisseria meningitidis* through both repression and activation mechanisms. *Mol. Microbiol.* 70:1152–65
41. Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nat. Rev. Microbiol.* 11:443–54
42. Imlay JA, Chin SM, Linn S. 1988. Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science* 240:640–42
43. Inbaraj JJ, Chignell CF. 2004. Cytotoxic action of juglone and plumbagin: a mechanistic study using HaCaT keratinocytes. *Chem. Res. Toxicol.* 17:55–62
44. Jacobson FS, Morgan RW, Christman MF, Ames BN. 1989. An alkyl hydroperoxide reductase from *Salmonella typhimurium* involved in the defense of DNA against oxidative damage: purification and properties. *J. Biol. Chem.* 264:1488–96
45. Jacquamet L, Traore DA, Ferrer JL, Proux O, Testemale D, et al. 2009. Structural characterization of the active form of PerR: insights into the metal-induced activation of PerR and Fur proteins for DNA binding. *Mol. Microbiol.* 73:20–31
46. Jang S, Imlay JA. 2007. Micromolar intracellular hydrogen peroxide disrupts metabolism by damaging iron-sulfur enzymes. *J. Biol. Chem.* 282:929–37
47. Jang S, Imlay JA. 2010. Hydrogen peroxide inactivates the *Escherichia coli* Isc iron-sulphur assembly system, and OxyR induces the Suf system to compensate. *Mol. Microbiol.* 78:1448–67
48. Jo I, Chung I-Y, Bae H-W, Kim J-S, Song S, et al. 2015. Structural details of OxyR peroxide-sensing mechanism. *PNAS* 112:6443–48
49. Kallifidas D, Thomas D, Doughty P, Paget MSB. 2010. The  $\sigma^R$  regulon of *Streptomyces oelicolor* A3(2) reveals a key role in protein quality control during disulphide stress. *Microbiology* 156:1661–72
50. Kehres DG, Janakiraman A, Schlauch JM, Maguire ME. 2002. Regulation of *Salmonella enterica* serovar Typhimurium *mntH* transcription by  $H_2O_2$ ,  $Fe^{2+}$ , and  $Mn^{2+}$ . *J. Bacteriol.* 184:3151–58
51. Kim J, Park W. 2014. Oxidative stress response in *Pseudomonas putida*. *Appl. Microbiol. Biotechnol.* 98:6933–46
52. Kim M-S, Dufour YS, Yoo JS, Cho Y-B, Park J-H, et al. 2012. Conservation of thiol-oxidative stress responses regulated by SigR orthologues in actinomycetes. *Mol. Microbiol.* 85:326–44
53. Kim S, Bang Y-J, Kim D, Lim JG, Oh MH, Choi SH. 2014. Distinct characteristics of OxyR2, a new OxyR-type regulator, ensuring expression of peroxiredoxin 2 detoxifying low levels of hydrogen peroxide in *Vibrio vulnificus*. *Mol. Microbiol.* 93:992–1009
54. Kobayashi K, Fujikawa M, Kozawa T. 2014. Binding of promoter DNA to SoxR protein decreases the reduction potential of the [2Fe-2S] cluster. *Biochemistry* 54:334–39
55. Kobayashi K, Tagawa S. 1999. Isolation of reductase for SoxR that governs an oxidative response regulon from *Escherichia coli*. *FEBS Lett.* 451:227–30
56. Koo MS, Lee JH, Ray SY, Yeo WS, Lee JW, et al. 2003. A reducing system of the superoxide sensor SoxR in *Escherichia coli*. *EMBO J.* 22:2614–22
57. Korshunov SS, Imlay JA. 2002. A potential role for periplasmic superoxide dismutase in blocking the penetration of external superoxide into the cytosol of phagocytosed bacteria. *Mol. Microbiol.* 43:95–106
58. Krapp AR, Humbert MV, Carrillo N. 2011. The *soxRS* response of *Escherichia coli* can be induced in the absence of oxidative stress and oxygen by modulation of NADPH content. *Microbiology* 157:957–65



59. Krapp AR, Rodriguez RE, Poli HU, Paladini DH, Palatnik JF, Carillo N. 2002. The flavoenzyme ferredoxin (flavodoxin)-NADP(H) reductase modulates NADP(H) homeostasis during the *soxRS* response of *Escherichia coli*. *J. Bacteriol.* 184:1474–80
60. Kuo CF, Mashino T, Fridovich I. 1987.  $\alpha,\beta$ -dihydroxyisovalerate dehydratase: a superoxide-sensitive enzyme. *J. Biol. Chem.* 262:4724–27
61. Lee C, Lee SM, Mukhopadhyay P, Kim SJ, Lee SC, et al. 2004. Redox regulation of OxyR requires specific disulfide bond formation involving a rapid kinetic reaction path. *Nat. Struct. Mol. Biol.* 11:1179–85
62. Lee JH, Lee KL, Yeo WS, Park SJ, Roe JH. 2009. SoxRS-mediated lipopolysaccharide modification enhances resistance against multiple drugs in *Escherichia coli*. *J. Bacteriol.* 191:4441–50
63. Lee J-W, Helmann JD. 2006. Biochemical characterization of the structural  $\text{Zn}^{2+}$  site in the *Bacillus subtilis* peroxide sensor PerR. *J. Biol. Chem.* 281:23567–78
64. Lee J-W, Helmann JD. 2006. The PerR transcription factor senses  $\text{H}_2\text{O}_2$  by metal-catalyzed histidine oxidation. *Nature* 440:363–67
65. Lemke RSS, Peterson AC, Ziegelhoffer EC, Westphall MS, Tjellström H, et al. 2014. Synthesis and scavenging role of furan fatty acids. *PNAS* 111:E3450–57
66. Lin CN, Syu WJ, Sun WS, Chen JW, Chen TH, et al. 2010. A role of *ygfZ* in the *Escherichia coli* response to plumbagin challenge. *J. Biomed. Sci.* 17:84
67. Liochev SI, Benov L, Touati D, Fridovich I. 1999. Induction of the *soxRS* regulon of *Escherichia coli* by superoxide. *J. Biol. Chem.* 274:9479–81
68. Liochev SI, Fridovich I. 1992. Fumarase C, the stable fumarase of *Escherichia coli*, is controlled by the *soxRS* regulon. *PNAS* 89:5892–96
69. Liochev SI, Fridovich I. 1997. Lucigenin luminescence as a measure of intracellular superoxide dismutase activity in *Escherichia coli*. *PNAS* 94:2891–96
70. Liochev SI, Hausladen A, Fridovich I. 1999. Nitroreductase A is regulated as a member of the *soxRS* regulon of *Escherichia coli*. *PNAS* 96:3537–39
71. Liu Y, Bauer SC, Imlay JA. 2011. The YaaA protein of the *Escherichia coli* OxyR regulon lessens hydrogen peroxide toxicity by diminishing the amount of intracellular unincorporated iron. *J. Bacteriol.* 193:2186–96
72. Loprosert S, Fuangthong M, Whangsuk W, Atichartpongkul S, Mongkolsuk S. 2000. Molecular and physiological analysis of an OxyR-regulated *abpC* promoter in *Xanthomonas campestris* pv. Phaseoli. *Mol. Microbiol.* 37:1504–14
73. Luo L, Qi MS, Yao SY, Cheng HP, Zhu JB, Yu GQ. 2005. Role of oxyR from *Sinorhizobium meliloti* in regulating the expression of catalases. *Acta Biochim. Biophys. Sin.* 37:421–28
74. Lynch R, Fridovich I. 1978. Permeation of the erythrocyte stroma by superoxide radical. *J. Biol. Chem.* 253:4697–99
75. Ma D, Cook DN, Alberti M, Pon NG, Nikaido H, Hearst JE. 1995. Genes *acrA* and *acrB* encode a stress-induced efflux system of *Escherichia coli*. *Mol. Microbiol.* 16:45–55
76. Mahavithakanont A, Charoenlap N, Namchaiw P, Eiamphungporn W, Chattrakarn S, et al. 2012. Novel roles of SoxR, a transcriptional regulator from *Xanthomonas campestris*, in sensing redox-cycling drugs and regulating a protective gene that have overall implications for bacterial stress physiology and virulence on a host plant. *J. Bacteriol.* 194:209–17
77. Mancini S, Imlay JA. 2015. The induction of two biosynthetic enzymes helps *Escherichia coli* sustain heme synthesis and activate catalase during hydrogen peroxide stress. *Mol. Microbiol.* 96:744–63
78. Martin RG, Rosner JL. 2011. Promoter discrimination at class I MarA regulon promoters mediated by glutamic acid 89 of the MarA transcriptional activator of *Escherichia coli*. *J. Bacteriol.* 193:506–15
79. Mehdy MC. 1994. Active oxygen species in plant defense against pathogens. *Plant Physiol.* 105:467–72
80. Mielecki D, Grzesiuk E. 2014. Ada response—a strategy for repair of alkylated DNA in bacteria. *FEMS Microbiol. Lett.* 355:1–11
81. Nakayama T, Yonekura S-I, Yonei S, Zhang-Akiyama Q-M. 2013. *Escherichia coli* pyruvate:flavodoxin oxidoreductase, YdbK—regulation of expression and biological roles in protection against oxidative stress. *Genes Genet. Syst.* 88:175–88



82. Naseer N, Shapiro JA, Chander M. 2014. RNA-Seq analysis reveals a six-gene SoxR regulon in *Streptomyces coelicolor*. *PLOS ONE* 9:e106181
83. Nelson KJ, Parsonage D, Hall A, Karplus PA, Poole LB. 2008. Cysteine pK<sub>a</sub> values for the bacterial peroxiredoxin AhpC. *Biochemistry* 47:12860–68
84. Nunoshiba T, Derojaswalker T, Wishnok JS, Tannenbaum SR, Demple B. 1993. Activation by nitric oxide of an oxidative stress response that defends *Escherichia coli* against activated macrophages. *PNAS* 90:9993–97
85. Outten FW, Djaman O, Storz G. 2004. A suf operon requirement for Fe-S cluster assembly during iron starvation in *Escherichia coli*. *Mol. Microbiol.* 52:861–72
86. Paget MS, Molle V, Cohen G, Aharonowitz Y, Buttner MJ. 2001. Defining the disulphide stress response in *Streptomyces coelicolor* A3(2): identification of the  $\sigma^R$  regulon. *Mol. Microbiol.* 42:1007–20
87. Paiva SRD, Figueiredo MR, Aragão TV, Kaplan MAC. 2003. Antimicrobial activity in vitro of plumbagin isolated from *Plumbago* species. *Mem. Inst. Oswaldo Cruz* 98:959–61
88. Palma M, Zurita J, Ferreras JA, Worgall S, Larone DH, et al. 2005. *Pseudomonas aeruginosa* SoxR does not conform to the archetypal paradigm for SoxR-dependent regulation of the bacterial oxidative stress adaptive response. *Infect. Immun.* 73:2958–66
89. Parent A, Caux-Thang C, Signor L, Clemancey M, Sethu R, et al. 2013. Single glutamate to aspartate mutation makes ferric uptake regulator (Fur) as sensitive to H<sub>2</sub>O<sub>2</sub> as peroxide resistance regulator (PerR). *Angew. Chem. Int. Ed.* 52:10339–43
90. Pomposiello PJ, Koutsolioutsou A, Carrasco D, Demple B. 2003. SoxRS-regulated expression and genetic analysis of the *yggX* gene of *Escherichia coli*. *J. Bacteriol.* 185:6624–32
91. Poole LB. 2007. The catalytic mechanism of peroxiredoxins. *Subcell. Biochem.* 44:61–81
92. Pullan ST, Gidley MD, Jones RA, Barrett J, Stevanin TM, et al. 2007. Nitric oxide in chemostat-cultured *Escherichia coli* is sensed by Fnr and other global regulators: Unaltered methionine biosynthesis indicates lack of S nitrosation. *J. Bacteriol.* 189:1845–55
93. Putnam CD, Arvai AS, Bourne Y, Tainer JA. 2000. Active and inhibited human catalase structures: ligand and NADPH binding and catalytic mechanism. *J. Mol. Biol.* 296:295–309
94. Rai P, Cole TD, Wemmer DE, Linn S. 2001. Localization of Fe<sup>2+</sup> at an RTGR sequence within a DNA duplex explains preferential cleavage by Fe<sup>2+</sup> and H<sub>2</sub>O<sub>2</sub>. *J. Mol. Biol.* 312:1089–101
95. Rau J, Stolz A. 2003. Oxygen-insensitive nitroreductases NfsA and NfsB of *Escherichia coli* function under anaerobic conditions as lawsone-dependent Azo reductases. *Appl. Environ. Microbiol.* 69:3448–55
96. Runde S, Moliere N, Heinz A, Maisonneuve E, Janczkowski A, et al. 2014. The role of thiol oxidative stress response in heat-induced protein aggregate formation during thermotolerance in *Bacillus subtilis*. *Mol. Microbiol.* 91:1036–52
97. Seaver LC, Imlay JA. 2001. Hydrogen peroxide fluxes and compartmentalization inside growing *Escherichia coli*. *J. Bacteriol.* 183:7182–89
98. Seki M, Iida K, Saito M, Nakayama H, Yoshida S. 2004. Hydrogen peroxide production in *Streptococcus pyogenes*: involvement of lactase oxidase and coupling with aerobic utilization of lactate. *J. Bacteriol.* 186:2046–51
99. Seth D, Hausladen A, Wang YJ, Stamler JS. 2012. Endogenous protein S-nitrosylation of *E. coli*: regulation by OxyR. *Science* 336:470–73
100. Sheplock R, Recinos DA, Mackow N, Dietrich LE, Chander M. 2012. Species-specific residues calibrate SoxR sensitivity to redox-active molecules. *Mol. Microbiol.* 87:368–81
101. Shin JH, Singh AK, Cheon DJ, Roe J-H. 2011. Activation of the SoxR regulon in *Streptomyces coelicolor* by the extracellular form of the pigmented antibiotic actinorhodin. *J. Bacteriol.* 193:75–81
102. Singh AK, Shin J-H, Lee K-L, Imlay JA, Roe J-H. 2013. Comparative study of SoxR activation by redox-active compounds. *Mol. Microbiol.* 90:983–96
103. Sobota JM, Gu M, Imlay JA. 2014. Intracellular hydrogen peroxide and superoxide poison 3-deoxy-D-arabinoheptulosonate 7-phosphate synthase, the first committed enzyme in the aromatic biosynthetic pathway of *Escherichia coli*. *J. Bacteriol.* 196:1980–91
104. Sobota JM, Imlay JA. 2011. Iron enzyme ribulose-5-phosphate 3-epimerase in *Escherichia coli* is rapidly damaged by hydrogen peroxide but can be protected by manganese. *PNAS* 108:5402–7

105. Teramoto H, Inui M, Yukawa H. 2013. OxyR acts as a transcriptional repressor of hydrogen peroxide-inducible antioxidant genes in *Corynebacterium glutamicum* R. *FEBS J.* 280:3298–312
106. Tosatto SCE, Bosello V, Fogolari F, Mauri P, Roveri A, et al. 2008. The catalytic site of glutathione peroxidases. *Antiox. Redox Rep.* 10:1515–26
107. Traore DA, Ghaouani AE, Ilango S, Dupuy J, Jacquamet L, et al. 2006. Crystal structure of the apo-PerR-Zn protein from *Bacillus subtilis*. *Mol. Microbiol.* 61:1211–19
108. Tsaneva IR, Weiss B. 1990. *soxR*, a locus governing a superoxide response regulon in *Escherichia coli* K-12. *J. Bacteriol.* 172:4197–205
109. Turner JM, Messenger AJ. 1986. Occurrence, biochemistry, and physiology of phenazine pigment production. *Adv. Microb. Physiol.* 27:211–75
110. Varghese S, Tang Y, Imlay JA. 2003. Contrasting sensitivities of *Escherichia coli* aconitases A and B to oxidation and iron depletion. *J. Bacteriol.* 185:221–30
111. Varghese S, Wu A, Park S, Imlay KRC, Imlay JA. 2007. Submicromolar hydrogen peroxide disrupts the ability of Fur protein to control free-iron levels in *Escherichia coli*. *Mol. Microbiol.* 64:822–30
112. Wang Y, Kern SE, Newman DK. 2010. Endogenous phenazine antibiotics promote anaerobic survival of *Pseudomonas aeruginosa* via extracellular electron transfer. *J. Bacteriol.* 192:365–69
113. Wang Y, Wilks JC, Danhorn T, Ramos I, Croal L, Newman DK. 2011. Phenazine-1-carboxylic acid promotes bacterial biofilm development via ferrous iron acquisition. *J. Bacteriol.* 193:3606–17
114. Watanabe S, Kita A, Kobayashi K, Miki K. 2008. Crystal structure of the [2Fe-2S] oxidative-stress sensor SoxR bound to DNA. *PNAS* 105:4121–26
115. Winterbourn CC, Hampton MB, Livesey JH, Kettle AJ. 2006. Modeling the reactions of superoxide and myeloperoxidase in the neutrophil phagosome: implications for microbial killing. *J. Biol. Chem.* 281:39860–69
116. Winterbourn CC, Metodiewa D. 1999. Reactivity of biologically important thiol compounds with superoxide and hydrogen peroxide. *Free Rad. Biol. Med.* 27:322–28
117. Yamamoto K, Ishihama A, Busby SJ, Grainger DC. 2011. The *Escherichia coli* K-12 MntR miniregulon includes *dps*, which encodes the major stationary-phase DNA-binding protein. *J. Bacteriol.* 193:1477–80
118. Zheng M, Aslund F, Storz G. 1998. Activation of the OxyR transcription factor by reversible disulfide bond formation. *Science* 279:1718–21
119. Zheng M, Doan B, Schneider TD, Storz G. 1999. OxyR and SoxRS regulation of *fur*. *J. Bacteriol.* 181:4639–43
120. Zheng M, Wang X, Templeton LJ, Smulski DR, LaRossa RA, Storz G. 2001. DNA microarray-mediated transcriptional profiling of the *Escherichia coli* response to hydrogen peroxide. *J. Bacteriol.* 183:4562–70
121. Zuber P. 2009. Management of oxidative stress in *Bacillus*. *Annu. Rev. Microbiol.* 63:575–97



# Contents

My Lifelong Passion for Biochemistry and Anaerobic Microorganisms <i>Rudolf Kurt Thauer</i> .....	1
The Pyromaniac Inside You: <i>Salmonella</i> Metabolism in the Host Gut <i>Fabian Rivera-Chávez and Andreas J. Bäumler</i> .....	31
Regulation of Transcript Elongation <i>Georgiy A. Belogurov and Irina Artsimovitch</i> .....	49
<i>Candida albicans</i> Biofilms and Human Disease <i>Clarissa J. Nobile and Alexander D. Johnson</i> .....	71
Transcription Factors That Defend Bacteria Against Reactive Oxygen Species <i>James A. Imlay</i> .....	93
Bacterial Proteasomes <i>Jordan B. Jastrab and K. Heran Darwin</i> .....	109
The Organellar Genomes of <i>Chromera</i> and <i>Vitrella</i> , the Phototrophic Relatives of Apicomplexan Parasites <i>Miroslav Oborník and Julius Lukeš</i> .....	129
The Gut Microbiota of Termites: Digesting the Diversity in the Light of Ecology and Evolution <i>Andreas Brune and Carsten Dietrich</i> .....	145
Microsporidia: Eukaryotic Intracellular Parasites Shaped by Gene Loss and Horizontal Gene Transfers <i>Nicolas Corradi</i> .....	167
Anthrax Pathogenesis <i>Mabtab Moayeri, Stephen H. Leppla, Catherine Vrentas, Andrei P. Pomerantsev, and Shibui Liu</i> .....	185
CRISPR-Cas: New Tools for Genetic Manipulations from Bacterial Immunity Systems <i>Wenyan Jiang and Luciano A. Marraffini</i> .....	209

Perception and Homeostatic Control of Iron in the Rhizobia and Related Bacteria <i>Mark R. O'Brian</i> .....	229
Thymineless Death Lives On: New Insights into a Classic Phenomenon <i>Arkady Khodursky, Elena C. Guzmán, and Philip C. Hanawalt</i> .....	247
Membrane-Coupled mRNA Trafficking in Fungi <i>Carl Haag, Benedikt Steuten, and Michael Feldbrügge</i> .....	265
Molecular Pathogenesis of <i>Ehrlichia chaffeensis</i> Infection <i>Yasuko Rikihisa</i> .....	283
Microbiology Meets Big Data: The Case of Gut Microbiota–Derived Trimethylamine <i>Gwen Falony, Sara Vieira-Silva, and Jeroen Raes</i> .....	305
The Unique Molecular Choreography of Giant Pore Formation by the Cholesterol-Dependent Cytolysins of Gram-Positive Bacteria <i>Rodney K. Tweten, Eileen M. Hotze, and Kristin R. Wade</i> .....	323
Ion Regulation in the Malaria Parasite <i>Kiaran Kirk</i> .....	341
Physics of Intracellular Organization in Bacteria <i>Ned S. Wingreen and Kerwyn Casey Huang</i> .....	361
Stochastic Switching of Cell Fate in Microbes <i>Thomas M. Norman, Nathan D. Lord, Johan Paulsson, and Richard Losick</i> .....	381
Assembly of the Mycobacterial Cell Wall <i>Monika Jankute, Jonathan A.G. Cox, James Harrison, and Gurdyal S. Besra</i> .....	405
Mechanisms of Bacterial Colonization of the Respiratory Tract <i>Steven J. Siegel and Jeffrey N. Weiser</i> .....	425
Interactions Between the Gastrointestinal Microbiome and <i>Clostridium difficile</i> <i>Casey M. Theriot and Vincent B. Young</i> .....	445
Lytic Cycle of <i>Toxoplasma gondii</i> : 15 Years Later <i>Ira Blader, Bradley Coleman, Chun-Ti Chen, and Marc-Jan Gubbels</i> .....	463
Septins and Generation of Asymmetries in Fungal Cells <i>Anum Khan, Molly McQuilken, and Amy S. Gladfelter</i> .....	487
How Is Fe-S Cluster Formation Regulated? <i>Erin L. Mettert and Patricia J. Kiley</i> .....	505

Protein Phosphatases of Pathogenic Bacteria: Role in Physiology  
and Virulence  
*Andaleeb Sajid, Gunjan Arora, Anshika Singhal, Vipin C. Kalia,  
and Yogendra Singh* ..... 527

**Index**

Cumulative Index of Contributing Authors, Volumes 65–69 ..... 549

**Errata**

An online log of corrections to *Annual Review of Microbiology* articles may be found at  
<http://www.annualreviews.org/errata/micro>