

# Optimization Theory and Algorithms

Instructor: Prof. LIAO, Guocheng (廖国成)

Email: [liaogch6@mail.sysu.edu.cn](mailto:liaogch6@mail.sysu.edu.cn)

**School of Software Engineering  
Sun Yat-sen University**


# Outline

- Convex function
- Equivalent characterization
- Examples of convex function
- Properties of convex function
- Convexity-preserving operations

# Convex function

A function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if **dom**  $f$  is a convex set and for all  $x, y$ , and  $0 \leq \theta \leq 1$ :

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$



Function lies below the line segment between  $(x, f(x))$  and  $(y, f(y))$ .

**Strict convex** : for all  $x \neq y$ , and  $0 < \theta < 1$ :

$$f(\theta x + (1 - \theta)y) < \theta f(x) + (1 - \theta)f(y)$$

A function  $f$  is concave if  $-f$  is convex.


# Examples

- Univariate function:
  - Exponential function  $f(x) = e^{ax}$ , for any  $a$
  - Logarithmic function  $f(x) = \log x$  over  $\mathbb{R}_{++}$
  - Quadratic function  $f(x) = ax^2 + bx + c$ ,  $a > 0$
  - Power function  $f(x) = x^a$  is convex for  $a \geq 1$  or  $a \leq 0$ , and is concave for  $0 \leq a \leq 1$ .
- Affine function:  $f(x) = Ax + b$
- Quadratic function  $f(x) = x^T Qx + q^T x$ , where  $Q \succcurlyeq 0$
- Squared loss  $f(x) = \|Ax - b\|_2^2$  ( $A^T A$  is always positive semidefinite)
- Norm:  $\|x\|_p \triangleq (\sum_{i=1}^n |x_i|^p)^{1/p}$ , where  $p \geq 1$

# First-order characterization

Suppose  $f$  is differentiable. Then  $f$  is convex *if and only if*  $\text{dom } f$  is convex and  $f(y) \geq f(x) + \nabla f(x)^T (y - x)$ , for all  $x, y \in \text{dom } f$ .

$$\nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{bmatrix} \in \mathbb{R}^n$$



# Second-order characterization

Suppose  $f$  is twice differentiable. Then  $f$  is convex **if and only if**  $\text{dom } f$  is convex and its Hessian matrix is positive semidefinite:

$$\nabla^2 f(x) \succcurlyeq 0, \text{ for all } x \in \text{dom } f.$$

$$\nabla^2 f(x) = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix} \in \mathbb{R}^{n \times n}$$

Strict convex  $\Leftrightarrow \nabla^2 f(x) \succ 0$   
 Concave  $\Leftrightarrow \nabla^2 f(x) \preccurlyeq 0$

Quadratic function:  $f(x) = \frac{1}{2}x^T Qx + q^T x$

$$\nabla f(x) = Qx + q \quad \nabla^2 f(x) = Q \quad f(x) \text{ is convex} \Leftrightarrow Q \succcurlyeq 0$$

Squared loss:  $f(x) = \|Ax - b\|_2^2$

$$\nabla f(x) = 2A^T(Ax - b) \quad \nabla^2 f(x) = 2A^T A \quad f(x) \text{ is convex}$$

# Examples

Univariate function:

- Exponential function  $f(x) = e^{ax}$ , for any  $a$ .  $f''(x) = a^2 e^{ax} > 0$
- Logarithmic function  $f(x) = \log x$  over  $\mathbb{R}_{++}$ .  $f''(x) = -\frac{1}{x^2} < 0$
- Quadratic function  $f(x) = ax^2 + bx + c$ ,  $a > 0$ .  $f''(x) = a > 0$
- Power function  $f(x) = x^a$  is convex for  $a \geq 1$  or  $a \leq 0$ , and is concave for  $0 \leq a \leq 1$ .  
 $f''(x) = a(a-1)x^{a-2}$
- Negative entropy  $f(x) = x \log x$ .  $f'(x) = \log x + 1$   $f''(x) = \frac{1}{x} > 0$

# Examples

- Quadratic-over-linear function:  $f(x, y) = x^2/y$ , where  $y > 0$ .

$$\nabla^2 f(x) = \begin{bmatrix} \frac{2}{y} & -\frac{2x}{y^2} \\ -\frac{2x}{y^2} & \frac{2x^2}{y^3} \end{bmatrix} = \frac{2}{y^3} \begin{bmatrix} y^2 & -xy \\ -xy & x^2 \end{bmatrix} = \frac{2}{y^3} \begin{bmatrix} y \\ -x \end{bmatrix} \begin{bmatrix} y \\ -x \end{bmatrix}^T \succcurlyeq 0$$

- Norm:  $f(x) = \|x\|_p \triangleq (\sum_{i=1}^n |x_i|^p)^{1/p}$ , where  $p \geq 1$

$$f(\theta x + (1 - \theta)y) \leq f(\theta x) + f((1 - \theta)y) = \theta f(x) + (1 - \theta)f(y)$$

- Max:  $f(x) = \max_i x_i$

$$\begin{aligned} f(\theta x + (1 - \theta)y) &= \max_i (\theta x_i + (1 - \theta)y_i) \leq \theta \max_i x_i + (1 - \theta) \max_i y_i \\ &= \theta f(x) + (1 - \theta)f(y) \end{aligned}$$



# Restriction to a line

$f: \mathbb{R}^n \rightarrow \mathbb{R}$  is convex if and only if the function  $g: \mathbb{R} \rightarrow \mathbb{R}$

$$g(t) = f(x + ty), \text{ dom } g = \{t | x + ty \in \text{dom } f\}$$

is convex in  $t$  for any  $x \in \text{dom } f$ .

Example: Log-determinant.  $f: \mathbf{S}_{++}^n \rightarrow \mathbb{R}$ ,  $f(X) = \log \det X$  is concave

$$\begin{aligned} g(t) &= f(X + tV) = \log \det(X + tV) \\ &= \log \det(X^{1/2}IX^{1/2} + tX^{1/2}X^{-1/2}VX^{-1/2}X^{1/2}) \\ &= \log \det(X^{1/2}(I + tX^{-1/2}VX^{-1/2})X^{1/2}) \\ &= 2\log \det(X^{1/2}) + \log \det(I + tX^{-1/2}VX^{-1/2}) \\ &= \log \prod_{i=1}^n (1 + t\lambda_i) + 2\log \det(X^{1/2}) \\ &= \sum_{i=1}^n \log(1 + t\lambda_i) + 2\log \det(X^{1/2}) \end{aligned}$$

Eigenvalue of  $X^{-1/2}VX^{-1/2}$

$$g'(t) = \sum_{i=1}^n \frac{\lambda_i}{1 + t\lambda_i} \quad g''(t) = -\sum_{i=1}^n \frac{\lambda_i^2}{(1 + t\lambda_i)^2} < 0$$

# Jensen's inequality

Basic: if  $f$  is convex, then for any  $0 \leq \theta \leq 1$ :

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$

Extension to more than two points: if  $f$  is convex, then for any  $\theta_1, \dots, \theta_k \in [0,1]$  such that  $\sum_{i=1}^k \theta_i = 1$

$$f\left(\sum_{i=1}^k \theta_i x_i\right) \leq \sum_{i=1}^k \theta_i f(x_i)$$

Expression with expectation: if  $f$  is convex, then:

$$f(E[x]) \leq E[f(x)]$$

Example:  $\frac{a+b}{2} \geq \sqrt{ab}$      $f(x) = -\log x$  is convex  $\Rightarrow -\log \frac{a+b}{2} \leq -\frac{1}{2}(\log a + \log b)$   
i.e.,  $-\frac{a+b}{2} \geq -\sqrt{ab}$

# Sublevel set and epigraph:

$t$  – sublevel set of a function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$ :


$$C_t = \{x \in \mathbf{dom} f \mid f(x) \leq t\}$$

Sublevel sets of a convex function are convex sets (inverse is false).

A function is quasi-convex if all its sublevel sets are convex.

Epigraph of a function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$ :

$$\mathbf{epi} f = \{(x, t) \in \mathbb{R}^{n+1} \mid x \in \mathbf{dom} f, f(x) \leq t\}$$



$f$  is convex if and only if  $\mathbf{epi} f$  is a convex set

# Convexity-preserving operations

- Non-negative weighted sum: let  $f_1, \dots, f_m$  be convex functions and  $\alpha_1, \dots, \alpha_m \in \mathbb{R}$  be non-negative values. Then the following function is convex:

$$f(x) = \sum_{i=1}^m \alpha_i f_i(x)$$

- Affine composition: let  $f$  be a convex function. Then  $g(x) \triangleq f(Ax + b)$  is convex

E.g.,  $f(x) = \|Ax + b\|_2^2$  is convex

# Convexity-preserving operations

Composition: let  $f$  be the composition of  $g: \mathbb{R}^n \rightarrow \mathbb{R}$  and  $h: \mathbb{R} \rightarrow \mathbb{R}$ ,

$$f(x) = h(g(x))$$

- $g$  is **convex** and  $h$  is **non-decreasing**: then  $f$  is convex.
- $g$  is **concave** and  $h$  is **non-increasing**: then  $f$  is convex.

Examples:

- $\exp g(x)$  is convex if  $g$  is convex.
- $1/g(x)$  is convex if  $g$  is concave and positive.

# Convexity-preserving operations

Pointwise maximum: if  $f_1, \dots, f_m$  are convex function, then

$$f(x) = \max\{f_1(x), \dots, f_m(x)\}$$

is convex

Examples:

- Piecewise-linear function:  $f(x) = \max_i (a_i^T x + b_i)$  is convex.
- Sum of  $r$  largest components of  $x \in \mathbb{R}^n$ : let  $x_{[1]} \leq x_{[2]} \leq \dots \leq x_{[n]}$ . Then

$$f(x) = \sum_{i=1}^r x_{[i]}$$

is convex.



$$\max\{x_{i_1} + x_{i_2} + \dots + x_{i_r} \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_r\}$$

Sum of any  $r$  different components of  $x$  is convex function  
 $\Rightarrow$  pointwise maximum of them is convex

# Convexity-preserving operations

If  $f(x, y)$  is convex in  $x$  for each  $y \in C$ , then

$$g(x) = \sup_{y \in C} f(x, y)$$

is convex

Examples:

- Distance to the farthest point of a set:  $f(x) = \sup_{y \in S} \|x - y\|$
- Support function of a set:  $S_C(x) = \sup_{y \in C} y^T x$

# How to verify convexity of a function

- By definition
- Hessian matrix is positive semidefinite
- By convexity-preserving operations




# Conjugate function

The conjugate function of a function  $f$  is

$$f^*(y) = \sup_{x \in \text{dom } f} y^T x - f(x)$$

The domain of conjugate function is where the supremum is finite.



In one-dimensional space, the conjugate function is the maximum gap between the linear function  $xy$  and  $f(x)$ .

The conjugate function is always convex (pointwise supremum of a family of affine functions)

# Conjugate function: examples

- Affine function  $f(x) = ax + b$

$$f^*(y) = \sup_{x \in \mathbb{R}} xy - ax - b = b, y = a$$

If  $y \neq a$ , then  $xy - ax - b$  is **unbounded** for  $x \in \mathbb{R}$

- Negative logarithmic  $f(x) = -\log x$

$$f^*(y) = \sup_{x>0} xy + \log x = -1 - \log(-y), y < 0$$

If  $y \geq 0$ , then  $xy + \log x$  is **unbounded** as  $x \rightarrow +\infty$

- Exponential function  $f(x) = e^x$

$$f^*(y) = \sup_x xy - e^x = y \log(y) - y, y \geq 0$$

If  $y < 0$ , then  $xy - e^x$  is **unbounded** for  $x \rightarrow -\infty$

- Strictly convex quadratic  $f(x) = \frac{1}{2}x^T Qx$  with  $Q \succ 0$

$$f^*(y) = \sup_x y^T x - \frac{1}{2}x^T Qx = \frac{1}{2}y^T Q^{-1}y, \text{ for all } y$$