

UECM1304 TEST 2 MARKING GUIDE

Name: _____ Student ID: _____ Mark: _____ /20

COURSE CODE & COURSE TITLE: UECM1304 DISCRETE MATHEMATICS WITH APPLICATIONS

FACULTY: LKC FES, UTAR COURSE: AM

TRIMESTER: JUN 2025 LECTURER: LIEW HOW HUI

Instruction: Answer all questions in the space provided. **If you do not write your answer in the space provided, you will get ZERO mark.** An answer without working steps may also receive ZERO mark.

1. CO3: Demonstrate various proof-techniques.C3

(a) Use direct proof to show that there are integers m and n such that $2674m + 5257n = 7$.

[**Note:** You need to use extended Euclidean algorithm to find out m and n rather than simply guessing. Guessing the answer is regarded as cheating and only 0.5 marks will be awarded.] (4 marks)

Ans. $5257 = 2674 \times 1 + 2583$

$$2674 = 2583 \times 1 + 91$$

$$2583 = 91 \times 28 + 35$$

$$91 = 35 \times 2 + 21$$

$$35 = 21 \times 1 + 14$$

$$21 = 14 \times 1 + 7$$

$$14 = 7 \times 2 + 0$$

[1.5 marks]

$$7 = 21 - 14 \times 1$$

$$7 = 21 - (35 - 21) = 21 \times 2 - 35$$

$$7 = (91 - 35 \times 2) \times 2 - 35 = 91 \times 2 - 35 \times 5$$

$$7 = 91 \times 2 - (2583 - 91 \times 28) \times 5 = 91 \times 142 - 2583 \times 5$$

$$7 = (2674 - 2583) \times 142 - 2583 \times 5 = 2674 \times 142 - 2583 \times 147$$

$$7 = 2674 \times 142 - (5257 - 2674) \times 147 = 2674 \times 289 - 5257 \times 147$$
 [2 marks]

Therefore, there are $m = 289$ and $n = -147$ such that $2674m + 5257n = 7$[0.5 mark]

- (b) Use contrapositive proof to show that for all integers n , if $7n + 3$ is even then n is odd. (2 marks)

Ans. Suppose n is an even integer. Then $n = 2k$ for some integer k [0.5 mark]
 $7n + 3 = 14k + 3 = 2(7k + 1) + 1 = 2k' + 1$. Here $k' = 7k + 1$ is some integer. ... [1 mark]
 By definition, $7n + 3$ is odd. [0.5 mark]

- (c) Use the Principle of Mathematical Induction to prove the equality

$$2 + 3 + \cdots + n = \frac{(n-1)(n+2)}{2}$$

for integers $n \geq 2$. (3 marks)

Ans. Base Step:

When $n = 2$, $\text{RHS} = \frac{(2-1)(2+2)}{2} = \frac{4}{2} = 2 = \text{LHS}$ [1 mark]

Induction Step:

Assume $2 + 3 + \cdots + k = \frac{(k-1)(k+2)}{2}$ for $k \geq 2$ [0.3 mark]

$$2 + 3 + \cdots + k + (k+1) = \frac{(k-1)(k+2)}{2} + (k+1) \quad [0.5 \text{ mark}]$$

$$= \frac{(k-1)(k+2) + 2(k+1)}{2} = \frac{k^2 + k - 2 + 2k + 2}{2} \quad [0.5 \text{ mark}]$$

$$= \frac{k^2 + 3k}{2} = \frac{k(k+3)}{2} = \frac{((k+1)-1)((k+1)+2)}{2} \quad [0.5 \text{ mark}]$$

Hence, by the Principle of Mathematical Induction, $2 + 3 + \cdots + n = \frac{(n-1)(n+2)}{2}$ for $n \geq 2$.
 [0.2 mark]

(d) Use the Principle of Mathematical Induction to prove that the inequality

$$\prod_{i=1}^n \left(1 - \frac{1}{3^i}\right) \geq \frac{1}{9} + \frac{1}{3^{n+1}}$$

for integers $n \geq 1$. Here $\prod_{i=1}^n f(i) = f(1) \times f(2) \times \cdots \times f(n)$ is the product of n terms for some expression f . (3 marks)

Ans. Base Step:

When $n = 1$, LHS = $1 - \frac{1}{3} = \frac{2}{3} \geq \frac{1}{9} + \frac{1}{9} = \frac{2}{9}$ = RHS [1 mark]

Induction Step:

Assume $\prod_{i=1}^k \left(1 - \frac{1}{3^i}\right) \geq \frac{1}{9} + \frac{1}{3^{k+1}}$ for $k \geq 1$ [0.3 mark]

$$\prod_{i=1}^{k+1} \left(1 - \frac{1}{3^i}\right) = \left[\prod_{i=1}^k \left(1 - \frac{1}{3^i}\right) \right] \left(1 - \frac{1}{3^{k+1}}\right) \geq \left[\frac{1}{9} + \frac{1}{3^{k+1}} \right] \left(1 - \frac{1}{3^{k+1}}\right) \quad [0.5 \text{ mark}]$$

$$= \frac{1}{9} + \frac{1}{3^{k+1}} - \frac{1}{3^{k+3}} - \frac{1}{3^{2k+2}} = \frac{1}{9} + \frac{3}{3^{k+2}} - \frac{\frac{1}{3}}{3^{k+2}} - \frac{\frac{1}{3^k}}{3^{k+2}} \quad [0.5 \text{ mark}]$$

$$\geq \frac{1}{9} + \frac{1}{3^{k+2}} \left[3 - \frac{1}{3} - \frac{1}{3} \right] \geq \frac{1}{9} + \frac{1}{3^{k+2}} \quad [0.5 \text{ mark}]$$

Note that $k \geq 1 \Rightarrow 3^k \geq 3 \Rightarrow -\frac{1}{3^k} \geq -\frac{1}{3}$.

Hence, by the Principle of Mathematical Induction, $\prod_{i=1}^n \left(1 - \frac{1}{3^i}\right) \geq \frac{1}{9} + \frac{1}{3^{n+1}}$ for $n \geq 1$.

..... [0.2 mark]

2. CO4. Express relations correctly with their mathematical properties. C2

(a) A set $A = \{a, b, c, d, e\}$ has a relation

$$R = \{(a, a), (a, c), (a, d), (b, b), (b, c), (b, e), (c, a), (c, c), (d, a), (d, b), (d, d), (e, a), (e, e)\}$$

Answer the following questions by filling the box with Yes or No.

- | | |
|---|----------------------------------|
| i. Is R reflexive? | <input type="text" value="Yes"/> |
| ii. Is R symmetric? | <input type="text" value="No"/> |
| iii. Is R a partial order relation? | <input type="text" value="No"/> |

Provide justifications for the above questions in the box below if your answer is No.

($3 \times 0.3 + 0.6 = 1.5$ mark)

Ans. ii. R is not symmetric because $(b, c) \in R$ but $(c, b) \notin R$ [0.3 mark]

iii. R is not a partial order because it is not antisymmetric since $(a, c), (c, a) \in R$ but $a \neq c$ [0.3 mark]


Note that the lecturer uses a symmetric pair for disclaiming anti-symmetry in lecture.
This may not be precise enough.

Anti-symmetry: $\forall x \forall y (xRy \wedge yRx \rightarrow x = y)$

Not anti-symmetry: $\sim [\forall x \forall y (xRy \wedge yRx \rightarrow x = y)] \equiv \exists x \exists y [(xRy \wedge yRx \wedge x \neq y)]$

(b) Let A be the set of all the monic polynomial factors of the monic polynomial $(x+2)^2(x+3)(x+4)$ and the partial order relation $p \preceq q$ be p divides q for any monic polynomials p, q . Sketch the Hasse diagram of the poset (A, \preceq) . (1.5 marks)

Ans.



List out all elements of A in proper ordering [0.7 mark]

Correct links in the Hasse diagram [0.8 mark]

(c) Consider a set $A = \{a, b, c, d, e, f\}$ with the following relation

$$R = \{(a, c), (a, f), (b, c), (b, e), (c, b), (d, c), (e, d), (f, a)\}.$$

- i. Write down the **matrix representation** of R and then (I) determine if the relation R is **irreflexive** with justification; (II) determine if the relation R is **anti-symmetric** with justifications. (2 marks)

Ans. $M_R =$

	a	b	c	d	e	f
a	0	0	1	0	0	1
b	0	0	1	0	1	0
c	0	1	0	0	0	0
d	0	0	1	0	0	0
e	0	0	0	1	0	0
f	1	0	0	0	0	0

..... [1 mark]

(I) Since the diagonal of M_R are all zero, R is irreflexive. [0.5 mark]

(II) Since $(b, c), (c, b) \in R$ but $b \neq c$, R is **not anti-symmetric**. [0.5 mark]

- ii. Apply the **Warshall algorithm** to find the matrix representation of the **transitive closure** of R . (3 marks)

Ans. Step 1:

	a	b	c	d	e	f
a	0	0	1	0	0	1
b	0	0	1	0	1	0
c	0	1	0	0	0	0
d	0	0	1	0	0	0
e	0	0	0	1	0	0
f	1	0	1	0	0	1

..... [0.5 mark]

Step 2:

	a	b	c	d	e	f
a	0	0	1	0	0	1
b	0	0	1	0	1	0
c	0	1	1	0	1	0
d	0	0	1	0	0	0
e	0	0	0	1	0	0
f	1	0	1	0	0	1

..... [0.5 mark]

Step 3:

	a	b	c	d	e	f
a	0	1	1	0	1	1
b	0	1	1	0	1	0
c	0	1	1	0	1	0
d	0	1	1	0	1	0
e	0	0	0	1	0	0
f	1	1	1	0	1	1

..... [0.5 mark]

Step 4:

$$\begin{array}{c}
 \\
 \\
 \\
 \\
 \\
 \\
 \end{array}
 \begin{array}{c}
 a \\
 b \\
 c \\
 d \\
 e \\
 f
 \end{array}
 \begin{array}{c}
 a \quad b \quad c \quad d \quad e \quad f \\
 \left(\begin{array}{cccccc}
 0 & 1 & 1 & 0 & 1 & 1 \\
 0 & 1 & 1 & 0 & 1 & 0 \\
 0 & 1 & 1 & 0 & 1 & 0 \\
 0 & 1 & 1 & 0 & 1 & 0 \\
 0 & \boxed{1} & \boxed{1} & 1 & \boxed{1} & 0 \\
 1 & 1 & 1 & 0 & 1 & 1
 \end{array} \right)
 \end{array}
 \dots\dots\dots [0.5 \text{ mark}]$$

Step 5:

$$\begin{array}{c}
 \\
 \\
 \\
 \\
 \\
 \\
 \end{array}
 \begin{array}{c}
 a \\
 b \\
 c \\
 d \\
 e \\
 f
 \end{array}
 \begin{array}{c}
 a \quad b \quad c \quad d \quad e \quad f \\
 \left(\begin{array}{cccccc}
 0 & 1 & 1 & \boxed{1} & 1 & 1 \\
 0 & 1 & 1 & \boxed{1} & 1 & 0 \\
 0 & 1 & 1 & \boxed{1} & 1 & 0 \\
 0 & 1 & 1 & \boxed{1} & 1 & 0 \\
 0 & 1 & 1 & 1 & 1 & 0 \\
 1 & 1 & 1 & \boxed{1} & 1 & 1
 \end{array} \right)
 \end{array}
 \dots\dots\dots [0.5 \text{ mark}]$$

Step 6: The matrix representation of the transitive closure is

$$\begin{array}{c}
 \\
 \\
 \\
 \\
 \\
 \\
 \end{array}
 \begin{array}{c}
 a \\
 b \\
 c \\
 d \\
 e \\
 f
 \end{array}
 \begin{array}{c}
 a \quad b \quad c \quad d \quad e \quad f \\
 \left(\begin{array}{cccccc}
 \boxed{1} & 1 & 1 & 1 & 1 & 1 \\
 0 & 1 & 1 & 1 & 1 & 0 \\
 0 & 1 & 1 & 1 & 1 & 0 \\
 0 & 1 & 1 & 1 & 1 & 0 \\
 0 & 1 & 1 & 1 & 1 & 0 \\
 1 & 1 & 1 & 1 & 1 & 1
 \end{array} \right)
 \end{array}
 \dots\dots\dots [0.5 \text{ mark}]$$