


Lab 1: Convolution

ILIAS
TOUGUI

We want to convolve a rectangle with itself
a rectangle is defined as:

$$f(t) = \begin{cases} 1 & ; -1 \leq t \leq 1 \\ 0 & ; \text{otherwise} \end{cases}$$



1) the convolution is defined mathematically as:

$$g(x) = f(x) * h(x) = \int_{-\infty}^{+\infty} f(t) \cdot h(x-t) dt$$

In this case we have $f(t) = h(t)$

2) Let's define the support of each function


$f(t)$ is non zero in $-1 \leq t \leq 1$ $[-1, 1]$

$f(x-t)$ is non zero in $-1 \leq x-t \leq 1$
 $-x-1 \leq -t \leq -x+1$
 $x-1 \leq t \leq x+1$

the support of $h(x-t)$ is $[x-1, x+1]$

3) Compute the convolution for each of the following cases :


a) No overlap



We have no overlap when : $x-1 = -3 \Rightarrow x = -2$
 or $x+1 = -1 \Rightarrow x = -2$
 and $x-1 = 1 \Rightarrow x = 2$
 or $x+1 = 3 \Rightarrow x = +2$

$$g(x) = 0 ; x \leq -2 \text{ or } x \geq 2$$

b) 1st partial overlap




$$g(x) = \int_{-1}^{x+1} f(t) \cdot h(x-t) dt$$

$$= \int_{-1}^{x+1} 1 \times 1 dt = [t]_{-1}^{x+1} = x+1 - (-1) = x+2$$

$$g(x) = x+2 \text{ when } -2 \leq x \leq 0$$

Case 3) : Full overlap




We have a full overlap at $x=0$

$$g(x) = \int_{-a}^a f(t) \cdot h(x-t) dt$$

$$= \int_{-1}^1 1 \times 1 dt = [t]_{-1}^1 = 1 - (-1) = 2$$

$$\text{at } x=0 \quad g(x) = 2$$

d) Case 4 : 2nd partial overlap.




$$g(x) = \int_{x-1}^1 f(t) \cdot h(x-t) dt$$

$$= \int_{x-1}^1 1 \times 1 dt = [t]_{x-1}^1 = 1 - x + 1 = -x + 2$$

$$g(x) = +2 - x \quad \text{when } 0 \leq x \leq 2$$

$$g(x) = \begin{cases} x+2 & ; -2 \leq x \leq 0 \\ 2 & ; x = 0 \\ -x+2 & ; 0 \leq x \leq 2 \\ 0 & ; \text{otherwise} \end{cases}$$



Lab A : Convolution

we want to convolve a rectangular pulse with a triangle, the convolution is defined as:

$$g(x) = f(x) * h(x) = \int_{-\infty}^{+\infty} f(t) \cdot h(x-t) dt$$


because the convolution is commutative meaning that:

$$f * g = g * f$$

$$g(x) = h(x) * f(x) = \int_{-\infty}^{+\infty} h(t) \cdot f(x-t) dt$$


1) let's define and draw the functions:

$$f(t) = \begin{cases} 1+t & ; -1 \leq t \leq 0 \\ 1-t & ; 0 \leq t \leq 1 \\ 0 & ; \text{otherwise} \end{cases}$$



$$h(x-t) = \begin{cases} 1 & ; -1 \leq x-t \leq 1 \\ 0 & ; \text{otherwise} \end{cases}$$


$$= \begin{cases} 1 & ; x-1 \leq t \leq x+1 \\ 0 & ; \text{otherwise} \end{cases}$$




Triangle support : $[-1, 1]$
 Rectangle support : $[x-1, x+1]$

2) Draw and compute the convolution for each of the following cases:

a) case 1) : No overlaps



$$x-1 = -3 \text{ and } x+1 = -1 \\ x = -2 \text{ and } x = -2$$




$$x-1 = 1 \text{ and } x+1 = 3 \\ x = 2 \text{ and } x = 2$$

No overlaps when $x \leq -2$ and $x \geq 2$

$$g(x) = 0$$

b) case 2 : 1st partial overlap

Subcase 1b:



$$\text{at } -2 \leq x \leq -1$$

$$g(x) = \int_{-n}^{x+n} f(t) \cdot h(x-t) dt$$

$$= \int_{-n}^{x+n} (1+t) \cdot 1 dt$$

$$= \int_{-n}^{x+n} 1 dt + \int_{-n}^{x+n} t dt$$


$$= \left[t \right]_{-n}^{x+n} + \left[\frac{t^2}{2} \right]_{-n}^{x+n}$$

$$= x+n - (-1) + \frac{(x+n)^2}{2} - \frac{(-1)^2}{2}$$

$$= x+2 + \frac{1}{2}x^2 + \frac{2}{2}x + \frac{1}{2} - \frac{1}{2}$$

$$g(x) = \frac{1}{2}x^2 + 2x + 2$$

subcase 2b: 1st partial overlap



$$g(x) = \int_{-n}^0 f(t) \cdot h(x-t) dt + \int_0^{x+n} f(t) \cdot h(x-t) dt$$

$$= \int_{-n}^0 (1+t) \cdot 1 dt + \int_0^{x+n} (1-t) \cdot 1 dt$$

$$= \int_{-n}^0 1 dt + \int_{-n}^0 t dt + \int_0^{x+n} 1 dt - \int_0^{x+n} t dt$$

$$g(x) = [t]_{-1}^0 + \left[\frac{t^2}{2}\right]_{-1}^0 + [t]_0^{x+1} - \left[\frac{t^2}{2}\right]_0^{x+1}$$


$$= 0 - (-1) + \frac{0^2}{2} - \frac{(-1)^2}{2} + x+1 - 0 - \frac{(x+1)^2}{2} - \frac{0^2}{2}$$

$$= 1 - \frac{1}{2} + x+1 - \frac{x^2}{2} - \frac{2x}{2} - \frac{1}{2}$$

$$g(x) = 1 - \frac{1}{2}x^2$$

when $-1 \leq x \leq 0$

case 3: Full overlap



we have a full overlap at $x=0$, so $g(x)$ is

$$g(x) = \int_{-1}^0 f(t) \cdot h(x-t) dt + \int_0^1 f(t) \cdot h(x-t) dt$$

$$= \int_{-1}^0 (1+t) \cdot 1 dt + \int_0^1 (1-t) \cdot 1 dt$$


$$= \int_{-1}^0 1 dt + \int_{-1}^0 t dt + \int_0^1 1 dt - \int_0^1 t dt$$

$$= [t]_{-1}^0 + \left[\frac{t^2}{2}\right]_{-1}^0 + [t]_0^1 - \left[\frac{t^2}{2}\right]_0^1$$

$$\begin{aligned}
 g(x) &= 0 - (-1) + \frac{0^2}{2} - \frac{(-1)^2}{2} + 1 - 0 - \frac{1^2}{2} + \frac{0^2}{2} \\
 &= 1 - \frac{1}{2} + 1 - \frac{1}{2} \\
 &= 2 - 1 = 1
 \end{aligned}$$

$$\boxed{\text{at } x=0 ; g(x) = 1}$$

d) case 4: 2nd partial overlap
subcase 4a)




$$\begin{aligned}
 g(x) &= \int_{x-1}^0 f(t) \cdot h(x-t) dt + \int_0^1 f(t) \cdot h(x-t) dt \\
 &= \int_{x-1}^0 (1+t) \cdot 1 dt + \int_0^1 (1-t) \cdot 1 dt \\
 &= \int_{x-1}^0 1 dt + \int_{x-1}^0 t dt + \int_0^1 1 dt - \int_0^1 t dt \\
 &= [t]_{x-1}^0 + \left[\frac{t^2}{2}\right]_{x-1}^0 + [t]_0^1 - \left[\frac{t^2}{2}\right]_0^1 \\
 &= 0 - (x-1) + \frac{0^2}{2} - \frac{(x-1)^2}{2} + 1 - 0 - \frac{1^2}{2} + \frac{0^2}{2}
 \end{aligned}$$

$$\begin{aligned}
 g(x) &= -x+1 - \left(\frac{x^2 - 2x + 1}{2} \right) + 1 - \frac{1}{2} \\
 &= -x+1 - \frac{1}{2}x^2 + \frac{2}{2}x - \frac{1}{2} + 1 - \frac{1}{2}
 \end{aligned}$$

$$g(x) = -\frac{1}{2}x^2 + 1 \quad \text{when} \quad 0 < x < 1$$

Case (b):



$$g(x) = \int_{x-1}^1 f(t) \cdot h(x-t) dt$$

$$= \int_{x-1}^1 (1-t) \cdot 1 dt$$

$$= \int_{x-1}^1 1 dt - \int_{x-1}^1 t dt$$

$$= \left[t \right]_{x-1}^1 - \left[\frac{t^2}{2} \right]_{x-1}^1$$

$$= 1 - (x-1) - \frac{1^2}{2} + \frac{(x-1)^2}{2}$$

$$g(x) = 1 - x + 1 - \frac{1}{2} + \frac{(x^2 - 2x + 1)}{2}$$

$$= 1 - x + 1 - \frac{1}{2} + \frac{1}{2}x^2 - x + \frac{1}{2}$$

$$= \frac{1}{2}x^2 - 2x + 2$$

$$g(x) = \frac{1}{2}x^2 - 2x + 2 \quad \text{when } 1 < x < 2$$

so

$$g(x) = \begin{cases} 0 & ; x < -2 \\ \frac{1}{2}x^2 + 2x + 2 & ; -2 < x < -1 \\ 1 - \frac{1}{2}x^2 & ; -1 < x < 1 \\ \frac{1}{2}x^2 - 2x + 2 & ; 1 < x < 2 \\ 0 & ; x > 2 \end{cases}$$

