

1  
2 Stable isotopes contain substantial additive information about terrestrial  
3 carbon and water cycling.

4  
5 Bonan Li<sup>a,b,c,d</sup> Stephen P. Good<sup>a,b</sup>, Richard P. Fiorella<sup>e,f,g</sup>, Catherine E. Finkenbiner<sup>a,b</sup>,  
6 Gabriel J. Bowen<sup>e,f</sup>, David C. Noone<sup>h</sup>, Christopher J. Stilli<sup>i</sup>, William R.L. Anderegg<sup>f,j,h</sup>

7 <sup>a</sup>Department of Biological & Ecological Engineering, Oregon State University

8 <sup>b</sup>Water Resources Graduate Program, Oregon State University

9 <sup>c</sup>College of Earth Ocean and Atmospheric Sciences, Oregon State University

10 <sup>d</sup>Department of Biological & Agricultural Engineering, University of Arkansas

11 <sup>e</sup>Department of Geology and Geophysics, University of Utah

12 <sup>f</sup>Global Change and Sustainability Center, University of Utah

13 <sup>g</sup>Earth and Environmental Sciences Division, Los Alamos National Laboratory

14 <sup>h</sup>Department of Physics, University of Auckland

15 <sup>i</sup>Department of Forest Ecosystems and Society, Oregon State University

16 <sup>j</sup>School of Biological Sciences, University of Utah

17 <sup>h</sup>Wilkes Center for Climate Science and Policy, University of Utah

18 Corresponding author: Bonan Li

19 [Email:libon@oregonstate.edu](mailto:libon@oregonstate.edu)

## Abstract

Stable isotope ratios of H ( $\delta^2H$ ), O ( $\delta^{18}O$ ), and C ( $\delta^{13}C$ ) are linked to key biogeochemical processes of the water and carbon cycles; however, the degree to which isotope-associated processes are reflected in macroscale ecosystem flux observations remains unquantified. Here through formal information assessment, new measurements of  $\delta^{13}C$  of net ecosystem exchange ( $NEE$ ) as well as  $\delta^2H$  and  $\delta^{18}O$  of latent heat ( $LH$ ) fluxes across the United States National Ecological Observation Network are used to determine conditions under which isotope measurements are informative of environmental exchanges. We find all three isotopic datasets individually contain comparable amounts of information about  $NEE$  and  $LH$  fluxes as wind speed observations. Such information from isotope measurements, however, is largely unique. Generally,  $\delta^{13}C$  provides more information about  $LH$  as aridity increases or mean annual precipitation decreases;  $\delta^2H$  provides more information about  $LH$  as temperatures or mean annual precipitation decreases, and also provides more information about  $NEE$  as temperatures decrease. Overall, we show that the stable isotope datasets collected by NEON contribute non-trivial amounts of new information about bulk environmental fluxes useful for interpreting biogeochemical and ecohydrological processes at landscape scales. However, the utility of this new information varies with environmental conditions at continental scales. This study provides an approach for quantifying the value adding non-traditional sensing approaches to environmental monitoring sites and These results demonstrate isotopic variability reflecting biophysical controls on  $LH$ , and to a lesser extent  $NEE$ , fluxes is stronger under low precipitation, arid, and cooler conditions. The patterns identified here in this study are expected to aid in modeling and data interpretation efforts focused on constraining carbon and water cycles' mechanisms.

**Keywords:** isotope, carbon flux, water flux, NEON, information theory

## 1. Introduction

Understanding the interactions and drivers of water and carbon exchanges between terrestrial ecosystems and the atmosphere is crucial to illuminate processes driving Earth's current climate as well as forecasting impacts of future change on ecosystems and the climate itself (Jung et al 2011, Piao et al 2020)[1,2]. To date, significant efforts have been made to monitor terrestrial carbon and water fluxes, including the widespread development of macroscale eddy covariance (EC) networks to measure ecosystem fluxes (Baldocchi 2014, Schimel and Schneider 2019)[3,4]. EC flux towers can measure continuous net ecosystem exchange ( $NEE$ ) of  $CO_2$  between the land surface and atmosphere at various frequency-time domains. Similarly, EC measurements of latent heat flux ( $LH$ ), representing evaporation and transpiration from soils, water bodies, and plant canopies, provides valuable information for understanding regional and global water

budgets as well as agricultural applications (Zhou et al 2018, Zeng et al 2020)[5,6]. Flux measurements have been used for a variety of environmental applications such as calibrating and validating remotely sensed flux estimations (Jia et al 2012)[7], parameterizing land surface models (Williams et al 2009)[8], modeling seasonal crop coefficients (Li et al 2008)[9], and investigating disturbance impacts such as post-fire carbon balance (Lupascu et al 2020)[10]. While measurements of  $LH$  and  $NEE$  can quantify fluxes themselves, new kinds of data are needed to refine knowledge of the processes driving these fluxes which are central to the carbon and water cycles.

To improve understanding of Earth system processes, the geoscience community has developed a wide array of advanced measurements to complement EC flux data to help constrain environmental processes. These include studies focused on stable isotope fluxes (Dubbert and Werner 2019)[11], Carbonyl Sulfide (COS) (Whelan et al 2018)[12], various radiometric indices such as thermal (Still et al 2021)[13] and solar induced fluorescence (SIF) (Guan et al 2016)[14], and even environmental DNA (URYcki et al 2022)[15]. Prominent among these techniques, naturally occurring water and carbon isotopes measurements have been shown to be a powerful tool for understanding a wide array of ecohydrological and biophysical processes because distinct processes are, and are not, often associated with known isotope transformations (i.e., fractionation effects) (Bowen and Good 2015)[16]. Water isotope ratios ( $\delta^2H$  and  $\delta^{18}O$  in water) have been used to partition evapotranspiration into evaporation and transpiration, as evaporated and transpired fluxes from the same ecosystem may have distinct isotope ratios (Xiao et al 2018, Berkelhammer et al 2013)[17,18].  $\delta^{13}C$  values of  $CO_2$  have also been applied to separate  $NEE$  into its constituent fluxes, as the isotopic composition of photosynthesis can differ from that of ecosystem respiration (Lee et al 2020)[19]. Previous network-based studies of  $\delta^2H$ ,  $\delta^{18}O$  and  $\delta^{13}C$  examined patterns across distinct ecosystems using cryogenic baths and flask samples; however, the poor temporal sampling and spatial coverage has limited these approaches to understand ecosystem-scale processes these approaches are constrained in their ability to provide information about ecosystem-scale processes, which generally requires finer temporal and spatial sampling coverage (Orlowski et al 2018, Gemery et al 1996)[20,21]. The development of automated laser spectroscopy systems mounted on EC towers provides new opportunities to obtain long term spatially and temporally resolved atmosphere profiles of these isotopes (Fiorella et al 2021)[22]. The recently launched National Ecological Observatory Network (NEON) provides the first standardized measurements of the stable isotope ratios of  $H_2O$  vapor and  $CO_2$  for ecosystems across the USA that can be used to estimate  $\delta^2H$  and  $\delta^{18}O$  of  $LH$  and  $\delta^{13}C$  of  $NEE$  (Finkenbiner et al 2022)[23].

The development of advanced ecosystem measurements across networks such as NEON presents new scientific possibilities; yet this also raises the fundamental question of how useful new and often expensive data streams are for constraining targeted environmental processes. Many advanced measurements are made at considerable cost and effort, yet their full value as a source of information beyond traditional meteorological observations (e.g., vapor pressure deficit, [VPD], air temperature, [T], global radiation,

$\{R_g\}$ , and windspeed  $\{u\}$ , is rarely demonstrated in a formal sense, especially within continental-scale networks where variability in environmental conditions occurs across a much wider range than individual sites. Here we capitalize on recent advances in information theory to assess the information content of NEON stable isotope data. These advances allow for the formal quantification of linear and nonlinear interactions between variables (termed mutual information) (Cover and Thomas 2005)[24], as well as approaches to diagnose how unique the information provided by new data sources is relative to others (Goodwell and Kumar 2017, Williams and Beer 2010)[25,26]. This study addresses three related questions: (1) Do new observations (here  $\delta^2H$ ,  $\delta^{18}O$ , and  $\delta^{13}C$  values) contain useful information about the bulk *NEE* and *LH* fluxes across North America? (2) Can any of the information provided by new (isotope) measurements be obtained from other meteorological variables? And (3) Under which environmental conditions do these new measurements provide the most additional information? In doing so, this study provides a generalizable approach for evaluating the conditions under which novel geoscience data is helpful for understanding the Earth system. It also formally quantifies the conditions under which environmental processes associated with transformations of stable isotope ratios, as measured systematically within continental scale networks, are a greater contribution to overall environmental exchanges. This approach thereby provides key process level benchmarks for advancing research into Earth's integrated carbon and water cycles.

## 2 Materials and methods

### 2.1 Study sites and data preparations

This study was conducted ~~at terrestrial sites that are part of~~ at part of terrestrial sites of National Ecological Observatory Network (NEON), which is a continental scale research platform for understanding the ecological responses to climate change, land use change and species invasion (Barnett et al 2019)[27]. We used the 30-minute aggregated *NEE*, *LH*, global radiation ( $R_g$ ), air temperature ( $T$ ), and the two-dimensional wind speed ( $u$ ) datasets from the NEON's eddy covariance bundled datasets (National Ecological Observatory Network (NEON) 2022a)[28]. The vapor pressure deficit (VPD) data were derived based on NEON's relative humidity, temperature, and barometric pressure products (NEON 2022b). The *NEE* and *LH* data were filtered for periods of low turbulence based on friction velocities ( $u^*$ ) then gap-filled using the marginal distribution sampling method (Wutzler et al 2018). The gap-filled and  $u^*$  filtered 30-minute fluxes were further averaged on a daily scale to facilitate future analysis. All 30-minute meteorological variables except wind speed were gap-filled using the marginal distribution sampling mentioned above. The gap-filled meteorological variables were then averaged to daily scale. Extreme values in daily flux and meteorological datasets were further processed using an inter-quantile filter (Goodwell and Kumar 2017). More details can be found in Supplemental information. The vapor pressure deficit (VPD) data were derived based on NEON's relative humidity product [29]. These 30-minute variables were gap-filling and

~~further processed to daily scale. More details can be found in Supplemental information.~~  
Daily stable isotope ratios of *NEE* and *LH* were obtained from a recently published  
datasets (Finkenbiner et al 2022)[30], ~~which was derived based on the isotope~~  
~~composition of carbon dioxide and water vapor from EC tower profiles across NEON sites.~~  
~~The  $\delta^{18}\text{O}$  values were converted to deuterium excess (*d*) via  $d = \delta^2\text{H} - 8 * \delta^{18}\text{O}$~~   
~~(Dansgaard 1964).~~  
~~which was derived based on the surface isotope composition of carbon dioxide and water~~  
~~vapor across NEON sites.~~

## 2.2 Information measures

~~In this study the mutual information metric,  $I(X;Y)$ , was chosen to analyze how different~~  
~~meteorological variables share information about ecosystem fluxes because it has the~~  
~~advantage over traditional metric (e.g., correlation coefficient) of capturing both linear and~~  
~~non-linear dependencies between two variables. It represents the reduction in~~  
~~uncertainties of one variable given the knowledge of another variable. Formally, Mutual~~  
information is a measure of how two random variables are probabilistically dependent on  
each other in the unit of bits (Cover and Thomas 2005)[24]. Probabilistically, the mutual  
information can be expressed as:

$$I(X;Y) = \sum p(x,y) \log_2 \left( \frac{p(x,y)}{p(x)p(y)} \right) \quad (1)$$

where  $p(x)$ ,  $p(y)$ , and  $p(x,y)$  are the probability density functions of random variables  $X$ ,  $Y$ ,  
and  $\{X,Y\}$  respectively.

The multivariate mutual information of a single random variable ( $Z$ ) and a set of random  
variables  $\{X, Y\}$  characterizes the amount of uncertainty in  $Z$  that can be reduced by the  
knowledge of  $\{X, Y\}$  and can be expressed as:

$$I(X,Y;Z) = \sum p(x,y,z) \log_2 \left( \frac{p(x,y,z)}{p(x,y)p(z)} \right) \quad (2)$$

where  $p(z)$ ,  $p(x,y)$ , and  $p(x,y,z)$  are the probability density functions of variables  $Z$ ,  $\{X,Y\}$ ,  
and  $\{X,Y,Z\}$ , respectively and were estimated using a kernel density estimation (KDE)  
method with a gaussian kernel and Silverman bandwidth selection method (Silverman  
2018)[34]. ~~To evaluate the above information metrics, We-rescaledwe rescaled~~ each data  
point to a common range of [0, 1] before using KDE. We then evaluate the probability  
density functions from 0 to 1 with a step size of 0.05.

We computed the pairwise mutual information (~~[e.g.,  $I(NEE;VPD)$ ,  $I(LH;VPD)$ , ...etc]~~)  
shared among  $VPD$ ,  $T$ ,  $R_g$ ,  $u$ ,  $\delta^{13}\text{C}$ ,  $\delta^2\text{H}$ , and  $d$  about *NEE* and *LH* iteratively. Due to the  
limitation of isotope datasets, we computed the mutual information of each variable with  
the *NEE* and *LH* by subsampling 100 data points without replacement 500 times to ensure  
constituent data counts in mutual information calculations. Then, the mutual information  
of the variable of interest and the flux is computed as the average mutual information  
across 500 resamplings. The mutual information contents computed above are  
~~testedevaluated~~ for statistical significance (refer to Supplemental information for details).



## 2.3 Partial information decomposition

The multivariate mutual information can be decomposed into different informational components via a partial information decomposition framework (PID) (Goodwell and Kumar 2017a, Goodwell et al 2018, Williams and Beer 2010)[25,26,32]. This framework captures how the different source variables interactively influence a target variable of interest, which can possibly reveal the process that relates source variables and a target without any modeling assumptions. The PID can decompose  $I(X,Y;Z)$  into: (1) unique information (U) that is only provided by X or Y solely to the Z; (2) synergistic information (S) that is the information provided to the Z when X and Y act jointly; (3) redundant information (R) that is the overlapping information provided both by X and Y to the Z (Goodwell and Kumar 2017-[33]). The PID framework can be formulated as

$$I(X,Y; Z) = U_X + U_Y + R + S \quad (3)$$

$$I(X; Z) = U_X + R \quad (4)$$

$$I(Y; Z) = U_Y + R \quad (5)$$

Where  $U_X$  and  $U_Y$  are the unique information of X and Y to Z, respectively. R and S are the redundant and synergistic information of X and Y to Z, respectively. All PID components are non-negative real numbers in unit of bits (Goodwell and Kumar 2017)[33].

In this study, we quantified the information flow between each flux and each isotope flux ratio by leveraging the PID framework (Goodwell and Kumar 2017)[25]. We defined the decomposed information components that the isotope ratios provided to the bulk fluxes as the averaged unique information across all meteorological variables ( $VPD$ ,  $T$ ,  $R_g$ , and  $u$ ). As with computing the individual mutual information, we also subsampled 100 data points from each dataset without replacement 500 times. The partial information components of the isotopes were then computed as the averaged information components from 500 iterations. The significance tests were performed similarly to mutual information (refer to Supplemental information for details).

## 3. Results

Informational analysis shows that isotope data ( $\delta^{13}C$ ,  $\delta^2H$ , and  $d$ ) and traditional meteorological data ( $R_g$ ,  $T$ ,  $VPD$ ,  $u$ ) each contain significant information about temporal variation in  $NEE$  and  $LH$  fluxes (Fig. 1) ~~throughout the NEON sites. This formally demonstrates that  $NEE$  and  $LH$  become less uncertain given the knowledge of isotope data or meteorological data throughout the NEON sites.~~ We find that  $R_g$ ,  $T$ , and  $VPD$  observations consistently contain more information about environmental fluxes than either isotope data or wind speed ( $u$ ), which provides comparable amount information about  $NEE$  and  $LH$  fluxes (Fig. 1). Though the information provided by  $R_g$  is larger than the information from  $u$  and the isotopes,  $u$  is nevertheless one of the well-established drivers of surface-atmosphere water and carbon exchange and is commonly measured at meteorological stations worldwide (Yusup and Liu 2020)[34].

In general, individual variables tend to share more information with  $LH$  than  $NEE$  (Fig. 1). This indicates that  $LH$  is generally more easily constrained and predicted based on these environmental observations, possibly because it more strongly captures isotopic differences in the contributing one-way flux compared to  $NEE$  which is the net sum of two opposing fluxes with less distinct isotope ratios. Instead of  $\delta^{13}C$  values best constraining  $NEE$  and  $\delta^2H$  or  $d$  values best constraining  $LH$ , we find that  $\delta^2H$  values on average provide slightly more mutual information than  $\delta^{13}C$  values for both  $NEE$  and  $LH$  fluxes; however, both these (i.e.,  $\delta^2H$  and  $\delta^{13}C$ ) are more informative than  $d$ .  $NEE$  is a quantity that encompasses downward carbon uptake via plant photosynthesis and carbon release upward through respiration (Reichstein *et al* 2005) while water, as represented by  $LH$ , is evaporated upward during evapotranspiration.  $\delta^2H$  links with the phase transformation of water that is strongly temperature dependent (Xiao *et al* 2018). Therefore,  $\delta^2H$  is more likely to carry slightly more information about  $LE$  and  $NEE$  than  $\delta^{13}C$ . The amount of information that can be inferred from isotopes (and other variables) about  $NEE$  and  $LH$  is highly unlikely to be obtained by random processes ( $p < 0.01$ ).

We decomposed and evaluated the multivariate mutual information between environmental fluxes, isotope ratios, and other variables (Fig. 2). These results demonstrate that most of the information provided by the isotopes about  $NEE$  and  $LH$  is unique to these measurements ( $\delta^{13}C$  and  $\delta^2H$ ). This unique information (i.e., the information contribution that is contributed only by one variable to the target variable) provided by  $\delta^{13}C$  and  $\delta^2H$  values about  $LH$  is generally higher than the unique information provided about  $NEE$ . The unique information provided by  $\delta^{13}C$  and  $\delta^2H$  values is higher than that contained within  $d$  values for both  $LH$  and  $NEE$  fluxes. The unique information is found to vary spatially across the NEON sites (Supplemental Fig. S1). All the unique information provided by the isotope ratios is statistically significant and highly unlikely to be obtained at random ( $p < 0.01$ ).

In addition to the unique information that  $\delta^{13}C$ ,  $\delta^2H$ , and  $d$  values contain about  $NEE$  and  $LH$  fluxes, a smaller amount of synergistic (i.e., the information component when isotope and other variables act jointly to provide information about ecosystem fluxes) –and redundant information (i.e., the overlapping information that isotope or other variables contribute to ecosystem fluxes) is also presented (Supplemental Fig. S2 and S3). Among all the isotopes, the synergistic component of  $d$  values is slightly larger for  $NEE$  and  $\delta^{13}C$  is marginally larger for  $LH$ . In general, redundant information tends to be smaller than the unique and synergistic components. The unique and redundant information linking isotopes with  $NEE$  and  $LH$  are statistically significant ( $p < 0.01$ ).

The total additional information, represented by the sum of the synergistic information and the unique information ( $U+S$ ), provided by each flux isotope composition to  $LH$  and  $NEE$  varies spatially across NEON sites (Fig. 3)  $\delta^{13}C$  contributes the most substantial information about  $NEE$  and  $LH$  in the Northeastern US (i.e., New Hampshire) and southwestern US (i.e., New Mexico), respectively (Fig. 3a and Fig. 3d). In northern Alaska,  $\delta^2H$  contributes the largest amount of additive information to  $NEE$  (Fig. 3b). There is an

increased possibility of observing more additive information of  $\delta^2H$  about  $LH$  at site with higher latitude (Fig. 3e). The highest additional information that  $d$  provides to  $NEE$  and  $LH$  were observed in Virginia (Fig. 3c) and Wyoming (Fig. 3f), respectively. The fraction of information for isotopes about  $NEE$  that is additive, i.e.  $(U+S)/(U+S+R)$ , is 0.95 for  $\delta^{13}C$ , 0.92 for  $\delta^2H$ , and 0.99 for  $d$ , respectively). For  $LH$ ,  $\delta^2H$  and  $\delta^{13}C$  provided more additive information than  $d$  (Fig. 3aA). The fraction of additive information about  $LH$  is 0.89 for  $\delta^{13}C$ , 0.84 for  $\delta^2H$ , and 0.94 for  $d$ , respectively. The additive information of  $\delta^{13}C$ ,  $\delta^2H$  and  $d$  relating to  $LH$  has larger variability among sites than that relating to  $NEE$  (Fig. 3a-3b and Fig. 3d-3e3), and there is less variability in the additive information of  $d$  about  $NEE$  (Fig. 3c and Fig. 3f) than in  $LH$ . All the additive information of these isotopes relating to  $NEE$  and  $LH$  is statistically significant ( $p < 0.01$ ).

## 4. Discussion

Our analysis provides a rigorous evaluation of the quantitative value of isotope ratios to provide useful information about carbon and water fluxes across continental scale gradients. For these bulk fluxes, we showed that the information individually provided by these isotopes was similar to the amount of information provided by wind speed measurements, while providing less information than atmospheric vapor pressure deficit, air temperature, and radiation measurements. The meteorological observations evaluated here are commonly used to drive forecasts of environmental processes (Cosgrove et al 2003, Rodell et al 2004)[35,36] and thus serve as a benchmark for environmental data. A prior  $NEE$  simulation showed that radiation was consistently the most sensitive predictor for the simulation of  $NEE$  at ~~maze~~-maize fields with distinct irrigation practices (Safa et al 2019)[37]. Similarly, a sensitivity analysis on global evapotranspiration models indicated that net radiation was ~~found to be~~ one of the influential input variables (Talsma et al 2018)[38]. Our results are consistent with the fundamental notion that solar radiation is the basis for all ecosystem functions (Yetemen et al 2015)[39] (excluding rare energy transformations) and drives most diurnal variation in air temperature and vapor pressure deficit and therefore is more likely to share higher amount of mutual information individually with  $LH$  and  $NEE$ , with temperature and moisture levels of secondary importance and isotope metrics and wind speed of tertiary importance.

The meteorological variables evaluated here are known to be inter-related to some extent. For instance, the vapor pressure deficit is strongly dependent on air temperature due to the Clausius-Clapeyron relationship (Clausius 1850)[40] and air temperature is tightly related to the amount of radiation as well as to sensible heat fluxes. Past studies have highlighted how  $NEE$  and  $LH$  respond to changes in vapor pressure deficit, air temperature and radiation across various scales, seasons, and ecosystems (Chen et al 2020, Niu et al 2012, Gu et al 2006)[41–43]. Vapor pressure deficit was found to have direct effect on surface energy partitioning as high vapor pressure deficit represents high atmosphere demand and hence high  $LH$  with constant surface conductance (Tong et al 2022, Wang et al 2019)[44,45]. Yet, high vapor pressure deficit can reduce stomatal



conductance and thereby reduce plant photosynthesis (Grossiord *et al* 2020)[46]. Wind speed can modulate the rate of evapotranspiration and thereby *LH* (Yang *et al* 2019, Liu and Zhang 2013)[47,48]. The different effect of vapor pressure deficit and wind speed on *LH* may be underrepresented by other metrics but can be captured if evaluated using information theory-based metrics like those explored here.

Information carried by these isotope ratios was found to be distinct from the traditional meteorological variables examined here~~unique~~. It is also crucial to understand how different variables interactively provide information to a target of interest because knowledge of the interdependencies ~~interactive dependencies~~ between the inputs and outputs of a studied system is fundamental for model uncertainty characterization (Ruddell *et al* 2019, Li and Good 2021, Gong *et al* 2013)[49–51]. In fact, one of the challenges for land surface models is increasing process complexity with the integration of a set of sub-models with the expansion of input dimensions (Fisher and Koven 2020)[52], which can increase the risk of model “equifinality”. Moreover, numerous models have been developed to estimate ecosystem fluxes (Wood 2021, Su 2002, Veroustraete *et al* 1996)[53–55].—However, these methods often require some assumptions or simplifications, which can be subject to significant uncertainty (Papale *et al* 2006, Zhao *et al* 2020)[56,57]. In general, it may be more desirable for each of most of the inputs in a model to provide unique or synergistic pieces of information (Wibral *et al* 2017)[58], which can potentially capture different processes relating to the target (Goodwell *et al* 2018)[32]. Therefore, the construction and simplification of ecosystem models should move ~~be~~ towards a direction that maximizes unique information of each input.

The decomposition of the multivariate mutual information between isotopes, other meteorological variables, and the bulk fluxes offers an opportunity to elucidate how much of the information from isotopes is transferred to the bulk fluxes (*NEE* and *LH*). In this study, the portion of unique information from isotopes measurements for carbon and water isotopes was statistically significant. This suggests that processes driving variation in isotope ratios may influence these fluxes via distinct pathways. ~~suggesting that isotope ratios of the fluxes may influence these fluxes via distinct pathways.~~ We observed inter-site variations in the unique information provided by the isotopes, indicating that the unique information may be dependent on site-specific conditions ~~(e.g., such as aridity and precipitation).~~ There is a higher chance that  $\delta^{13}\text{C}$  contributes more information about *LH* under drier or lower precipitation conditions (Supplemental Fig. S4). Additionally, both  $\delta^{13}\text{C}$  and  $\delta^2\text{H}$  tend to provide more distinct insights into *NEE* and *LH* in cooler or lower precipitation conditions.—This suggests that the patterns of bulk fluxes can potentially be better characterized and predicted with the isotopes included as an additional constraint.

The additional information provided by isotopes to these bulk fluxes are described by the sum of unique information and synergistic information. Our analysis demonstrate that fusing isotope data products can potentially lead to better monitoring and prediction of *NEE* and *LH* in a process modeling framework, as these isotope datasets provide

additional information beyond traditional meteorological variables and are associated with known physical mechanisms. The incorporation of isotope datasets into artificial intelligence (AI) and machine learning (ML) models, especially explainable AI models, can potentially improve predictive accuracies and enhance our understanding of ecosystem fluxes. Nevertheless, uncertainties can be introduced when incorporating isotope dataset to models with larger spatial scale. It is challenging to include isotope datasets to models that require larger spatial scale isotope datasets, as they are often hard to acquire. Researchers might also consider different incorporation strategies in different ecoclimate regions. HoweverIn addition, the amount of added information of the isotope datasets is likely to vary across sites, climate, and ecosystems. To assess this, we evaluated the additive information of isotopes based on NEON site conditions via a simple linear regression analysis (Fig. 4). We showed that the additive information that  $\delta^{13}\text{C}$  provides about  $LH$  is influenced by mean annual precipitation, aridity, and site elevation (Fig. 4d), as indicated by a significant slope value from the linear regression.  $\delta^{13}\text{C}$  is likely to provide more useful information about  $LH$  in locations with higher atmospheric evaporative demand relative to precipitation or in locations with less annual precipitation or with higher altitude. The additive information  $\delta^2\text{H}$  provides about  $NEE$  was shown to be mainly influenced by the site mean annual temperature (Fig. 4b).  $\delta^2\text{H}$  tends to be more informative about  $NEE$  in locations with cooler climates. Similarly, there is more opportunity for  $\delta^2\text{H}$  to provide additional knowledge about  $LH$  at locations with cooler climates or less mean annual precipitation (Fig. 4e). No significant relationship was found between the additional information of the  $d$  provided to either  $NEE$  or  $LH$ .

Variations in additional information across NEON sites indicate differences in conditional dependencies of ecosystem fluxes on processes related to isotope fluxes. Changes in ecosystem structure and climate affect the ecosystem's adaptability to environmental changes (Weiskopf et al 2020) that influences the biochemical processes responsible for isotope fractionation, which can intensify or weaken these conditional dependencies. It's worth noting that this study primarily examined how each isotope contributes additional information to  $NEE$  and  $LH$  with an emphasis on atmospheric centric conditions. However, it is crucial to acknowledge that ecosystems broadly have a wide range of inherent complexities, such as geomorphology (e.g., slope, aspect) subsurface dynamics (e.g., depth to water table), vegetation species and traits (e.g. plant hydraulic traits), and soil physics (e.g., soil texture), which might play a role in shaping the way of how isotope observations provide extra information about  $NEE$  and  $LH$ .

One of key motivations for measuring stable isotopes of water and carbon fluxes is that they may provide unique and novel knowledge about key mechanisms across ecosystems (Good et al 2014, Conrad et al 2012, Wang et al 2010). Such hypothesis has not been formally tested until this study. One of key motivations for measuring stable isotopes of water and carbon fluxes is that they may provide a unique constraint across ecosystems, which has not been formally tested until this study, which allow for the partitioning of bulk fluxes into their respective constituents (Good et al 2014, Conrad et al 2012, Wang et al 2010)[59–64]. This is because the flux isotope ratios are influenced by

distinct biophysical processes, and thus larger amounts of new mutual information between isotopes and environmental fluxes quantifies the conditions under which these processes are more dominant ~~dominate~~ components of overall bulk fluxes. In this light, the trends described above (and in Fig 4) are consistent with prior knowledge of isotope geophysics. For instance, equilibrium fractionation factors are sensitive to temperature, particularly at low values (Bowen and Good 2015)[46], with broad decreases in vapor  $\delta^2H$  observed poleward at continental scales (Good et al 2015)[62]. Similarly, evaporation is expected to play a larger role in *LH* fluxes under low vegetation, more arid climates (Wang et al 2014)[63], and this study provides a new way to quantify the relative importance of these isotope processes on bulk fluxes.

It is important to acknowledge that our analysis focused on how daily isotope datasets are informative of bulk ecosystem fluxes. It might be worthwhile to analyze how similar observations are informative of ecosystem fluxes at finer temporal scales. For instance, how lags in isotope dataset responses are influenced ecosystem processes, and correspondingly how do the partial information components change with different lag timescales can possibly reveal more detailed linkages between ecosystem fluxes and isotope fluxes. In this study, we considered abiotic variables ( $VPD$ ,  $T$ ,  $u$ ,  $R_g$ ) as the confounding part in the partial information decomposition. It might also be worthwhile to explore how other biotic variables such as ecosystem structure, species composition, and plant hydraulic traits, rooting depth can influence the total additive information of isotope dataset to the bulk fluxes.

This analysis is based on current available data products and quality control methods. As more NEON data becomes available, future studies to may investigate if and how the results vary with longer timeseries data and a wider range of environmental conditions. It may be also worthwhile for future studies to investigate if and how the results vary with more available datasets and a wider range of environmental conditions. However, given the power of isotopes for tackling fundamental problems in carbon and water cycling and projecting the future of terrestrial ecosystem function under a rapidly changing climate (Bowen and Good 2015, Bowling et al 2008)[46,64], our results can be useful to provide guidance for improving model results after the incorporation of isotope flux ratios.

## Acknowledgements

The authors want to acknowledge the funding support of the United States National Science Foundation (DEB1802885 and DEB1802880). RPF also received support from the Laboratory Directed Research and Development program of Los Alamos National Laboratory under project number 20210961PRD3.

## Author contribution

BL and SPG designed the study. RPF provided flux datasets and gap-filled meteorological datasets and wrote part of the data processing steps in Supplementary material. BL analyzed the data and wrote the manuscript. SPG, RPF, CEF, GJB, DCN, CJS, and WRLA reviewed the manuscript.

## Data availability statement

The datasets that are associated with this study is publicly available at <https://data.neonscience.org/> and <https://www.hydroshare.org/resource/e74edc35d45441579d51286ea01b519f/>. All materials associated with this study will be made available at [https://github.com/libonancaesar/ERL\\_info\\_isotope](https://github.com/libonancaesar/ERL_info_isotope)

## Competing interest statement

The authors declare no conflicts of interest.

## References

- Baldocchi D 2014 Measuring fluxes of trace gases and energy between ecosystems and the atmosphere - the state and future of the eddy covariance method *Glob Chang Biol* **20** 3600–9 Online: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.12649>
- Barnett D T, Adler P B, Chemel B R, Duffy P A, Enquist B J, Grace J B, Harrison S, Peet R K, Schimel D S, Stohlgren T J and Vellend M 2019 The plant diversity sampling design for The National Ecological Observatory Network *Ecosphere* **10** Online: <https://onlinelibrary.wiley.com/doi/10.1002/ecs2.2603>

- Berkelhammer M, Hu J, Bailey A, Noone D C, Still C J, Barnard H, Gochis D, Hsiao G S, Rahn T and Turnipseed A 2013 The nocturnal water cycle in an open-canopy forest *Journal of Geophysical Research: Atmospheres* **118** 10,225-10,242 Online: <http://doi.wiley.com/10.1002/jgrd.50701>
- Bowen G J and Good S P 2015 Incorporating water isoscapes in hydrological and water resource investigations *WIREs Water* **2** 107–19 Online: <https://onlinelibrary.wiley.com/doi/10.1002/wat2.1069>
- Bowling D R, Pataki D E and Randerson J T 2008 Carbon isotopes in terrestrial ecosystem pools and CO<sub>2</sub> fluxes *New Phytologist* **178** 24–40 Online: <https://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2007.02342.x>
- Chen J, Wen J, Kang S, Meng X, Tian H, Ma X and Yuan Y 2020 Assessments of the factors controlling latent heat flux and the coupling degree between an alpine wetland and the atmosphere on the Qinghai-Tibetan Plateau in summer *Atmos Res* **240** 104937 Online: <https://linkinghub.elsevier.com/retrieve/pii/S016980951931573X>
- Clausius R 1850 Ueber die bewegende Kraft der Wärme und die Gesetze, welche sich daraus für die Wärmelehre selbst ableiten lassen *Annalen der Physik und Chemie* **155** 500–24 Online: <https://onlinelibrary.wiley.com/doi/10.1002/andp.18501550403>
- Conrad R, Klose M, Yuan Q, Lu Y and Chidthaisong A 2012 Stable carbon isotope fractionation, carbon flux partitioning and priming effects in anoxic soils during methanogenic degradation of straw and soil organic matter *Soil Biol Biochem* **49** 193–9 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0038071712000958>
- Cosgrove B A, Lohmann D, Mitchell K E, Houser P R, Wood E F, Schaake J C, Robock A, Marshall C, Sheffield J, Duan Q, Luo L, Higgins R W, Pinker R T, Tarpley J D and Meng J 2003 Real-time and retrospective forcing in the North American Land Data Assimilation System (NLDAS) project *Journal of Geophysical Research: Atmospheres* **108** 2002JD003118 Online: <https://onlinelibrary.wiley.com/doi/abs/10.1029/2002JD003118>
- Cover T M and Thomas J A 2005 *Elements of Information Theory* (Wiley) Online: <https://onlinelibrary.wiley.com/doi/book/10.1002/047174882X>
- Dansgaard W 1964 Stable isotopes in precipitation *Tellus A: Dynamic Meteorology and Oceanography* **16** 436–68
- Dubbert M and Werner C 2019 Water fluxes mediated by vegetation: emerging isotopic insights at the soil and atmosphere interfaces *New Phytologist* **221** 1754–63 Online: <https://onlinelibrary.wiley.com/doi/10.1111/nph.15547>
- Finkenbiner C E, Li B, Spencer L, Butler Z, Haagsma M, Fiorella R P, Allen S T, Anderegg W, Still C J, Noone D, Bowen G J and Good S P 2022 The NEON Daily



Isotopic Composition of Environmental Exchanges Dataset *Sci Data* **9** 353 Online:  
<https://www.nature.com/articles/s41597-022-01412-4>

Fiorella R P, Good S P, Allen S T, Guo J S, Still C J, Noone D C, Anderegg W R L,  
Florian C R, Luo H, Pingintha-Durden N and Bowen G J 2021 Calibration  
Strategies for Detecting Macroscale Patterns in NEON Atmospheric Carbon  
Isotope Observations *J Geophys Res Biogeosci* **126**

Fisher R A and Koven C D 2020 Perspectives on the Future of Land Surface Models  
and the Challenges of Representing Complex Terrestrial Systems *J Adv Model  
Earth Syst* **12** Online: <https://onlinelibrary.wiley.com/doi/10.1029/2018MS001453>

Gemery P A, Trolier M and White J W C 1996 Oxygen isotope exchange between  
carbon dioxide and water following atmospheric sampling using glass flasks  
*Journal of Geophysical Research: Atmospheres* **101** 14415–20

Gong W, Gupta H V., Yang D, Sricharan K and Hero A O 2013 Estimating epistemic  
and aleatory uncertainties during hydrologic modeling: An information theoretic  
approach *Water Resour Res* **49** 2253–73 Online:  
<http://doi.wiley.com/10.1002/wrcr.20161>

Good S P, Noone D, Kurita N, Benetti M and Bowen G J 2015 D/H isotope ratios in the  
global hydrologic cycle *Geophys Res Lett* **42** 5042–50 Online:  
<http://doi.wiley.com/10.1002/2015GL064117>

Good S P, Soderberg K, Guan K, King E G, Scanlon T M and Caylor K K 2014  $\delta^2\text{H}$   
isotopic flux partitioning of evapotranspiration over a grass field following a water  
pulse and subsequent dry down *Water Resour Res* **50** 1410–32 Online:  
<http://doi.wiley.com/10.1002/2013WR014333>

Goodwell A E and Kumar P 2017 Temporal information partitioning: Characterizing  
synergy, uniqueness, and redundancy in interacting environmental variables *Water  
Resour Res* **53** 5920–42 Online:  
<https://onlinelibrary.wiley.com/doi/10.1002/2016WR020216>

Goodwell A E, Kumar P, Fellows A W and Flerchinger G N 2018 Dynamic process  
connectivity explains ecohydrologic responses to rainfall pulses and drought  
*Proceedings of the National Academy of Sciences* **115** Online:  
<https://pnas.org/doi/full/10.1073/pnas.1800236115>

Grossiord C, Buckley T N, Cernusak L A, Novick K A, Poulter B, Siegwolf R T W, Sperry  
J S and McDowell N G 2020 Plant responses to rising vapor pressure deficit *New  
Phytologist* **226** 1550–66 Online:  
<https://onlinelibrary.wiley.com/doi/10.1111/nph.16485>

Gu L, Meyers T, Pallardy S G, Hanson P J, Yang B, Heuer M, Hosman K P, Riggs J S,  
Sluss D and Wullschlegel S D 2006 Direct and indirect effects of atmospheric  
conditions and soil moisture on surface energy partitioning revealed by a prolonged

drought at a temperate forest site *J Geophys Res* **111** D16102 Online:  
<http://doi.wiley.com/10.1029/2006JD007161>

Guan K, Berry J A, Zhang Y, Joiner J, Guanter L, Badgley G and Lobell D B 2016  
Improving the monitoring of crop productivity using spaceborne solar-induced  
fluorescence *Glob Chang Biol* **22** 716–26 Online:  
<https://onlinelibrary.wiley.com/doi/10.1111/gcb.13136>

Jia Z, Liu S, Xu Z, Chen Y and Zhu M 2012 Validation of remotely sensed  
evapotranspiration over the Hai River Basin, China *Journal of Geophysical  
Research: Atmospheres* **117** n/a-n/a Online:  
<http://doi.wiley.com/10.1029/2011JD017037>

Jung M, Reichstein M, Margolis H A, Cescatti A, Richardson A D, Arain M A, Arneth A,  
Bernhofer C, Bonal D, Chen J, Gianelle D, Gobron N, Kiely G, Kutsch W, Lasslop  
G, Law B E, Lindroth A, Merbold L, Montagnani L, Moors E J, Papale D,  
Sottocornola M, Vaccari F and Williams C 2011 Global patterns of land-atmosphere  
fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy  
covariance, satellite, and meteorological observations *J Geophys Res* **116** G00J07  
Online: <http://doi.wiley.com/10.1029/2010JG001566>

Lee S-C, Christen A, Black T A, Jassal R S, Ketler R and Nesic Z 2020 Partitioning of  
net ecosystem exchange into photosynthesis and respiration using continuous  
stable isotope measurements in a Pacific Northwest Douglas-fir forest ecosystem  
*Agric For Meteorol* **292–293** 108109 Online:  
<https://linkinghub.elsevier.com/retrieve/pii/S0168192320302112>

Li B and Good S P 2021 Information-based uncertainty decomposition in dual-channel  
microwave remote sensing of soil moisture *Hydrol Earth Syst Sci* **25** 5029–45  
Online: <https://hess.copernicus.org/articles/25/5029/2021/>

Li S, Kang S, Li F and Zhang L 2008 Evapotranspiration and crop coefficient of spring  
maize with plastic mulch using eddy covariance in northwest China *Agric Water  
Manag* **95** 1214–22 Online:  
<https://linkinghub.elsevier.com/retrieve/pii/S0378377408001169>

Liu X and Zhang D 2013 Trend analysis of reference evapotranspiration in Northwest  
China: The roles of changing wind speed and surface air temperature *Hydrol  
Process* **27** 3941–8 Online: <https://onlinelibrary.wiley.com/doi/10.1002/hyp.9527>

Lupascu M, Akhtar H, Smith T E L and Sukri R S 2020 Post-fire carbon dynamics in the  
tropical peat swamp forests of Brunei reveal long-term elevated CH<sub>4</sub> flux *Glob  
Chang Biol* **26** 5125–45 Online:  
<https://onlinelibrary.wiley.com/doi/10.1111/gcb.15195>

National Ecological Observatory Network (NEON) 2022a Bundled data products - eddy  
covariance (DP4.00200.001) Online: <https://data.neonscience.org>

National Ecological Observatory Network (NEON) 2022b Relative humidity (DP1.00098.001) Online: <https://data.neonscience.org>

Niu S, Luo Y, Fei S, Yuan W, Schimel D, Law B E, Ammann C, Altaf Arain M, Arneth A, Aubinet M, Barr A, Beringer J, Bernhofer C, Andrew Black T, Buchmann N, Cescatti A, Chen J, Davis K J, Dellwik E, Desai A R, Etzold S, Francois L, Gianelle D, Gielen B, Goldstein A, Groenendijk M, Gu L, Hanan N, Helfter C, Hirano T, Hollinger D Y, Jones M B, Kiely G, Kolb T E, Kutsch W L, Lafleur P, Lawrence D M, Li L, Lindroth A, Litvak M, Loustau D, Lund M, Marek M, Martin T A, Matteucci G, Migliavacca M, Montagnani L, Moors E, William Munger J, Noormets A, Oechel W, Olejnik J, U K T P, Pilegaard K, Rambal S, Raschi A, Scott R L, Seufert G, Spano D, Stoy P, Sutton M A, Varlagin A, Vesala T, Weng E, Wohlfahrt G, Yang B, Zhang Z and Zhou X 2012 Thermal optimality of net ecosystem exchange of carbon dioxide and underlying mechanisms *New Phytologist* **194** 775–83 Online: <https://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2012.04095.x>

Orlowski N, Breuer L, Angeli N, Boeckx P, Brumbt C, Cook C S, Dubbert M, Dyckmans J, Gallagher B, Gralher B, Herbstritt B, Hervé-Fernández P, Hissler C, Koeniger P, Legout A, Macdonald C J, Oyarzún C, Redelstein R, Seidler C, Siegwolf R, Stump C, Thomsen S, Weiler M, Werner C and McDonnell J J 2018 Inter-laboratory comparison of cryogenic water extraction systems for stable isotope analysis of soil water *Hydrol Earth Syst Sci* **22** 3619–37

Papale D, Reichstein M, Aubinet M, Canfora E, Bernhofer C, Kutsch W, Longdoz B, Rambal S, Valentini R, Vesala T and Yakir D 2006 Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation *Biogeosciences* **3** 571–83 Online: <https://bg.copernicus.org/articles/3/571/2006/>

Piao S, Wang X, Wang K, Li X, Bastos A, Canadell J G, Ciais P, Friedlingstein P and Sitch S 2020 Interannual variation of terrestrial carbon cycle: Issues and perspectives *Glob Chang Biol* **26** 300–18 Online: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.14884>

Reichstein M, Falge E, Baldocchi D, Papale D, Aubinet M, Berbigier P, Bernhofer C, Buchmann N, Gilmanov T, Granier A, Grunwald T, Havrankova K, Ilvesniemi H, Janous D, Knohl A, Laurila T, Lohila A, Loustau D, Matteucci G, Meyers T, Miglietta F, Ourcival J-M, Pumpanen J, Rambal S, Rotenberg E, Sanz M, Tenhunen J, Seufert G, Vaccari F, Vesala T, Yakir D and Valentini R 2005 On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm *Glob Chang Biol* **11** 1424–39 Online: <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2486.2005.001002.x>

Rodell M, Houser P R, Jambor U, Gottschalck J, Mitchell K, Meng C-J, Arsenault K, Cosgrove B, Radakovich J, Bosilovich M, Entin J K, Walker J P, Lohmann D and

- Toll D 2004 The Global Land Data Assimilation System *Bull Am Meteorol Soc* **85** 381–94 Online: <https://journals.ametsoc.org/doi/10.1175/BAMS-85-3-381>
- Ruddell B L, Drewry D T and Nearing G S 2019 Information Theory for Model Diagnostics: Structural Error is Indicated by Trade-Off Between Functional and Predictive Performance *Water Resour Res* **55** 6534–54 Online: <https://onlinelibrary.wiley.com/doi/10.1029/2018WR023692>
- Safa B, Arkebauer T J, Zhu Q, Suyker A and Irmak S 2019 Net Ecosystem Exchange (NEE) simulation in maize using artificial neural networks *IFAC Journal of Systems and Control* **7** 100036 Online: <https://linkinghub.elsevier.com/retrieve/pii/S2468601817302584>
- Schimel D and Schneider F D 2019 Flux towers in the sky: global ecology from space *New Phytologist* **224** 570–84 Online: <https://onlinelibrary.wiley.com/doi/10.1111/nph.15934>
- Silverman B W 2018 *Density Estimation for Statistics and Data Analysis* (Routledge) Online: <https://www.taylorfrancis.com/books/9781351456173>
- Still C J, Rastogi B, Page G F M, Griffith D M, Sibley A, Schulze M, Hawkins L, Pau S, Detto M and Helliker B R 2021 Imaging canopy temperature: shedding (thermal) light on ecosystem processes *New Phytologist* **230** 1746–53 Online: <https://onlinelibrary.wiley.com/doi/10.1111/nph.17321>
- Su Z 2002 The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes *Hydrol Earth Syst Sci* **6** 85–100 Online: <https://hess.copernicus.org/articles/6/85/2002/>
- Talsma C, Good S, Miralles D, Fisher J, Martens B, Jimenez C and Purdy A 2018 Sensitivity of Evapotranspiration Components in Remote Sensing-Based Models *Remote Sens (Basel)* **10** 1601 Online: <http://www.mdpi.com/2072-4292/10/10/1601>
- Tong B, Guo J, Xu H, Wang Y, Li H, Bian L, Zhang J and Zhou S 2022 Effects of soil moisture, net radiation, and atmospheric vapor pressure deficit on surface evaporation fraction at a semi-arid grass site *Science of The Total Environment* **849** 157890 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0048969722049890>
- URycki D R, Bassiouni M, Good S P, Crump B C and Li B 2022 The streamwater microbiome encodes hydrologic data across scales *Science of The Total Environment* **849** 157911 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0048969722050100>
- Veroustraete F, Patyn J and Myneni R B 1996 Estimating net ecosystem exchange of carbon using the normalized difference vegetation index and an ecosystem model *Remote Sens Environ* **58** 115–30 Online: <https://linkinghub.elsevier.com/retrieve/pii/0034425795002588>

- Wang L, Caylor K K, Villegas J C, Barron-Gafford G A, Breshears D D and Huxman T E 2010 Partitioning evapotranspiration across gradients of woody plant cover: Assessment of a stable isotope technique *Geophys Res Lett* **37** n/a-n/a Online: <http://doi.wiley.com/10.1029/2010GL043228>
- Wang L, Good S P and Caylor K K 2014 Global synthesis of vegetation control on evapotranspiration partitioning *Geophys Res Lett* **41** 6753–7 Online: <http://doi.wiley.com/10.1002/2014GL061439>
- Wang P, Li D, Liao W, Rigden A and Wang W 2019 Contrasting Evaporative Responses of Ecosystems to Heatwaves Traced to the Opposing Roles of Vapor Pressure Deficit and Surface Resistance *Water Resour Res* **55** 4550–63 Online: <https://onlinelibrary.wiley.com/doi/10.1029/2019WR024771>
- Weiskopf S R, Rubenstein M A, Crozier L G, Gaichas S, Griffis R, Halofsky J E, Hyde K J W, Morelli T L, Morissette J T, Muñoz R C, Pershing A J, Peterson D L, Poudel R, Staudinger M D, Sutton-Grier A E, Thompson L, Vose J, Weltzin J F and Whyte K P 2020 Climate change effects on biodiversity, ecosystems, ecosystem services, and natural resource management in the United States *Science of The Total Environment* **733** 137782
- Whelan M E, Lennartz S T, Gimeno T E, Wehr R, Wohlfahrt G, Wang Y, Kooijmans L M J, Hilton T W, Belviso S, Peylin P, Commane R, Sun W, Chen H, Kuai L, Mammarella I, Maseyk K, Berkelhammer M, Li K-F, Yakir D, Zumkehr A, Katayama Y, Ogée J, Spielmann F M, Kitz F, Rastogi B, Kesselmeier J, Marshall J, Erkkilä K-M, Wingate L, Meredith L K, He W, Bunk R, Launois T, Vesala T, Schmidt J A, Fichot C G, Seibt U, Saleska S, Saltzman E S, Montzka S A, Berry J A and Campbell J E 2018 Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water cycles *Biogeosciences* **15** 3625–57 Online: <https://bg.copernicus.org/articles/15/3625/2018/>
- Wibral M, Priesemann V, Kay J W, Lizier J T and Phillips W A 2017 Partial information decomposition as a unified approach to the specification of neural goal functions *Brain Cogn* **112** 25–38 Online: <https://linkinghub.elsevier.com/retrieve/pii/S027826261530021X>
- Williams M, Richardson A D, Reichstein M, Stoy P C, Peylin P, Verbeeck H, Carvalhais N, Jung M, Hollinger D Y, Kattge J, Leuning R, Luo Y, Tomelleri E, Trudinger C M and Wang Y-P 2009 Improving land surface models with FLUXNET data *Biogeosciences* **6** 1341–59 Online: <https://bg.copernicus.org/articles/6/1341/2009/>
- Williams P L and Beer R D 2010 Nonnegative Decomposition of Multivariate Information Online: <http://arxiv.org/abs/1004.2515>
- Wood D A 2021 Net ecosystem carbon exchange prediction and insightful data mining with an optimized data-matching algorithm *Ecol Indic* **124** 107426 Online: <https://linkinghub.elsevier.com/retrieve/pii/S1470160X21000911>



- Wutzler T, Lucas-Moffat A, Migliavacca M, Knauer J, Sickel K, Šigut L, Menzer O and Reichstein M 2018 Basic and extensible post-processing of eddy covariance flux data with REddyProc *Biogeosciences* **15** 5015–30 Online: <https://bg.copernicus.org/articles/15/5015/2018/>
- Xiao W, Wei Z and Wen X 2018 Evapotranspiration partitioning at the ecosystem scale using the stable isotope method—A review *Agric For Meteorol* **263** 346–61 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0168192318303009>
- Yang Y, Cui Y, Bai K, Luo T, Dai J, Wang W and Luo Y 2019 Short-term forecasting of daily reference evapotranspiration using the reduced-set Penman-Monteith model and public weather forecasts *Agric Water Manag* **211** 70–80 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0378377418314732>
- Yetemen O, Istanbuluoglu E, Flores-Cervantes J H, Vivoni E R and Bras R L 2015 Ecohydrologic role of solar radiation on landscape evolution *Water Resour Res* **51** 1127–57 Online: <http://doi.wiley.com/10.1002/2014WR016169>
- Yusup Y and Liu H 2020 Effects of persistent wind speeds on turbulent fluxes in the water-atmosphere interface *Theor Appl Climatol* **140** 313–25 Online: <http://link.springer.com/10.1007/s00704-019-03084-4>
- Zeng S, Xia J, Chen X, Zou L, Du H and She D 2020 Integrated land-surface hydrological and biogeochemical processes in simulating water, energy and carbon fluxes over two different ecosystems *J Hydrol (Amst)* **582** 124390 Online: <https://linkinghub.elsevier.com/retrieve/pii/S0022169419311254>
- Zhao W L, Qiu G Y, Xiong Y J, Paw U K T, Gentine P and Chen B Y 2020 Uncertainties Caused by Resistances in Evapotranspiration Estimation Using High-Density Eddy Covariance Measurements *J Hydrometeorol* **21** 1349–65 Online: <https://journals.ametsoc.org/view/journals/hydr/21/6/JHM-D-19-0191.1.xml>
- Zhou S, Yu B, Zhang Y, Huang Y and Wang G 2018 Water use efficiency and evapotranspiration partitioning for three typical ecosystems in the Heihe River Basin, northwestern China *Agric For Meteorol* **253–254** 261–73 Online: <https://linkinghub.elsevier.com/retrieve/pii/S016819231830039X>
1. Jung M, Reichstein M, Margolis HA, Cescatti A, Richardson AD, Arain MA, et al. Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations. *J Geophys Res* [Internet]. 2011 Sep 3;116:G00J07. Available from: <http://doi.wiley.com/10.1029/2010JG001566>
2. Piao S, Wang X, Wang K, Li X, Bastos A, Canadell JG, et al. Interannual variation of terrestrial carbon cycle: Issues and perspectives. *Glob Chang Biol* [Internet]. 2020 Jan 29;26(1):300–18. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.14884>

3. Baldocchi D. Measuring fluxes of trace gases and energy between ecosystems and the atmosphere—the state and future of the eddy covariance method. *Glob Chang Biol* [Internet]. 2014 Dec;20(12):3600–9. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.12649>
4. Schimel D, Schneider FD. Flux towers in the sky: global ecology from space. *New Phytologist* [Internet]. 2019 Oct 3;224(2):570–84. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/nph.15934>
5. Zhou S, Yu B, Zhang Y, Huang Y, Wang G. Water use efficiency and evapotranspiration partitioning for three typical ecosystems in the Heihe River Basin, northwestern China. *Agric For Meteorol* [Internet]. 2018 May;253–254:261–73. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S016819231830039X>
6. Zeng S, Xia J, Chen X, Zou L, Du H, She D. Integrated land surface hydrological and biogeochemical processes in simulating water, energy and carbon fluxes over two different ecosystems. *J Hydrol (Amst)* [Internet]. 2020 Mar;582:124390. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0022169419311254>
7. Jia Z, Liu S, Xu Z, Chen Y, Zhu M. Validation of remotely sensed evapotranspiration over the Hai River Basin, China. *Journal of Geophysical Research: Atmospheres* [Internet]. 2012 Jul 16;117(D13):n/a–n/a. Available from: <http://doi.wiley.com/10.1029/2011JD017037>
8. Williams M, Richardson AD, Reichstein M, Stoy PC, Peylin P, Verbeeck H, et al. Improving land surface models with FLUXNET data. *Biogeosciences* [Internet]. 2009 Jul 30;6(7):1341–59. Available from: <https://bg.copernicus.org/articles/6/1341/2009/>
9. Li S, Kang S, Li F, Zhang L. Evapotranspiration and crop coefficient of spring maize with plastic mulch using eddy covariance in northwest China. *Agric Water Manag* [Internet]. 2008 Nov;95(11):1214–22. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0378377408001169>
10. Lupascu M, Akhtar H, Smith TEL, Sukri RS. Post-fire carbon dynamics in the tropical peat swamp forests of Brunei reveal long-term elevated CH<sub>4</sub> flux. *Glob Chang Biol* [Internet]. 2020 Sep 15;26(9):5125–45. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.15195>
11. Dubbert M, Werner C. Water fluxes mediated by vegetation: emerging isotopic insights at the soil and atmosphere interfaces. *New Phytologist* [Internet]. 2019 Mar 19;221(4):1754–63. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/nph.15547>
12. Whelan ME, Lennartz ST, Gimeno TE, Wehr R, Wohlfahrt G, Wang Y, et al. Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and

- water cycles. *Biogeosciences* [Internet]. 2018 Jun 18;15(12):3625–57. Available from: <https://bg.copernicus.org/articles/15/3625/2018/>
13. Still CJ, Rastogi B, Page GFM, Griffith DM, Sibley A, Schulze M, et al. Imaging canopy temperature: shedding (thermal) light on ecosystem processes. *New Phytologist* [Internet]. 2021 Jun 2;230(5):1746–53. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/nph.17321>
14. Guan K, Berry JA, Zhang Y, Joiner J, Guanter L, Badgley G, et al. Improving the monitoring of crop productivity using spaceborne solar-induced fluorescence. *Glob Chang Biol* [Internet]. 2016 Feb 10;22(2):716–26. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/gcb.13136>
15. URycki DR, Bassiouni M, Good SP, Crump BC, Li B. The streamwater microbiome encodes hydrologic data across scales. *Science of The Total Environment* [Internet]. 2022 Nov;849:157911. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0048969722050100>
16. Bowen GJ, Good SP. Incorporating water isoscapes in hydrological and water resource investigations. *WIREs Water* [Internet]. 2015 Mar 16;2(2):107–19. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/wat2.1069>
17. Xiao W, Wei Z, Wen X. Evapotranspiration partitioning at the ecosystem scale using the stable isotope method—A review. *Agric For Meteorol* [Internet]. 2018 Dec;263:346–61. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0168192318303009>
18. Berkelhammer M, Hu J, Bailey A, Noone DC, Still CJ, Barnard H, et al. The nocturnal water cycle in an open-canopy forest. *Journal of Geophysical Research: Atmospheres* [Internet]. 2013 Sep 16;118(17):10,225–10,242. Available from: <http://doi.wiley.com/10.1002/jgrd.50701>
19. Lee SC, Christen A, Black TA, Jassal RS, Ketler R, Nesic Z. Partitioning of net ecosystem exchange into photosynthesis and respiration using continuous stable isotope measurements in a Pacific Northwest Douglas-fir forest ecosystem. *Agric For Meteorol* [Internet]. 2020 Oct;292–293:108109. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0168192320302112>
20. Orlowski N, Breuer L, Angeli N, Boeckx P, Brumbt C, Cook CS, et al. Inter-laboratory comparison of cryogenic water extraction systems for stable isotope analysis of soil water. *Hydrol Earth Syst Sci*. 2018 Jul 6;22(7):3619–37.
21. Gemery PA, Trolier M, White JWC. Oxygen isotope exchange between carbon dioxide and water following atmospheric sampling using glass flasks. *Journal of Geophysical Research: Atmospheres*. 1996 Jun 20;101(D9):14415–20.

22. Fiorella RP, Good SP, Allen ST, Guo JS, Still CJ, Noone DC, et al. Calibration Strategies for Detecting Macroscale Patterns in NEON Atmospheric Carbon Isotope Observations. *J Geophys Res Biogeosci*. 2021 Mar 26;126(3).
23. Finkenbiner CE, Li B, Spencer L, Butler Z, Haagsma M, Fiorella RP, et al. The NEON Daily Isotopic Composition of Environmental Exchanges Dataset. *Sci Data*. 2022 Dec 21;9(1):353.
24. Cover TM, Thomas JA. Elements of Information Theory [Internet]. Wiley; 2005. Available from: <https://onlinelibrary.wiley.com/doi/book/10.1002/047174882X>
25. Goodwell AE, Kumar P. Temporal information partitioning: Characterizing synergy, uniqueness, and redundancy in interacting environmental variables. *Water Resour Res* [Internet]. 2017 Jul 24;53(7):5920–42. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/2016WR020216>
26. Williams PL, Beer RD. Nonnegative Decomposition of Multivariate Information. 2010 Apr 14; Available from: <http://arxiv.org/abs/1004.2515>
27. Barnett DT, Adler PB, Chemel BR, Duffy PA, Enquist BJ, Grace JB, et al. The plant diversity sampling design for The National Ecological Observatory Network. *Ecosphere* [Internet]. 2019 Feb 25;10(2). Available from: <https://onlinelibrary.wiley.com/doi/10.1002/ecs2.2603>
28. National Ecological Observatory Network (NEON). Bundled data products – eddy covariance (DP4.00200.001) [Internet]. National Ecological Observatory Network (NEON); 2022. Available from: <https://data.neonscience.org>
29. National Ecological Observatory Network (NEON). Relative humidity (DP1.00098.001). National Ecological Observatory Network (NEON); 2022.
30. Finkenbiner CE, Li B, Spencer L, Butler Z, Haagsma M, Fiorella RP, et al. The NEON Daily Isotopic Composition of Environmental Exchanges Dataset. *Sci Data* [Internet]. 2022 Dec 21;9(1):353. Available from: <https://www.nature.com/articles/s41597-022-01412-4>
31. Silverman BW. Density Estimation for Statistics and Data Analysis [Internet]. Routledge; 2018. Available from: <https://www.taylorfrancis.com/books/9781351456173>
32. Goodwell AE, Kumar P, Fellows AW, Flerchinger GN. Dynamic process connectivity explains ecohydrologic responses to rainfall pulses and drought. *Proceedings of the National Academy of Sciences* [Internet]. 2018 Sep 11;115(37). Available from: <https://pnas.org/doi/full/10.1073/pnas.1800236115>
33. Goodwell AE, Kumar P. Temporal information partitioning: Characterizing synergy, uniqueness, and redundancy in interacting environmental variables. *Water Resour Res*. 2017 Jul 24;53(7):5920–42.

34. Yusup Y, Liu H. Effects of persistent wind speeds on turbulent fluxes in the water-atmosphere interface. *Theor Appl Climatol* [Internet]. 2020 Apr 13;140(1–2):313–25. Available from: <http://link.springer.com/10.1007/s00704-019-03084-4>
35. Cosgrove BA, Lohmann D, Mitchell KE, Houser PR, Wood EF, Schaake JC, et al. Real-time and retrospective forcing in the North American Land Data Assimilation System (NLDAS) project. *Journal of Geophysical Research: Atmospheres* [Internet]. 2003 Nov 27;108(D22):2002JD003118. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1029/2002JD003118>
36. Rodell M, Houser PR, Jambor U, Gottschalck J, Mitchell K, Meng CJ, et al. The Global Land Data Assimilation System. *Bull Am Meteorol Soc* [Internet]. 2004 Mar;85(3):381–94. Available from: <https://journals.ametsoc.org/doi/10.1175/BAMS-85-3-381>
37. Safa B, Arkebauer TJ, Zhu Q, Suyker A, Irmak S. Net Ecosystem Exchange (NEE) simulation in maize using artificial neural networks. *IFAC Journal of Systems and Control* [Internet]. 2019 Mar;7:100036. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2468601817302584>
38. Talsma C, Good S, Miralles D, Fisher J, Martens B, Jimenez C, et al. Sensitivity of Evapotranspiration Components in Remote Sensing-Based Models. *Remote Sens (Basel)* [Internet]. 2018 Oct 9;10(10):1601. Available from: <http://www.mdpi.com/2072-4292/10/10/1601>
39. Yetemen O, Istanbuluoglu E, Flores-Cervantes JH, Vivoni ER, Bras RL. Ecohydrologic role of solar radiation on landscape evolution. *Water Resour Res* [Internet]. 2015 Feb;51(2):1127–57. Available from: <http://doi.wiley.com/10.1002/2014WR016169>
40. Clausius R. Ueber die bewegende Kraft der Wärme und die Gesetze, welche sich daraus für die Wärmelehre selbst ableiten lassen. *Annalen der Physik und Chemie* [Internet]. 1850;155(4):500–24. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/andp.18501550403>
41. Chen J, Wen J, Kang S, Meng X, Tian H, Ma X, et al. Assessments of the factors controlling latent heat flux and the coupling degree between an alpine wetland and the atmosphere on the Qinghai-Tibetan Plateau in summer. *Atmos Res* [Internet]. 2020 Aug;240:104937. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S016980951931573X>
42. Niu S, Luo Y, Fei S, Yuan W, Schimel D, Law BE, et al. Thermal optimality of net ecosystem exchange of carbon dioxide and underlying mechanisms. *New Phytologist* [Internet]. 2012 May 7;194(3):775–83. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2012.04095.x>




43. Gu L, Meyers T, Pallardy SG, Hanson PJ, Yang B, Heuer M, et al. Direct and indirect effects of atmospheric conditions and soil moisture on surface energy partitioning revealed by a prolonged drought at a temperate forest site. *J Geophys Res* [Internet]. 2006;111(D16):D16102. Available from: <http://doi.wiley.com/10.1029/2006JD007161>
44. Tong B, Guo J, Xu H, Wang Y, Li H, Bian L, et al. Effects of soil moisture, net radiation, and atmospheric vapor pressure deficit on surface evaporation fraction at a semi-arid grass site. *Science of The Total Environment* [Internet]. 2022 Nov;849:157890. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0048969722049890>
45. Wang P, Li D, Liao W, Rigden A, Wang W. Contrasting Evaporative Responses of Ecosystems to Heatwaves Traced to the Opposing Roles of Vapor Pressure Deficit and Surface Resistance. *Water Resour Res* [Internet]. 2019 Jun 4;55(6):4550–63. Available from: <https://onlinelibrary.wiley.com/doi/10.1029/2019WR024771>
46. Grossiord C, Buckley TN, Cernusak LA, Novick KA, Poulter B, Siegwolf RTW, et al. Plant responses to rising vapor pressure deficit. *New Phytologist* [Internet]. 2020 Jun 20;226(6):1550–66. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/nph.16485>
47. Yang Y, Cui Y, Bai K, Luo T, Dai J, Wang W, et al. Short-term forecasting of daily reference evapotranspiration using the reduced-set Penman-Monteith model and public weather forecasts. *Agric Water Manag* [Internet]. 2019 Jan;211:70–80. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0378377418314732>
48. Liu X, Zhang D. Trend analysis of reference evapotranspiration in Northwest China: The roles of changing wind speed and surface air temperature. *Hydrol Process* [Internet]. 2013 Dec 30;27(26):3941–8. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/hyp.9527>
49. Ruddell BL, Drewry DT, Nearing GS. Information Theory for Model Diagnostics: Structural Error is Indicated by Trade-Off Between Functional and Predictive Performance. *Water Resour Res* [Internet]. 2019 Aug 6;55(8):6534–54. Available from: <https://onlinelibrary.wiley.com/doi/10.1029/2018WR023692>
50. Li B, Good SP. Information-based uncertainty decomposition in dual-channel microwave remote sensing of soil moisture. *Hydrol Earth Syst Sci* [Internet]. 2021 Sep 17;25(9):5029–45. Available from: <https://hess.copernicus.org/articles/25/5029/2021/>
51. Gong W, Gupta H V., Yang D, Sricharan K, Hero AO. Estimating epistemic and aleatory uncertainties during hydrologic modeling: An information theoretic approach. *Water Resour Res* [Internet]. 2013 Apr;49(4):2253–73. Available from: <http://doi.wiley.com/10.1002/wrcr.20161>

52. Fisher RA, Koven CD. Perspectives on the Future of Land Surface Models and the Challenges of Representing Complex Terrestrial Systems. *J Adv Model Earth Syst* [Internet]. 2020 Apr 20;12(4). Available from: <https://onlinelibrary.wiley.com/doi/10.1029/2018MS001453>
53. Wood DA. Net ecosystem carbon exchange prediction and insightful data mining with an optimized data-matching algorithm. *Ecol Indic* [Internet]. 2021 May;124:107426. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1470160X21000914>
54. Su Z. The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes. *Hydrol Earth Syst Sci* [Internet]. 2002 Feb 28;6(1):85–100. Available from: <https://hess.copernicus.org/articles/6/85/2002/>
55. Veroustraete F, Patyn J, Myneni RB. Estimating net ecosystem exchange of carbon using the normalized difference vegetation index and an ecosystem model. *Remote Sens Environ* [Internet]. 1996 Oct;58(1):115–30. Available from: <https://linkinghub.elsevier.com/retrieve/pii/0034425795002588>
56. Papale D, Reichstein M, Aubinet M, Canfora E, Bernhofer C, Kutsch W, et al. Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation. *Biogeosciences* [Internet]. 2006 Nov 27;3(4):571–83. Available from: <https://bg.copernicus.org/articles/3/571/2006/>
57. Zhao WL, Qiu GY, Xiong YJ, Paw U KT, Gentile P, Chen BY. Uncertainties Caused by Resistances in Evapotranspiration Estimation Using High-Density Eddy Covariance Measurements. *J Hydrometeorol* [Internet]. 2020 Jun;21(6):1349–65. Available from: <https://journals.ametsoc.org/view/journals/hydr/21/6/JHM-D-19-0191.1.xml>
58. Wibral M, Priesemann V, Kay JW, Lizier JT, Phillips WA. Partial information decomposition as a unified approach to the specification of neural goal functions. *Brain Cogn* [Internet]. 2017 Mar;112:25–38. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S027826261530021X>
59. Good SP, Soderberg K, Guan K, King EG, Scanlon TM, Caylor KK.  $\delta^2\text{H}$  isotopic flux partitioning of evapotranspiration over a grass field following a water pulse and subsequent dry down. *Water Resour Res* [Internet]. 2014 Feb;50(2):1410–32. Available from: <http://doi.wiley.com/10.1002/2013WR014333>
60. Conrad R, Klose M, Yuan Q, Lu Y, Chidthaisong A. Stable carbon isotope fractionation, carbon flux partitioning and priming effects in anoxic soils during methanogenic degradation of straw and soil organic matter. *Soil Biol Biochem* [Internet]. 2012 Jun;49:193–9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0038071712000958>

61. Wang L, Caylor KK, Villegas JC, Barron-Gafford GA, Breshears DD, Huxman TE. Partitioning evapotranspiration across gradients of woody plant cover: Assessment of a stable isotope technique. *Geophys Res Lett* [Internet]. 2010 May;37(9):n/a-n/a. Available from: <http://doi.wiley.com/10.1029/2010GL043228>
62. Good SP, Noone D, Kurita N, Benetti M, Bowen GJ. D/H isotope ratios in the global hydrologic cycle. *Geophys Res Lett* [Internet]. 2015 Jun 28;42(12):5042–50. Available from: <http://doi.wiley.com/10.1002/2015GL064117>
63. Wang L, Good SP, Caylor KK. Global synthesis of vegetation control on evapotranspiration partitioning. *Geophys Res Lett* [Internet]. 2014 Oct 16;41(19):6753–7. Available from: <http://doi.wiley.com/10.1002/2014GL061439>
64. Bowling DR, Pataki DE, Randerson JT. Carbon isotopes in terrestrial ecosystem pools and CO<sub>2</sub> fluxes. *New Phytologist* [Internet]. 2008 Apr 7;178(1):24–40. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2007.02342.x>

973  
974  
975




976  
977  
978  
979  
980  
981

**Figure 1** Individual mutual information shared between net ecosystem exchange (*NEE*) and each individual meteorological variable (vapor pressure deficit [*VPD*], air temperature [*T*], global radiation [*R<sub>g</sub>*], windspeed [*u*]) (a). Individual information shared between latent heat flux (*LH*) and each individual meteorological variable (b). The mean and median

values of each boxplot are shown as black triangle and white line, respectively. The double asterisk indicates a significant p-value ( $<0.01$ ).

**Figure 1** (a) Individual mutual information,  $I(X;Y)$ , shared between net ecosystem exchange,  $NEE$ , and each individual meteorological variable (vapor pressure deficit,  $VPD$ , air temperature,  $T$ , global radiation,  $R_g$ , windspeed,  $u$ ). (b) Individual information shared between latent heat flux,  $LH$ , and each individual meteorological variable. Boxes of mutual information between meteorological variables and flux are consists of the same quantity that is calculated based on different isotope availability (e.g., box of  $I(VPD; NEE)$  consists of  $I(VPD; NEE)$  based on the availability of  $\delta^{13}C$ ,  $I(VPD; NEE)$  based on the availability of  $\delta^2H$ , and  $I(VPD; NEE)$  on the availability of  $d$ , collectively). The mean and median values of each boxplot are shown as black triangle and white line, respectively. The double asterisk indicates a significant p-value ( $<0.01$ ).






**Figure 2** (a) The unique information,  $U$ , synergistic information,  $S$ , and redundant information,  $R$ , of the  $\delta^{13}\text{C}$ ,  $\delta^2\text{H}$ , and  $d$  stable isotope flux ratios on the net ecosystem

exchange, *NEE*, and (b) latent heat flux, *LH*. The values of *S*, and *R* are calculated by averaging across different meteorological variables, indicated by *Y* (e.g., the average over  $S(\delta^2H, VPD; LH)$ ,  $S(\delta^2H, T; LH)$ ,  $S(\delta^2H, u; LH)$ , and  $S(\delta^2H, R_g; LH)$  for *S*). The mean and median values of each boxplot are shown as black triangle and white line, respectively. The double asterisk indicates a significant p-value (<0.01).

**Figure 2** The unique information (*U*), synergistic information (*S*), and redundant information (*R*) of the stable isotope flux ratios on the net ecosystem exchange [*NEE*] (a) and latent heat flux (*LH*) (b) fluxes. The mean and median values of each boxplot are shown as black triangle and white line, respectively. The double asterisk indicates a significant p-value (<0.01).



**Figure 3** The additive information of (a)  $\delta^{13}\text{C}$ , (b)  $\delta^2\text{H}$ , and (c)  $d$  isotope data about net ecosystem exchange,  $NEE$ . The additive information of (d)  $\delta^{13}\text{C}$ , (e)  $\delta^2\text{H}$ , and (f)  $d$  isotope data about latent heat flux,  $LH$ . The additive information is the sum unique,  $U$ , and synergistic,  $S$ , information added by each data source. **Figure 3** The additive information of  $\delta^{13}\text{C}$  about net ecosystem exchange ( $NEE$ ) (a) and latent heat flux ( $LH$ ) (d). The additive information of  $\delta^{13}\text{H}$  about  $NEE$  (b) and latent heat flux ( $LH$ ) (e). The additive information of  $d$  about  $NEE$  (c) and latent heat flux ( $LH$ ) (f).

1046

1047

1048

1049

1050

1051


1052

1053

1054

1055

1056





**Figure 4** The total added information of (a)  $\delta^{13}C$ , (b)  $\delta^2H$ , and (c)  $d$  isotope data about net ecosystem exchange,  $NEE$ , against scaled site-specific variables. The total added information of (d)  $\delta^{13}C$ , (e)  $\delta^2H$ , and (f)  $d$  isotope data about latent heat flux,  $LH$  against scaled site-specific variables. Solid lines indicate a significant p-values ( $< 0.05$ ) of the slopes.

~~**Figure 4** The total added information of  $\delta^{13}C$  (a),  $\delta^2H$  (b), and  $d$  (c) about net ecosystem exchange ( $NEE$ ) against scaled site-specific variables. The total added information of  $\delta^{13}C$  (d),  $\delta^2H$  (e), and  $d$  (f) about latent heat flux ( $LH$ ) against scaled site-specific variables. Solid lines indicate a significant p-values ( $< 0.05$ ) of the slopes.~~