

Theory & Notes

1. XX-MODEL

Consider the XX-Heisenberg model, with its Hamiltonian given in terms of the traditional $\frac{1}{2}$ -spin operators ie.

$$(1) \quad \mathbf{H} = J \sum_{i=1}^L (\mathbf{S}_i^x \mathbf{S}_{i+1}^x + \mathbf{S}_i^y \mathbf{S}_{i+1}^y) - \lambda \sum_{j=1}^L \mathbf{S}_j^z,$$

which describes interacting spins in a one-dimensional chain, with periodic boundary conditions. (1)'s first terms represents nearest neighbour interactions in the x and y -directions interactions, with J being either positive or negative and quantifying the strength and type of interactions, while the second term represents a magnetic field of strength λ , applied in the z -direction of the spins.

In order to solve this problem, it is necessary to rewrite (1) and apply a Jordan-Wigner transformation, mapping the spin problem into a fermionic problem. But first, it is convenient to write the spin-operators in terms of the raising and lowering $\mathfrak{su}(2)$ -operators, ie.

$$\mathbf{S}_j^\pm = \mathbf{S}_j^x \pm i\mathbf{S}_j^y \Rightarrow \begin{aligned} \mathbf{S}_j^x &= \frac{1}{2}(\mathbf{S}_j^+ + \mathbf{S}_j^-), \\ \mathbf{S}_j^y &= \frac{1}{2i}(\mathbf{S}_j^+ - \mathbf{S}_j^-). \end{aligned}$$

Then, the XX-Hamiltonian can be re-written as

$$(2) \quad \mathbf{H} = \frac{J}{2} \sum_{i=1}^L (\mathbf{S}_i^+ \mathbf{S}_{i+1}^- + \mathbf{S}_i^- \mathbf{S}_{i+1}^+) - \lambda \sum_{j=1}^L \mathbf{S}_j^z$$

where the interacting terms in the first summation, flip neighboring spins if said spins are anti-aligned¹. In addition, the XX-Hamiltonian has a total magnetization symmetry, since the Hamiltonian given by (2) commutes with the magnetization operator.

Now, we use a Jordan-Wigner transformation whereby the spin operators are mapped to fermionic operators, as follows

¹In effect, consider for example, a two-spin problem. Then, the interaction term is given by

$$\mathbf{S}_1^+ \mathbf{S}_2^- + \mathbf{S}_1^- \mathbf{S}_2^+,$$

and consider an anti-aligned state $|\downarrow\uparrow\rangle$. Then, the action of the previous two-spin operator over this state yields

$$(\mathbf{S}_1^+ \mathbf{S}_2^- + \mathbf{S}_1^- \mathbf{S}_2^+) |\downarrow\uparrow\rangle = |\uparrow\downarrow\rangle + 0,$$

since $\mathbf{S}_1^- \mathbf{S}_2^+$ destroys the state. Similarly, $(\mathbf{S}_1^+ \mathbf{S}_2^- + \mathbf{S}_1^- \mathbf{S}_2^+) |\uparrow\downarrow\rangle = |\downarrow\uparrow\rangle$. However, note that, should both spins be either up or down, the state remain invariant under the action of the two-spin operator.

$$(3) \quad (\mathbf{S}_1^+ \mathbf{S}_2^- + \mathbf{S}_1^- \mathbf{S}_2^+) |\downarrow\downarrow\rangle = |\downarrow\downarrow\rangle \text{ and } (\mathbf{S}_1^+ \mathbf{S}_2^- + \mathbf{S}_1^- \mathbf{S}_2^+) |\uparrow\uparrow\rangle = |\uparrow\uparrow\rangle$$

$$\begin{aligned}
\mathbf{S}_j^z &= f_j^\dagger f_j - \frac{1}{2} \\
(4) \quad \mathbf{S}_j^- &= \exp \left(i\pi \sum_{\ell=1}^{L-1} f_\ell^\dagger f_\ell \right) \\
\mathbf{S}_j^+ &= \exp \left(-i\pi \sum_{\ell=1}^{L-1} f_\ell^\dagger f_\ell \right)
\end{aligned}$$

Under the Jordan-Wigner map, nearest-neighbours spin flipping is translated into nearest-neighbours fermionic hopping, ie. $\mathbf{S}_j^+ \mathbf{S}_{j+1}^- = f_j^\dagger f_{j+1}$ and $\mathbf{S}_j^- \mathbf{S}_{j+1}^+ = f_{j+1}^\dagger f_j$. However, due to the boundary conditions' periodicity, the XX-Hamiltonian cannot be rewritten as a fermionic model yet since (it will contain an additional boundary term), for example, the fermionic counterparts to the $\mathbf{S}_L^+ \mathbf{S}_1^-$ interaction are highly non-local operators and are not desirable. Indeed, under the Jordan-Wigner mapping

$$\mathbf{S}_L^+ \mathbf{S}_1^- = f_L^\dagger \exp \left(-i\pi \sum_{\ell=1}^{L-1} f_\ell^\dagger f_\ell \right) f_1,$$

which is not problematic, since it accounts for all L -lattice sites. Let

$$\begin{aligned}
(5) \quad \mathbf{S}_L^+ \mathbf{S}_1^- &= \mathcal{Q} f_L^\dagger f_1, \\
\mathbf{S}_L^- \mathbf{S}_1^+ &= \mathcal{Q} f_1^\dagger f_L,
\end{aligned}$$

then (2) can be rewritten as

$$(6) \quad \mathbf{H} = \frac{J}{2} \sum_{i=1}^{L-1} \left(f_i^\dagger f_{i+1} + f_{i+1}^\dagger f_i \right) - \lambda \sum_{j=1}^L \left(f_j^\dagger f_j - \frac{1}{2} \right) + \frac{J}{2} \mathcal{Q} (f_L^\dagger f_1 + f_1^\dagger f_L),$$

where the first term accounts for fermionic nearest-neighbour hopping, the second term accounts for the magnetic field, and the third term being the non-local boundary term. Note that this fermionic Hamiltonian hasn't got any type of boundary conditions, since the L -lattice site is disconnected in any way whatsoever from the first lattice site. Then, the standard procedure is to add and subtract terms from the Hamiltonian, so that the nearest-neighbour hopping term in (6) can also have periodic boundary conditions, thus yielding

$$(7) \quad \mathbf{H} = \frac{J}{2} \sum_{i=1}^L \left(f_i^\dagger f_{i+1} + f_{i+1}^\dagger f_i \right) - \lambda \sum_{j=1}^L \left(f_j^\dagger f_j - \frac{1}{2} \right) + \frac{J}{2} (\mathcal{Q} - 1) (f_L^\dagger f_1 + f_1^\dagger f_L),$$

where now the fermionic hopping term has the standard boundary conditions. The third term, since it does not involve any type of summation over lattice sites, only contributes at $\mathcal{O}\left(\frac{1}{L}\right)$ -order to any microscopic quantity. In the thermodynamic limit, this non-local term can be dropped, thus yielding an $\mathcal{O}(L)$ -Hamiltonian given by

$$(8) \quad \mathbf{H} = \frac{J}{2} \sum_{i=1}^L \left(f_i^\dagger f_{i+1} + f_{i+1}^\dagger f_i - \lambda f_i^\dagger f_i \right) + \frac{\lambda L}{2},$$

which is now fully cyclic and where its operators obey fermionic algebras. This Hamiltonian can then be diagonalized via a discrete Fourier transform on the fermionic operators

$$(9) \quad f_j = \frac{1}{\sqrt{L}} \sum_{\substack{k=2\pi m/L \\ m \in \mathbb{Z}_{[1,L]}}} e^{ijk} d_k, \quad f_j^\dagger = \frac{1}{\sqrt{L}} \sum_{\substack{k=2\pi m/L \\ m \in \mathbb{Z}_{[1,L]}}} e^{-ijk} d_k^\dagger,$$

with the inverse transformation given by

$$(10) \quad d_k = \frac{1}{\sqrt{L}} \sum_{j=1}^L e^{-ikj} f_j \quad d_k^\dagger = \frac{1}{\sqrt{L}} \sum_{j=1}^L e^{ikj} f_j^\dagger.$$

Note that the d_k -operators follow the standard fermionic anticommutation algebra. Note as well, that the f_j -vacuum state, defined such that $f_j |0\rangle_f = 0$, $\forall j$, is the same as the d_k -vacuum state, defined such that $d_j |0\rangle_d = 0$, $\forall k$, ie. $|0\rangle_f = |0\rangle_d$. Another important relationship is the Fourier transform's consistency condition, ie.

$$\sum_{j=1}^L e^{i(k-q)j} = L\delta_{kq}.$$

Under the Fourier transform, (8)'s terms are mapped as follows

$$\begin{aligned} \sum_{j=1}^L f_j^\dagger f_{j+1} &= \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{-ikj} e^{iq(j+1)} d_k^\dagger d_q = \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{i(q-k)j} e^{iq} d_k^\dagger d_q \\ &= \sum_{k,q} \frac{1}{L} e^{iq} \delta_{qk} d_k^\dagger d_q = \sum_k e^{ik} d_k^\dagger d_k \\ \sum_{j=1}^L f_{j+1}^\dagger f_j &= \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{-ik(j+1)} e^{iqj} d_k^\dagger d_q = \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{i(q-k)j} e^{-ik} d_k^\dagger d_q \\ &= \sum_{k,q} \frac{1}{L} e^{-ik} \delta_{qk} d_k^\dagger d_q = \sum_k e^{-ik} d_k^\dagger d_k \\ \sum_{j=1}^L f_j^\dagger f_j &= \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{-ikj} e^{iqj} d_k^\dagger d_q = \sum_{j=1}^L \frac{1}{L} \sum_{k,q} e^{i(q-k)j} d_k^\dagger d_q \\ &= \sum_{k,q} \frac{1}{L} \delta_{qk} d_k^\dagger d_q = \sum_k d_k^\dagger d_k \end{aligned}$$

Therefore, using these identities, the new Hamiltonian is given by

$$(11) \quad \mathbf{H} = \frac{J}{2} \sum_k \left(e^{ik} d_k^\dagger d_k + e^{-ik} d_k^\dagger d_k - \lambda d_k^\dagger d_k \right) + \frac{\lambda L}{2}$$

$$(12) \quad = \sum_k \left(J \cos k - \lambda \right) d_k^\dagger d_k + \frac{\lambda L}{2},$$

which can be rewritten as

$$(13) \quad \mathbf{H} = \sum_k \epsilon_k d_k^\dagger d_k + \frac{\lambda L}{2} \quad \text{with the eigenvalues being given by } \epsilon_k = J \cos k - \lambda + \frac{\lambda L}{2} \quad \text{and the eigenvectors being given by } |E_n\rangle = \prod_n (d_k^\dagger)^n, \quad \text{with eigenvalue } E_n = \sum_n \epsilon_n,$$

where, $k = \frac{2\pi m}{L}$, $m = -\frac{L}{2} + 1, \dots, \frac{L}{2}$, ie. $k \in (-\pi, \pi]$, this is called the first Brillouin zone. Thus the problem has been solved

As for its thermal properties, since this is a fermionic model, the fermions will obey the Fermi-Dirac distribution, ie.

$$(14) \quad \mathcal{N}_{jk} = \langle d_j^\dagger d_k \rangle_{\text{th}} = \frac{1}{1 + e^{\beta \epsilon_k + \mu}} \delta_{jk}$$

1.1. Quantum Phase Transition. The system's groundstate is defined such that

$$(15) \quad \langle E_G | d_k^\dagger d_k | E_G \rangle = 1 \text{ if } \epsilon_k \leq 0 \text{ and } \langle E_G | d_k^\dagger d_k | E_G \rangle = 0 \text{ if } \epsilon_k > 0.$$

Then the groundstate energy is given by

$$(16) \quad E_G = \sum_k \epsilon_k \langle E_G | d_k^\dagger d_k | E_G \rangle.$$

Now, there exists a momenta k_F such that $\epsilon_{k_F} = 0$, with k_F being the Fermi momentum, ie. $\epsilon_{k_f} = J \cos k_f - \lambda$. The previous equation depends both on the coupling strength J and the magnetic field strength λ . From the previous equation some special cases arise, namely

- when $\lambda < -J$, which yields a Fermi momentum $k_F = \pi$,
- when $|\lambda| \leq J$, which yields a Fermi momentum $k_F = \arccos\left(\frac{\lambda}{J}\right)$,
- when $\lambda > J$ which yields $k_F = 0$.

Therefore, the system groundstate will be defined by the following relations,

$$(17) \quad \begin{aligned} \langle E_G | d_k^\dagger d_k | E_G \rangle &= 1 & \langle E_G | d_k^\dagger d_k | E_G \rangle &= 0 \\ &\text{and} & & \\ \text{if } k \in [-\pi, -k_F] \cup [k_F, \pi], & & \text{if } k \in [-k_F, k_F]. \end{aligned}$$

The total magnetization operator is defined as the sum of the S_z operators, along the entire chain ie. $M^z = \sum_{j=1} S_j^z$. Under the Jordan-Wigner transformation, $S_j^z = f_j^\dagger f_j - \frac{1}{2}$. In turn, these fermionic number operators can be mapped to fermionic number operators acting on momentum space, ie. $\sum_{j=1}^L f_j^\dagger f_j = \sum_k d_k^\dagger d_k$. This entails

$$(18) \quad M^z = \sum_k \left(d_k^\dagger d_k - \frac{1}{2} \right).$$

Then the total magnetization operator's expectation value for the groundstate can be calculated. Given the system's fermionic nature, momentum eigenmodes are either occupied or unoccupied, contributing only one or zeros respectively. Then, this expectation value yields

$$(19) \quad \langle E_G | M^z | E_G \rangle = \sum_{k=-\pi}^{-k_F} 1 + \sum_{k=k_F}^{\pi} 1 - \frac{1}{2} \sum_{k=-\pi}^{\pi} 1 = \frac{1}{2} \left(\sum_{k=-\pi}^{-k_F} 1 + \sum_{k=k_F}^{\pi} 1 - \sum_{k=-k_F}^{k_F} 1 \right),$$

where the total magnetization expectation value is just half the difference of two competing summations, related to the negative and positive eigenmodes. Defining the total magnetization per site operator, $m^z = \frac{1}{L} M^z$. Now, as the chain gets progressively bigger, with more and more lattice sites, the summations in (19) can be approximated by Riemannian integrals, ie.

$$\sum_k f(k) = \frac{1}{\Delta k} \sum_k f(k) \Delta k \xrightarrow{L \rightarrow \infty} \frac{L}{2\pi} \int_{k \in \text{BZ}} dk f(k) \text{ where } \Delta k = \frac{2\pi}{L}.$$

Using this trick, (19) can be rewritten as

$$(20) \quad \langle m^z \rangle_G = \frac{1}{4\pi} \left(\int_{-\pi}^{-k_F} dk + \int_{k_F}^{\pi} dk - \int_{-k_F}^{k_F} dk \right) = \frac{1}{4\pi} \left(-k_F + \pi + \pi - k_F - k_F + k_F \right)$$

$$(21) \quad = \frac{1}{2} - \frac{k_F}{\pi},$$

which yields the total magnetization per lattice site. Note that at $\lambda < J$, $k_F = \pi$ whereby there is total polarization ie. $\langle \mathbf{m}^z \rangle_G = -\frac{1}{2}$. Similarly, for $\lambda > J$, $k_F = 0$ and then $\langle \mathbf{m}^z \rangle_G = \frac{1}{2}$. However, if $|\lambda| \leq J$, then $k_F = \arccos\left(\frac{\lambda}{J}\right)$, which then yields a total magnetization per lattice site given by

$$(22) \quad \langle \mathbf{m}^z \rangle_G = \begin{cases} \frac{1}{2} - \frac{1}{\pi} \arccos\left(\frac{\lambda}{J}\right) & \text{if } |\lambda| \leq J \\ -\frac{1}{2} & \text{if } \lambda < J \\ \frac{1}{2} & \text{if } \lambda > J \end{cases}$$

Another interesting quantity is the total magnetic susceptibility, which can be written in terms of the total magnetization per lattice site as follows

$$(23) \quad \chi = \left(\frac{\partial \langle \mathbf{m}^z \rangle_G}{\partial \lambda} \right) \Rightarrow \chi = \begin{cases} \frac{1}{J\pi\sqrt{1-\frac{\lambda^2}{J^2}}} & \text{if } |\lambda| \leq J \\ 0 & \text{if } \lambda < J \\ 0 & \text{if } \lambda > J \end{cases},$$

which is divergent if $\lambda = J$. This shows that the XX-Heisenberg model has a second order (since there is no latent heat involved it can't be a first order phase transition) quantum phase transition, with an associated power law with a critical exponent.

2. NUMERICAL SOLUTION TO FERMIONIC MODELS

Consider a Hamiltonian describing a fermionic system, given by

$$(24) \quad \mathbf{H} = J \sum_{j=1}^{L-1} (f_j^\dagger f_{j+1} + f_{j+1}^\dagger f_j) + \sum_{j=1}^L \lambda_j f_j^\dagger f_j, \quad \text{with the usual commutation rules} \quad \begin{cases} \{f_j, f_k\} = \{f_j^\dagger, f_k^\dagger\} = 0 \\ \{f_j, f_k^\dagger\} = \delta_{jk} \end{cases}$$

where L indicates the number of lattice sites, J is the hopping strength, which could be either positive or negative, and where λ_j is the on-site potential strength². Said Hamiltonian has open boundaries conditions since there is no hopping term across the boundary. Note that we can rewrite (24) as

$$(25) \quad \mathbf{H} = \sum_{i,j=1}^L \mathcal{M}_{ij} f_i^\dagger f_j \quad \text{with} \quad \mathcal{M}_{ij} = \begin{cases} \lambda_i & \text{if } i = j \\ J & \text{if } j = i + 1 \text{ or } i = j + 1 \\ 0 & \text{otherwise} \end{cases}, \quad \mathcal{M} \in \text{GL}(L, \mathbb{R}),$$

which is a positive-defined tri-diagonal matrix. Let $\mathbf{f} = (f_1 \ f_2 \ \cdots \ f_L)^T$ be a vector of the L fermionic operators. Then, (24) can be rewritten as

$$(26) \quad \mathbf{H} = \mathbf{f}^\dagger \mathcal{M} \mathbf{f}.$$

Since \mathcal{M} is symmetric, then it can be diagonalized $\mathcal{M} = A D A^T$, where $A \in \mathbb{R}^{L \times L}$ is a real orthogonal matrix and with $\mathcal{D}_{ij} \in \mathbb{R}^{L \times L} \mid \mathcal{D}_{ij} = \epsilon_i \delta_{ij}$. In this context, the A -matrix acts on the fermionic operator as a Bogoliubov transformation, allowing for (24) to be rewritten as

²The λ_j -term frequently appears in many condensed matter models, with different numerical values and interpretations, eg.

- In the XX model, $\lambda_j = \lambda \ \forall j$.
- While for the Anderson model $\lambda_j \in \mathcal{U}_{\mathbb{R}[-W, W]}$, a uniform random variable, with W being the disorder strength.
- In the Aubry-André model, $\lambda_j = \lambda \cos(2\pi\sigma j)$, with $\sigma \in \mathbb{I}$ and λ quantifying the disorder strength.

$$(27) \quad \mathbf{H} = \mathbf{f}^\dagger A \mathcal{D} A^T \mathbf{f} = \mathbf{d}^\dagger \mathcal{D} \mathbf{d}$$

where $\mathbf{d} = A^T \mathbf{f}$. Since the A -matrix is orthogonal, the new d_k -operators are fermionic operators as well, satisfying (24)'s anti-commutation rules. Then, the new fermionic operators are

$$\begin{aligned} d_k &= \sum_{j=1}^L A_{jk} f_j \\ f_j &= \sum_{k=1}^L A_{jk} d_k \end{aligned} \quad \text{since } A^T A = \sum_{j,k=1}^L A_{jk} A_{kj} = \mathbb{1}_L.$$

Then, we can expand (27) in terms of the lattices, as follows

$$(28) \quad \mathbf{H} = \sum_{k=1}^L \epsilon_k d_k^\dagger d_k,$$

which is a sum of number operators with potentials. The eigenstates can then be constructed from the theory's vacuum state, by applying the d_k^\dagger -fermionic operators. In the Heisenberg-picture, the d_k -operators can be evolved via the Heisenberg equation of motion

$$(29) \quad \frac{d}{dt} d_k = i[\mathbf{H}, d_k],$$

and using that $d_k^2 = 0$, it turns out that (29)'s solution is simply $d_k(t) = e^{-i\epsilon_k t} d_k$. The system's correlation can be easily found by analyzing the following matrix. Let $\mathcal{N}_{jk} = \langle d_j^\dagger d_k \rangle$, where the expectation value is taken via calculating the operator's trace along the Fock space, which takes the following values

$$(30) \quad \mathcal{N}_{jk} = \langle d_j^\dagger d_k \rangle = \begin{cases} 0 \text{ or } 1 & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases},$$

ie. different lattice-sites are not correlated and there can only be a single fermion at most per lattice site, in accordance with Pauli's principle. A ground state, for example, would choose to turn on all fermions in the eigenmode d -space such that $\epsilon_k < 0$. If instead, the expectation value is taken with thermal states, the Fermi-Dirac distribution is returned,

$$(31) \quad \mathcal{N}_{jk} = \langle d_j^\dagger d_k \rangle_{\text{th}} = \frac{1}{1 + e^{\beta\epsilon_k + \mu}} \delta_{jk}.$$

Another interesting quality is a system with an initial configuration where the system's initial state, in real space, is known. In this setting, \mathcal{N}_{jk} is known for all lattices. Consider for example the Anderson model, where the system's initial state is given by a single tensor product of n -fermionic states in real space, with $n < L$. Then, for all lattice sites, we have that \mathcal{N}_{jj} is either zero or one. The \mathcal{N}_{jk} -matrix entries can then be evaluated as

$$\langle d_j^\dagger d_k \rangle = \sum_{i,j=1}^{n < L} A_{ik} A_{jl} \langle f_i^\dagger f_j \rangle = \sum_{j=1}^n A_{jk} A_{jl} \langle f_j^\dagger f_j \rangle,$$

which can then be numerically computed to obtain the LHS expectation value. In general, this \mathcal{N}_{jk} -matrix will not be diagonal, which is reasonable since the system's real configuration is not an eigenstate. In principle and in practice, by inverting (32), we can evolve any number operator or two-body correlation operator, ie.

$$(32) \quad \langle f_j^\dagger f_k \rangle = \sum_{k,l=1}^n A_{jk} A_{jl} \langle d_k^\dagger d_l \rangle.$$

This quantities' time evolution can then be found out to be

$$(33) \quad \langle f_j^\dagger(t) f_k(t) \rangle = \sum_{k,l=1}^L e^{i(\epsilon_k - \epsilon_l)t} A_{jk} A_{jl} \langle d_k^\dagger d_l \rangle,$$

which can then be numerically solved.

3. INDEPENDENT ELECTRONS AND STATIC CRYSTALS

3.1. Crystal lattices. The mathematical concept which best describes an actual crystal lattice is that of a Bravais lattice, defined as a set of mathematical points corresponding to the discrete positions in space given by

$$(34) \quad \{\mathbf{R} \in \mathbb{R}^3 \mid \mathbf{R} = \sum_{i=1}^3 n_i \mathbf{a}_i \text{ with } n_i \in \mathbb{Z}\},$$

where the $\{\mathbf{a}_i\}$ -vector are the so-called primitive vector. The Bravais lattice is invariant under the operation

$$\mathbf{R} \rightarrow \mathbf{R} + \mathbf{T} \text{ where } \mathbf{T} = \sum_{i=1}^3 L_i \mathbf{a}_i \mid L_i \in \mathbb{Z}$$

4. HEISENBERG XXX-CHAIN

Consider a one-dimensional lattice with N lattice sites and a $\frac{1}{2}$ -spin particle positioned at every lattice site, which have a nearest neighbour spin-spin interaction. Each particles either has spin up or spin down, generating a two-dimensional local Hilbert space \mathbb{V}_n . Strictly speaking there are two possible topologies for a one-dimensional chain, open or closed. Consider the XXX-Hamiltonian with closed topology, given by

$$(35) \quad \begin{aligned} \mathbf{H} &= J \sum_{n=1}^N \left(\vec{\mathbf{S}}_n \cdot \vec{\mathbf{S}}_{n+1} - \frac{1}{4} \mathbb{1}^{\otimes N} \right) \\ &= \frac{J}{4} \sum_{n=1}^N \left(\vec{\sigma}_n \cdot \vec{\sigma}_{n+1} - \frac{1}{4} \mathbb{1}^{\otimes N} \right). \end{aligned}$$

Note that $J < 0$ models a ferromagnetic state while $J > 0$ models an anti-ferromagnetic state. A ferromagnetic state refers to a state where all the spins are aligned while an anti-ferromagnetic state refers to a state where adjacent spins are anti-aligned within the domain.

5. BETHE ANSATZ

The main point of the Bethe solution is that XXX Heisenberg Hamiltonian, which defines the model, can be expressed in terms of a complex-valued \mathcal{R} -matrix which is a solution of the Yang-Baxter equation, thus leading to the model's integrability. The most convenient approach is to construct a transfer \mathcal{T} -matrix -a one parameter commutative family of operators acting on the full state space of the Heisenberg spin chain.

The Lax operator. Consider an XXX spin chain with N sites and a corresponding Hilbert space given by

$$(36) \quad \mathbb{H} \simeq \bigotimes_{n=1}^N \mathbb{V}_n \text{ where } \mathbb{V}_n \simeq \mathbb{C}^2 \quad \text{To these individual Hilbert spaces, additional auxiliary, non-physical, spaces } \mathbb{V}_a \simeq \mathbb{C}^2, \text{ can be added.}$$

The algebraic Bethe ansatz' basic tool is the lax operator \mathcal{L} , whose definition is as follows.

Definition 1. The lax operator \mathcal{L} can be defined as an operator which involves the local quantum space \mathbb{V}_n and the auxiliary space \mathbb{V}_a , as follows

$$(37) \quad \begin{aligned} \mathcal{L} : \mathbb{V}_n \otimes \mathbb{V}_a &\rightarrow \mathbb{V}_n \otimes \mathbb{V}_a && \text{where } \mathbb{1}_n \text{ and } \mathbf{S}_n^\alpha \text{ act on } \mathbb{V}_n \\ \mathcal{L}_{n,a} &= u(\mathbb{1}_n \otimes \mathbb{1}_a) + i \sum_{\alpha} \mathbf{S}_n^\alpha \otimes \sigma_a^\alpha, \quad \forall n \leq N && \text{while } \mathbb{1}_a \text{ and } \mathbf{S}_a^\alpha \text{ act upon } \mathbb{V}_a \text{ and where the} \\ &&& \text{complex-valued } u\text{-parameter is the spectral parameter.} \end{aligned}$$

Note that [equation \(37\)](#) can be rewritten as

$$(38) \quad \begin{aligned} \mathcal{L}_{n,a} &= u(\mathbb{1}_n \otimes \mathbb{1}_a) + i \sum_{\alpha} \mathbf{S}_n^\alpha \otimes \sigma_a^\alpha \\ &= u \begin{pmatrix} \mathbb{1}_n & 0 \\ 0 & \mathbb{1}_a \end{pmatrix} + i \mathbf{S}_n^x \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}_a + i \mathbf{S}_n^y \otimes \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}_a + i \mathbf{S}_n^z \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}_a \\ &= \begin{pmatrix} u\mathbb{1}_n + i\mathbf{S}_n^z & i\mathbf{S}_n^x + \mathbf{S}_n^y \\ i\mathbf{S}_n^x - \mathbf{S}_n^y & u\mathbb{1}_n - i\mathbf{S}_n^z \end{pmatrix}_a = \begin{pmatrix} u\mathbb{1}_n + i\mathbf{S}_n^z & i\mathbf{S}_n^- \\ i\mathbf{S}_n^+ & u\mathbb{1}_n - i\mathbf{S}_n^z \end{pmatrix}_a, \end{aligned}$$

which is a 2×2 -matrix in the auxiliary space \mathbb{V}_a and the matrix entries are operators acting on the physical Hilbert space \mathbb{V}_n .

Consider now the following mathematical object.

Definition 2. Let the permutation operator be

$$(39) \quad \mathcal{P}_{n,a} = \frac{1}{2} \left(\mathbb{1}_n \otimes \mathbb{1}_a + \sum_{\alpha} \sigma_n^{\alpha} \otimes \sigma_n^{\alpha} \right), = \frac{1}{2} \sigma^{\mu} \sigma_{\mu}$$

which³ acts on both the physical and auxiliary systems.

Using this definition for the permutation operator, the lax operator can then be rewritten as

$$(41) \quad \begin{aligned} \mathcal{L}_{n,a} &= u(\mathbb{1}_n \otimes \mathbb{1}_a) + i \sum_{\alpha} \mathbf{S}_n^{\alpha} \otimes \sigma_a^{\alpha} = u(\mathbb{1}_n \otimes \mathbb{1}_a) + \frac{i}{2} \sum_{\alpha} \sigma_n^{\alpha} \otimes \sigma_a^{\alpha} \\ &= u(\mathbb{1}_n \otimes \mathbb{1}_a) - \frac{i}{2} (\mathbb{1}_n \otimes \mathbb{1}_a) + \frac{i}{2} (\mathbb{1}_n \otimes \mathbb{1}_a) + \frac{i}{2} \sum_{\alpha} \sigma_n^{\alpha} \otimes \sigma_a^{\alpha} \\ &= \left(u - \frac{i}{2} \right) (\mathbb{1}_n \otimes \mathbb{1}_a) + \frac{i}{2} \left(\mathbb{1}_n \otimes \mathbb{1}_a + \sum_{\alpha} \sigma_n^{\alpha} \otimes \sigma_n^{\alpha} \right) \\ &= \left(u - \frac{i}{2} \right) \mathbb{1}_{n,a} + i \mathcal{P}_{n,a}. \end{aligned}$$

The permutation operator indeed permutes the factors in the tensor product the factors in the tensor product by calculating the sum of tensor product of the Pauli matrices with respect to the standard four-dimensional basis, ie. if $\mathbf{x} = \begin{pmatrix} a \\ b \end{pmatrix}$ and $\mathbf{y} = \begin{pmatrix} c \\ d \end{pmatrix}$ then $\mathbf{x} \otimes \mathbf{y} = \begin{pmatrix} ac & ad \\ bc & bd \end{pmatrix}$ and $\mathbf{y} \otimes \mathbf{x} = \begin{pmatrix} ac & bc \\ ad & bd \end{pmatrix}$. Therefore,

$$(42) \quad \mathcal{P}_{n,a}(\mathbf{x} \otimes \mathbf{y}) = \mathbf{y} \otimes \mathbf{x} \text{ and } \mathcal{P}_{n,a}(\mathbf{y} \otimes \mathbf{x}) = \mathbf{x} \otimes \mathbf{y}.$$

Fundamental commutation Relations and Monodromy Matrix. Consider the following two lax operators

$$\begin{aligned} \mathcal{L}_{n,a_1}(u_1) : \mathbb{V}_n \otimes \mathbb{V}_{a_1} &\rightarrow \mathbb{V}_n \otimes \mathbb{V}_{a_1} \\ \mathcal{L}_{n,a_2}(u_2) : \mathbb{V}_n \otimes \mathbb{V}_{a_2} &\rightarrow \mathbb{V}_n \otimes \mathbb{V}_{a_2} \end{aligned} \quad \begin{array}{l} \text{both of which act on the physical quantum} \\ \text{state and on two different auxiliary spaces as well.} \end{array}$$

The product of these two operator is then a triple tensor product in $\mathbb{V}_n \otimes \mathbb{V}_{a_1} \otimes \mathbb{V}_{a_2}$. Now, the following matrix arises

Definition 3. Consider an \mathcal{R} -matrix which relates the permutation and spectral parameters of two auxiliary spaces, which is given by

$$(43) \quad \mathcal{R} : \mathbb{V}_{a_1} \otimes \mathbb{V}_{a_2} \otimes \mathbb{V}_{a_1} \otimes \mathbb{V}_{a_2}$$

$$\mathcal{R}_{a_1,a_2}(u_1 - u_2) = (u_1 - u_2) \mathbb{1}_{a_1,a_2} + i \mathcal{P}_{a_1,a_2},$$

which⁴ doesn't act on the physical system at all, but on the auxiliary, phantom, systems.

³Note that the permutation operator can be written as a 2×2 -matrix in the auxiliary space \mathbb{V}_a ,

$$(40) \quad \begin{aligned} \mathcal{P}_{n,a} &= \frac{1}{2} \left[\begin{pmatrix} \mathbb{1}_n & 0 \\ 0 & \mathbb{1}_a \end{pmatrix} + \sigma_n^x \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}_a + \sigma_n^y \otimes \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}_a + \sigma_n^z \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}_a \right] \\ &= \begin{pmatrix} \mathbb{1}_n + \sigma_n^z & \sigma_n^x - i\sigma_n^y \\ \sigma_n^x + i\sigma_n^y & \mathbb{1}_n - \sigma_n^z \end{pmatrix}_a = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \end{aligned}$$

⁴Note that \mathcal{R} can be rewritten as a 2×2 -matrix in either of the auxiliary spaces, as follows

An interesting relationship between the products of the two lax operators and the \mathcal{R} -matrix can then be found⁵, consider

$$\begin{aligned}
& \mathcal{R}_{a_1, a_2}(u_1 - u_2) \mathcal{L}_{n, a_1}(u_1) \mathcal{L}_{n, a_2}(u_2) \\
&= \left((u_1 - u_2) \mathbb{1}_{a_1, a_2} + i \mathcal{P}_{a_1, a_2} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} + i \mathcal{P}_{n, a_1} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right) \\
&= \left((u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_1} + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} + i^2 \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} \right) \\
&\quad \times \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right) \\
&= (u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&\quad + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&\quad + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&\quad - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&= (u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} \times \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_2} \times \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \\
&\quad + (u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} \times i \mathcal{P}_{n, a_1} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_2} \times i \mathcal{P}_{n, a_1} \\
&\quad + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&\quad - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&= \left((u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_2} \right) \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \\
&\quad + \left((u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_2} \right) \times i \mathcal{P}_{n, a_1} \\
&\quad + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&\quad - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} i \mathcal{P}_{n, a_2}
\end{aligned}$$

$$(44) \quad \mathcal{R}_{a_1, a_2} = \begin{pmatrix} \left(u + \frac{i}{2} \right) \mathbb{1}_{a_2} + i \mathbf{S}_{a_2}^z & i \mathbf{S}_{a_2}^- \\ i \mathbf{S}_{a_2}^+ & \left(u + \frac{i}{2} \right) \mathbb{1}_{a_2} - i \mathbf{S}_{a_2}^z \end{pmatrix}_{a_1} = \begin{pmatrix} \left(u + \frac{i}{2} \right) \mathbb{1}_{a_1} + i \mathbf{S}_{a_1}^z & i \mathbf{S}_{a_1}^- \\ i \mathbf{S}_{a_1}^+ & \left(u + \frac{i}{2} \right) \mathbb{1}_{a_1} - i \mathbf{S}_{a_1}^z \end{pmatrix}_{a_2}.$$

⁵Note that the following permutation properties hold

$$(45) \quad \mathcal{P}_{n, a_1} \mathcal{P}_{n, a_2} = \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} = \mathcal{P}_{n, a_2} \mathcal{P}_{a_2, a_1} \text{ and } \mathcal{P}_{a, b} = \mathcal{P}_{b, a}$$

$$\begin{aligned}
&= \left((u_1 - u_2) \mathbb{1}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + (u_1 - u_2) \mathbb{1}_{a_1, a_2} i \mathcal{P}_{n, a_2} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_1} \right) \\
&+ i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{a_1, a_2} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&- \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} \left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} - \mathcal{P}_{a_1, a_2} \mathcal{P}_{n, a_1} i \mathcal{P}_{n, a_2} \\
&= \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_1} \right) (u_1 - u_2) \mathbb{1}_{a_1, a_2} \\
&+ \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_1} \right) i \mathcal{P}_{n, a_2} \\
&= \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_1} \right) \left(\left(u - \frac{i}{2} \right) \mathbb{1}_{n, a_2} + i \mathcal{P}_{n, a_2} \right),
\end{aligned}$$

which proves that

$$(46) \quad \mathcal{R}_{a_1, a_2}(u_1 - u_2) \mathcal{L}_{n, a_1}(u_1) \mathcal{L}_{n, a_2}(u_2) = \mathcal{L}_{n, a_1}(u_2) \mathcal{L}_{n, a_1}(u_1) \mathcal{R}_{a_1, a_2}(u_1 - u_2).$$

From the fundamental commutation relation it can then be shown that the \mathcal{R} -matrix follows the quantum Yang-Baxter equation. Consider then the product of the following \mathcal{L} -operators,

$$\begin{aligned}
(47) \quad \mathcal{L}_{n, 1} \mathcal{L}_{n, 2} \mathcal{L}_{n, 3} &= \mathcal{R}_{12}^{-1} \mathcal{L}_{n, 2} \mathcal{L}_{n, 1} \mathcal{L}_{n, 3} \mathcal{R}_{12} \\
&= \mathcal{R}_{12}^{-1} \mathcal{R}_{13}^{-1} \mathcal{L}_{n, 2} \mathcal{L}_{n, 3} \mathcal{L}_{n, 1} \mathcal{R}_{13} \mathcal{R}_{12} \\
&= \mathcal{R}_{12}^{-1} \mathcal{R}_{13}^{-1} \mathcal{R}_{23}^{-1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 2} \mathcal{L}_{n, 1} \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12} \\
&= (\mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12})^{-1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 2} \mathcal{L}_{n, 1} \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12},
\end{aligned}$$

and similarly

$$\begin{aligned}
(48) \quad \mathcal{L}_{n, 1} \mathcal{L}_{n, 2} \mathcal{L}_{n, 3} &= \mathcal{R}_{23}^{-1} \mathcal{L}_{n, 1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 2} \mathcal{R}_{23} \\
&= \mathcal{R}_{23}^{-1} \mathcal{R}_{13}^{-1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 1} \mathcal{L}_{n, 2} \mathcal{R}_{13} \mathcal{R}_{23} \\
&= \mathcal{R}_{23}^{-1} \mathcal{R}_{13}^{-1} \mathcal{R}_{12}^{-1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 2} \mathcal{L}_{n, 1} \mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23} \\
&= (\mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23})^{-1} \mathcal{L}_{n, 3} \mathcal{L}_{n, 2} \mathcal{L}_{n, 1} \mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23},
\end{aligned}$$

both of which then yield an important constraint on the \mathcal{R} -matrix, namely

$$(49) \quad \mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23} = \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12},$$

which is the quantum Yang-Baxter equation.

Definition 4. Let the monodromy matrix $\mathcal{T}_{N, a}$ be defined as a product of subsequent of \mathcal{L} -operators, ie.

$$\begin{aligned}
(50) \quad \mathcal{T}_{N, a} &= \prod_1^{j=N} \mathcal{L}_{j, a} \\
\mathcal{T}_{N, a} &= \prod_1^{j=N} \begin{pmatrix} u \mathbb{1}_j + i \mathcal{S}_j^z & i \mathcal{S}_j^- \\ i \mathcal{S}_j^+ & u \mathbb{1}_j - i \mathcal{S}_j^z \end{pmatrix} \equiv \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix},
\end{aligned}$$

This matrix obeys the fundamental relation as well⁶, ie.

$$(52) \quad \mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{T}_{N,a_2}(u_2) = \mathcal{T}_{N,a_2}(u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{R}_{a_1,a_2}(u_1 - u_2).$$

Definition 5. The transfer \mathfrak{t} -matrix is defined as the \mathcal{T} -matrix's partial trace over the auxiliary space \mathbb{V}_a , ie.

$$(53) \quad \mathfrak{t}(u) = \text{Tr}_{\mathbb{V}_a} \mathcal{T}_{N,a}(u) = \text{Tr}_{\mathbb{V}_a} \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix} = A(u) + D(u),$$

where \mathfrak{t}, A, B, C, D all are 2×2 -matrices in the physical system, the auxiliary systems being present no more.

With the previous definitions in mind, **equation (52)**'s double trace can then be written in terms of the transfer matrix, as follows

$$\begin{aligned} & \text{Tr}_{\mathbb{V}_{a_1}} \text{Tr}_{\mathbb{V}_{a_2}} \left[\mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{T}_{N,a_2}(u_2) \right] \\ & \Rightarrow \text{Tr}_{\mathbb{V}_{a_1}} \text{Tr}_{\mathbb{V}_{a_2}} \left[\mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{T}_{N,a_2}(u_2) - \mathcal{T}_{N,a_1}(u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{R}_{a_1,a_2}(u_1 - u_2) \right] = 0 \\ & \Rightarrow \text{Tr}_{\mathbb{V}_{a_1}} \text{Tr}_{\mathbb{V}_{a_2}} \left[\mathcal{R}_{a_1,a_2}(u_1 - u_2) \left(\mathcal{T}_{N,a_1}(u_1) \mathcal{T}_{N,a_2}(u_2) - \mathcal{T}_{N,a_2}(u_2) \mathcal{T}_{N,a_1}(u_1) \right) \right] = 0 \end{aligned}$$

which implies $\mathfrak{t}(u_1)\mathfrak{t}(u_2) = \mathfrak{t}(u_2)\mathfrak{t}(u_1)$.

Therefore the transfer matrix commutes with itself for different values of the spectral parameter. It turns out that the monodromy matrix can be expanded in a power series around any point $z_0 \in \mathbb{C}$, generating an infinite set of linearly independent commuting operators acting on the full quantum space. This formally leads to the model's integrability.

Monodromy matrix and the Hamiltonian. The Hamiltonian operator belongs to the family of transfer matrices. Let $z_0 = \frac{i}{2}$, then the monodromy matrix can be expanded around said complex point yielding (up to first order)

$$\begin{aligned} \mathcal{T}_{N,a}(z_0) &= \prod_1^{j=N} \left[\left(\cancel{z_0} - \frac{i}{2} \right) \mathbb{1}_{j,a} + i \mathcal{P}_{j,a} \right] = i^N \mathcal{P}_{N,a} \mathcal{P}_{N-1,a} \cdots \mathcal{P}_{1,a} = \\ (54) \quad &= i^N \mathcal{P}_{1,2} \mathcal{P}_{2,3} \cdots \mathcal{P}_{N-1,N} \mathcal{P}_{N,a} \\ &= i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \mathcal{P}_{N,a} \end{aligned}$$

⁶For example, consider $N = 2$, then

$$\begin{aligned} \mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{T}_{N,a_1}(u_1) \mathcal{T}_{N,a_2}(u_2) &= \mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{T}_{2,a_1}(u_1) \mathcal{T}_{2,a_2}(u_2) \\ &= \mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{L}_{2,a_1}(u_1) \mathcal{L}_{2,a_1}(u_1) \mathcal{L}_{2,a_2}(u_2) \mathcal{L}_{1,a_2}(u_2) \\ &= \left(\mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{L}_{2,a_1}(u_1) \mathcal{L}_{2,a_2}(u_2) \right) \mathcal{L}_{1,a_1}(u_1) \mathcal{L}_{1,a_2}(u_2) \\ (51) \quad &= \mathcal{L}_{2,a_2}(u_2) \mathcal{L}_{2,a_1}(u_1) \left(\mathcal{R}_{a_1,a_2}(u_1 - u_2) \mathcal{L}_{1,a_1}(u_1) \mathcal{L}_{1,a_2}(u_2) \right) \\ &= \mathcal{L}_{2,a_2}(u_2) \mathcal{L}_{2,a_1}(u_1) \mathcal{L}_{1,a_2}(u_2) \mathcal{L}_{1,a_1}(u_1) \mathcal{R}_{a_1,a_2}(u_1 - u_2) \\ &= \mathcal{L}_{2,a_2}(u_2) \mathcal{L}_{1,a_2}(u_2) \mathcal{L}_{2,a_1}(u_1) \mathcal{L}_{1,a_1}(u_1) \mathcal{R}_{a_1,a_2}(u_1 - u_2) \\ &= \mathcal{T}_{n,a_1}(u_2) \mathcal{T}_{n,a_1}(u_1) \mathcal{R}_{a_1,a_2}(u_1 - u_2), \end{aligned}$$

where the fundamental commutation relation given by (46) was used. Note as well that \mathcal{L} -operators acting on different spaces commute as well, eg. $[\mathcal{L}_{n_1,a_1}, \mathcal{L}_{n_2,a_2}] \propto \delta_{n_1,n_2}$. The general fundamental relation can then be proved via mathematical induction for the general case.

where the second line holds given the properties of the permutation operators. Taking the partial trace over the auxiliary space yields the transfer matrix, as follows

$$\begin{aligned}
 \mathfrak{t}(z_0) &= \text{Tr}_{\mathbb{V}_a} \mathcal{T}_{N,a}(z_0) = \text{Tr}_{\mathbb{V}_a} i^N \mathcal{P}_{1,2} i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \mathcal{P}_{N,a} \\
 (55) \quad &= i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \text{Tr}_{\mathbb{V}_a} \mathcal{P}_{N,a} = i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \text{Tr}_{\mathbb{V}_a} \frac{1}{2} \begin{pmatrix} \mathbb{1}_n + \sigma_n^z & \sigma_n^x - i\sigma_n^y \\ \sigma_n^x + i\sigma_n^y & \mathbb{1}_n - \sigma_n^z \end{pmatrix}_a \\
 &= i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \mathbb{1}_N.
 \end{aligned}$$

Then $\mathcal{U} = i^{-N} \mathfrak{t}(z_0) = \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1}$ is a shift operator in the full quantum Hilbert space \mathbb{H} . In effect, note that

$$\mathcal{P}_{n_1, n_2} \mathbf{X}_{n_2} \mathcal{P}_{n_1, n_2} = \mathbf{X}_{n_1},$$

ie. this permutation moves \mathbf{X} one step back. Furthermore, this operator is unitary (given that the permutations are, by definition, unitary operators). Then, according to Stone's theorem on one-parameter unitary groups, the shift operator is related to the momentum operator

$$(56) \quad \mathcal{U} = e^{iP}$$

The next order in the expansion of the transfer matrix can then be found via the derivative of the monodromy matrix at $z_0 = \frac{i}{2}$,

$$\begin{aligned}
 \frac{d\mathcal{T}_{N,a}(u)}{du} \Big|_{u=z_0} &= \frac{d}{du} \prod_1^{j=N} \begin{pmatrix} u\mathbb{1}_j + i\mathbf{S}_j^z & i\mathbf{S}_j^- \\ i\mathbf{S}_j^+ & u\mathbb{1}_j - i\mathbf{S}_j^z \end{pmatrix} \Big|_{u=z_0} \\
 &= \frac{d}{du} \left[\begin{pmatrix} u\mathbb{1}_N + i\mathbf{S}_N^z & i\mathbf{S}_N^- \\ i\mathbf{S}_N^+ & u\mathbb{1}_N - i\mathbf{S}_N^z \end{pmatrix} \right] \Big|_{u=z_0} \prod_1^{j=N-1} \begin{pmatrix} u\mathbb{1}_j + i\mathbf{S}_j^z & i\mathbf{S}_j^- \\ i\mathbf{S}_j^+ & u\mathbb{1}_j - i\mathbf{S}_j^z \end{pmatrix} \\
 &\quad + \begin{pmatrix} u\mathbb{1}_N + i\mathbf{S}_N^z & i\mathbf{S}_N^- \\ i\mathbf{S}_N^+ & u\mathbb{1}_N - i\mathbf{S}_N^z \end{pmatrix} \frac{d}{du} \left[\begin{pmatrix} u\mathbb{1}_{N-1} + i\mathbf{S}_{N-1}^z & i\mathbf{S}_{N-1}^- \\ i\mathbf{S}_{N-1}^+ & u\mathbb{1}_{N-1} - i\mathbf{S}_{N-1}^z \end{pmatrix} \right] \Big|_{u=z_0} \\
 &\quad \times \prod_1^{j=N-2} \begin{pmatrix} u\mathbb{1}_j + i\mathbf{S}_j^z & i\mathbf{S}_j^- \\ i\mathbf{S}_j^+ & u\mathbb{1}_j - i\mathbf{S}_j^z \end{pmatrix} \\
 &\quad + \cdots + \prod_3^{j=N} \begin{pmatrix} u\mathbb{1}_j + i\mathbf{S}_j^z & i\mathbf{S}_j^- \\ i\mathbf{S}_j^+ & u\mathbb{1}_j - i\mathbf{S}_j^z \end{pmatrix} \frac{d}{du} \left[\begin{pmatrix} u\mathbb{1}_2 + i\mathbf{S}_2^z & i\mathbf{S}_2^- \\ i\mathbf{S}_2^+ & u\mathbb{1}_2 - i\mathbf{S}_2^z \end{pmatrix} \right] \Big|_{u=z_0} \\
 &\quad \times \begin{pmatrix} u\mathbb{1}_1 + i\mathbf{S}_1^z & i\mathbf{S}_1^- \\ i\mathbf{S}_1^+ & u\mathbb{1}_1 - i\mathbf{S}_1^z \end{pmatrix} \\
 &\quad + \prod_2^{j=N} \begin{pmatrix} u\mathbb{1}_j + i\mathbf{S}_j^z & i\mathbf{S}_j^- \\ i\mathbf{S}_j^+ & u\mathbb{1}_j - i\mathbf{S}_j^z \end{pmatrix} \frac{d}{du} \left[\begin{pmatrix} u\mathbb{1}_1 + i\mathbf{S}_1^z & i\mathbf{S}_1^- \\ i\mathbf{S}_1^+ & u\mathbb{1}_1 - i\mathbf{S}_1^z \end{pmatrix} \right] \Big|_{u=z_0} \\
 &= i^{N-1} \left[\prod_1^{j=N-1} \mathcal{P}_j + \prod_N^{j=N} \mathcal{P}_j \times \prod_1^{j=N-2} \mathcal{P}_j + \cdots + \prod_k^{j=N} \mathcal{P}_j \times \prod_1^{k+2} \mathcal{P}_j \right. \\
 &\quad \left. + \cdots + \prod_3^{j=N} \mathcal{P}_j \times \prod_1^{j=1} \mathcal{P}_j + \prod_2^{j=N} \mathcal{P}_j \right] \\
 &= i^{N-1} \sum_{n \in \mathbb{N}} \mathcal{P}_{N,a} \cdots \mathcal{P}_{n-1,a} \mathcal{P}_{n+1,a} \cdots \mathcal{P}_{1,a}
 \end{aligned}$$

$$\begin{aligned}
&= i^{N-1} \sum_{n \in \mathbb{N}} \mathcal{P}_{1,2} \mathcal{P}_{2,3} \cdots \mathcal{P}_{n-1,n+1} \cdots \mathcal{P}_{N-1,N} \mathcal{P}_{N,a} \\
&= i^{N-1} \sum_{n \in \mathbb{N}} \prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1} \mathcal{P}_{N,a}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
(57) \quad \frac{d\mathbf{t}(u)}{du} \Big|_{u=z_0} &= \frac{d \operatorname{Tr}_{\mathbb{V}_a} \mathcal{T}_{N,a}(u)}{du} \Big|_{u=z_0} \\
&= i^{N-1} \sum_{n \in \mathbb{N}} \prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1} \times \operatorname{Tr}_{\mathbb{V}_a} \mathcal{P}_{N,a} \\
&= i^{N-1} \sum_{n \in \mathbb{N}} \prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1}.
\end{aligned}$$

Now, note that

$$\begin{aligned}
(58) \quad \frac{d}{du} \log(\mathbf{t}(u)) \Big|_{u=z_0} &= \frac{d}{du} \log \operatorname{Tr}_{\mathbb{V}_a} \mathcal{T}(u) \Big|_{u=z_0} \\
&= \frac{d}{du} \left(\mathbf{t}(u) \mathbf{t}(u)^{-1} \right) \Big|_{u=z_0} \\
&= i^{N-1} \sum_{n \in \mathbb{N}} \left(\prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1} \right) \times \left(i^N \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \mathbb{1}_N \right)^{-1} \\
&= \frac{1}{i} \sum_{n \in \mathbb{N}} \left(\prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1} \right) \times \prod_{j=1}^{N-1} \mathcal{P}_{N-1-j,N-j}^{-1} \\
&= \frac{1}{i} \sum_{n \in \mathbb{N}} \left(\prod_{j=1}^{n-2} \mathcal{P}_{j,j+1} \times \mathcal{P}_{n-1,n+1} \times \prod_{j=n+2}^{N-1} \mathcal{P}_{j,j+1} \right) \times \prod_{j=1}^{N-1} \mathcal{P}_{j,j+1} \\
&= \frac{1}{i} \sum_{n \in \mathbb{N}} \left(\cancel{\mathcal{P}_{1,2}} \cancel{\mathcal{P}_{2,3}} \cdots \mathcal{P}_{n-1,n+1} \cdots \cancel{\mathcal{P}_{N-1,N}} \times \cancel{\mathcal{P}_{N-1,N}} \cdots \mathcal{P}_{n,n+1} \cdots \cancel{\mathcal{P}_{2,3}} \cancel{\mathcal{P}_{1,2}} \right) \\
&= \frac{1}{i} \sum_{n \in \mathbb{N}} \mathcal{P}_{N-1,N}
\end{aligned}$$

Note that (35) can be rewritten in terms of the permutation operator as

$$(59) \quad \mathbf{H} = \frac{J}{2} \sum_n \left(\mathcal{P}_{n,n+1} - \mathbb{1}^{\otimes N} \right),$$

which in turn can be rewritten as

$$(60) \quad \mathbf{H} = \frac{J}{2} \left(i \frac{d}{du} \log(\mathbf{t}(u)) \Big|_{u=z_0} - N \mathbb{1}^{\otimes N} \right),$$

which shows that the XXX Hamiltonian belongs to the family of $N - 1$ commuting operators generated by the trace of the monodromy matrix \mathcal{T} . As a result, the Hamiltonian commutes with the transfer matrix $[\mathbf{H}, \mathbf{t}(u)] = 0$.

Diagonalizing the Hamiltonian. The only task left is to diagonalize the Hamiltonian, by diagonalizing the transfer matrix. The monodromy matrix $\mathbf{t}_{N,a}$ is a 2×2 -matrix in the physical space \mathbb{V}_a . Recalling the fundamental commutation relation between the \mathcal{R} and \mathcal{T} -matrices, then the following theorem holds,

Theorem 1. *Let \mathcal{T} be the monodromy matrix for the Heisenberg model. Then, $\forall u_1, u_2 \in \mathbb{C}$ then*

- $[\mathcal{T}_{ij}(u_1), \mathcal{T}_{ij}(u_2)] = 0, i, j = 1, 2$, which entails that $[B(u_1), B(u_2)] = \dots = [D(u_1), D(u_2)] = 0$
- $A(u_1)B(u_2) = f(u_1 - u_2)B(u_2)A(u_1) + g(u_1 - u_2)B(u_1)A(u_2)$
- $D(u_1)B(u_2) = h(u_1 - u_2)B(u_2)D(u_1) + k(u_1 - u_2)B(u_1)D(u_2)$,

where f, g, h , and k are given by

$$f(u) = \frac{u-i}{u}, \quad g(u) = \frac{i}{u}, \quad h(u) = \frac{u+i}{u}, \quad k(u) = -\frac{i}{u}.$$

Proof. Consider two different monodromy matrices, labelled $\mathcal{T}_{N,a_1}(u_1)$ and $\mathcal{T}_{N,a_2}(u_2)$, acting on two different auxiliary spaces, labelled \mathbb{V}_1 and \mathbb{V}_2 respectively. Then, these \mathcal{T} -matrices can be written as

$$(61) \quad \begin{aligned} \mathcal{T}_{N,a_1}(u_1) &= \begin{pmatrix} A(u_1) & B(u_1) \\ C(u_1) & D(u_1) \end{pmatrix} \otimes \mathbf{1}_{\mathbb{V}_2}, \quad \mathcal{T}_{N,a_2}(u_2) = \mathbf{1}_{\mathbb{V}_1} \otimes \begin{pmatrix} A(u_2) & B(u_2) \\ C(u_2) & D(u_2) \end{pmatrix} \\ &= \begin{pmatrix} A(u_1) & 0 & B(u_1) & 0 \\ 0 & A(u_1) & 0 & B(u_1) \\ C(u_1) & 0 & D(u_1) & 0 \\ 0 & C(u_1) & 0 & D(u_1) \end{pmatrix}, \quad \begin{pmatrix} A(u_2) & B(u_2) & 0 & 0 \\ C(u_2) & D(u_2) & 0 & 0 \\ 0 & 0 & A(u_2) & B(u_2) \\ 0 & 0 & C(u_2) & D(u_2) \end{pmatrix} \end{aligned}$$

Recall that the \mathcal{R} -matrix is given by (43), which can be rewritten as

$$\mathcal{R}_{a_1,a_2}(u_1 - u_2) = \begin{pmatrix} a(u_1 - u_2) & 0 & 0 & 0 \\ 0 & b(u_1 - u_2) & c(u_1 - u_2) & 0 \\ 0 & c(u_1 - u_2) & b(u_1 - u_2) & 0 \\ 0 & 0 & 0 & a(u_1 - u_2) \end{pmatrix} \text{ where } \begin{cases} a(\lambda) = \lambda + i \\ b(\lambda) = \lambda \\ c(\lambda) = i \end{cases},$$

Then the fundamental commutation relation (52) establishes a relationship between the monodromy \mathcal{T} -matrices and the \mathcal{R} -matrix. Then, plugging in the previous results yields

$$\begin{aligned} & \begin{pmatrix} a_{12} & 0 & 0 & 0 \\ 0 & b_{12} & c_{12} & 0 \\ 0 & c_{12} & b_{12} & 0 \\ 0 & 0 & 0 & a_{12} \end{pmatrix} \begin{pmatrix} A(u_1)A(u_2) & A(u_1)B(u_2) & B(u_1)A(u_2) & B(u_1)B(u_2) \\ A(u_1)C(u_2) & A(u_1)D(u_2) & B(u_1)C(u_2) & B(u_1)D(u_2) \\ C(u_1)A(u_2) & C(u_1)B(u_2) & D(u_1)A(u_2) & D(u_1)B(u_2) \\ C(u_1)C(u_2) & C(u_1)D(u_2) & D(u_1)C(u_2) & D(u_1)D(u_2) \end{pmatrix} \\ &= \begin{pmatrix} A(u_2)A(u_1) & B(u_2)A(u_1) & A(u_2)B(u_1) & B(u_2)B(u_1) \\ C(u_2)A(u_1) & D(u_2)A(u_1) & C(u_2)B(u_1) & D(u_2)B(u_1) \\ A(u_2)C(u_1) & B(u_2)C(u_1) & A(u_2)D(u_1) & B(u_2)D(u_1) \\ C(u_2)C(u_1) & D(u_2)C(u_1) & C(u_2)D(u_1) & D(u_2)D(u_1) \end{pmatrix} \begin{pmatrix} a_{12} & 0 & 0 & 0 \\ 0 & b_{12} & c_{12} & 0 \\ 0 & c_{12} & b_{12} & 0 \\ 0 & 0 & 0 & a_{12} \end{pmatrix} \\ \Rightarrow & \begin{pmatrix} a_{12}A_1A_2 & a_{12}A_1B_2 & a_{12}B_1A_2 & a_{12}B_1B_2 \\ b_{12}A_1C_2 + c_{12}C_1A_2 & b_{12}A_1D_2 + c_{12}C_1B_2 & b_{12}B_1C_2 + c_{12}D_1A_2 & b_{12}B_1D_2 + c_{12}D_1B_2 \\ c_{12}A_1C_2 + b_{12}C_1A_2 & b_{12}A_1D_2 + c_{12}C_1B_2 & c_{12}B_1C_2 + b_{12}D_1A_2 & c_{12}B_1D_2 + b_{12}D_1B_2 \\ a_{12}C_1C_2 & a_{12}C_1D_2 & a_{12}D_1C_2 & a_{12}D_1D_2 \end{pmatrix} \\ &= \begin{pmatrix} a_{12}A_2A_1 & b_{12}B_2A_1 + c_{12}A_2B_1 & c_{12}B_2A_1 + b_{12}A_2B_1 & a_{12}B_2B_1 \\ a_{12}C_2A_1 & b_{12}D_2A_1 + c_{12}C_2B_1 & c_{12}D_2A_1 + b_{12}C_2B_1 & a_{12}D_2B_1 \\ a_{12}A_2C_1 & b_{12}B_2C_1 + c_{12}A_2D_1 & c_{12}B_2C_1 + b_{12}A_2D_1 & a_{12}B_2D_1 \\ a_{12}C_2C_1 & b_{12}D_2C_1 + c_{12}C_2D_1 & c_{12}D_2C_1 + b_{12}C_2D_1 & a_{12}D_2D_1 \end{pmatrix} \end{aligned}$$

$$\Rightarrow \begin{pmatrix} a_{12}[A_1, A_2] & a_{12}A_1B_2 - (b_{12}B_2A_1 + c_{12}A_2B_1) & a_{12}B_1A_2 - (c_{12}B_2A_1 + c_{12}A_2B_1) & a_{12}[B_1, B_2] \\ (b_{12}A_1C_2 + c_{12}C_1A_2) - a_{12}C_2A_1 & b_{12}[A_1, D_2] + c_{12}[C_1B_2] & (b_{12}[B_1, C_2] + c_{12}[D_1, A_2]) & b_{12}B_1D_2 + c_{12}D_1B_2 - a_{12}D_2B_1 \\ c_{12}A_1C_2 + b_{12}C_1A_2 - a_{12}A_2C_1 & (b_{12}A_1D_2 + c_{12}C_1B_2) - b_{12}B_2C_1 + c_{12}A_2D_1 & c_{12}[B_1, C_2] + b_{12}[D_1, A_2] & c_{12}B_1D_2 + b_{12}D_1B_2 - a_{12}B_2D_1 \\ a_{12}[C_1, C_2] & a_{12}C_1D_2 - (b_{12}D_2C_1 + c_{12}C_2D_1) & a_{12}D_1C_2 - (c_{12}D_2C_1 + b_{12}C_2D_1) & a_{12}[D_1, D_2] \end{pmatrix} = \mathbf{0},$$

from which, it's been proven that $[\mathcal{T}_{ij}(u_1), \mathcal{T}_{ij}(u_2)] = 0$, $i, j = 1, 2$ as long as $a(u_1 - u_2) \neq 0$. From the other matrix-entries, it's immediately seen that

$$a(u_1 - u_2)B(u_1)A(u_2) = c(u_1 - u_2)B(u_2)A(u_1) + b(u_1 - u_2)A(u_2)B(u_1).$$

Swapping $u_1 \rightarrow u_2$ implies $u_1 - u_2 \rightarrow -(u_1 - u_2)$. Now, note that $c(u_1)/b(u_1) = g(u_1) = g(-u_1)$ and similarly $a(-u_1)/b(-u_1) = h(-u_1) \equiv f(u_1)$, which yields the second line of theorem 1. The third claim can then be similarly derived.

quod
erat
dem■

Eigenvalues of the Bethe Ansatz. Theorem 1's relations shows the model's inner structure, identifying an underlying algebraic structure somewhat similar to that of the Harmonic oscillator, where $A + D$ are associated with the eigenenergies, and where B and C can be interpreted as ladder operators.

Then, a Fock-like state space can be constructed for the N -body system, with its ground state satisfying $C\Omega^{\otimes N} = 0$. Then, $\omega_+^i = \begin{pmatrix} 1 & 0 \end{pmatrix}_i^\dagger$ describes i -th spin's state, with up spin. Then,

$$\mathbf{S}^+\omega_+^i = 0,$$

Therefore, consider the \mathcal{L} -matrix, defined by (41), its action on a tensor product of states can be written out as

$$(62) \quad \mathcal{L}_{N,a}v \otimes \omega_+^i \begin{pmatrix} \lambda + \frac{i}{2} & \cdots \\ 0 & \lambda - \frac{i}{2} \end{pmatrix} v \otimes \omega_+^i, \quad \forall v \in \mathbb{V}_a,$$

since \mathbf{S}^+ annihilates ω_+^i . Thus, the ground state can be written as the physical ferromagnetic vacuum state with all its spins up. Said state is an eigenstate of $A(u)$ and $D(u)$ and is annihilated by $C(u)$, ie.

$$(63) \quad \Omega^{\otimes N} = \bigotimes_{i=1}^N \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \text{ where } \begin{aligned} A(u)\Omega^{\otimes N} &= \left(u + \frac{i}{2}\right)^N \Omega^{\otimes N}, \\ C(u)\Omega^{\otimes N} &= 0, \\ D(u)\Omega^{\otimes N} &= \left(u - \frac{i}{2}\right)^N \Omega^{\otimes N}, \end{aligned}$$

The $B(u)$ operators can then be used to construct the Bethe state

$$(64) \quad |u_1, \dots, u_M\rangle = \prod_{j=1}^{M < N} B(u_j)\Omega^{\otimes N}, \text{ for some set } \{u_j\}_{j=1}^{M < N} \subset \mathbb{C}.$$

A priori, not all complex-valued sequences are allowed since the Bethe states must be eigenvectors of $A(u) + D(u)$, which yields some algebraic conditions.

Conditions on $\{u_j\}_{j=1}^{M < N}$. Consider the Bethe state given by equation (64). Then,

$$A(v)|u_1, \dots, u_M\rangle = A(v) \prod_{j=1}^{M < N} B(u_j)\Omega^{\otimes N}.$$

Using the fundamental commutation relations given theorem 1,

$$A(v)B(u) = f(v - u)B(u)A(v) + g(v - u)B(v)A(u).$$

Therefore,

$$\begin{aligned}
 (65) \quad A(v) |u_1, \dots, u_M\rangle &= \left[f(v - u_1)B(u_1)A(v) + g(v - u_1)B(v)A(u_1) \right] \prod_{j=2}^{M < N} B(u_j) \Omega^{\otimes N} \\
 &= \left[\prod_{k=1}^{\ell} f(v - u_k) \right] \alpha^N(v) |u_1, \dots, u_M\rangle + \mathcal{O}(A \times D),
 \end{aligned}$$

wherein, the first term is a scalar times the Bethe state, said scalar being the eigenvalue if and only if the second term, the term with products of A and D , cancels out. In that case, the Bethe state is an eigenvector of $A + D$. Then, continuing the previous calculation

$$\begin{aligned}
 (66) \quad A(v) |u_1, \dots, u_M\rangle &= \left[f(v - u_1)B(u_1)A(v) + g(v - u_1)B(v)A(u_1) \right] B(u_2) \prod_{j=3}^{M < N} B(u_j) \Omega^{\otimes N} \\
 &= \left[f(v - u_1)f(v - u_2)B(u_1)B(u_2)A(v) + f(v - u_1)g(v - u_2)B(u_1)B(v)A(u_2) \right. \\
 &\quad \left. + g(v - u_1)f(u_1 - u_2)B(v)B(u_2)A(u_1) + g(v - u_1)g(v - u_2)B(v)B(u_2) \right] \prod_{j=3}^{M < N} B(u_j) \Omega^{\otimes N}.
 \end{aligned}$$

Now, note that

$$\begin{aligned}
 (A(v) + D(v)) |u_1, \dots, u_M\rangle &= \left[\left(\prod_{k=1}^{\ell} f(v - u_k) \right) \alpha^N(v) + \left(\prod_{k=1}^{\ell} h(v - u_k) \right) \delta^N(v) \right] |u_1, \dots, u_M\rangle \\
 &\quad + \sum_{k=1}^{\ell} \left[\left(g(v - u_k) \left[\prod_{j \neq k}^{\ell} f(u_k - u_j) \right] \alpha^N(u_k) \right) + h(v - u_k) \left(\prod_{j \neq k}^{\ell} h(u_k - u_j) \right) \delta^N(u_j) \right] \\
 &\quad \times \prod_{\substack{j=1 \\ j \neq k}}^{\ell} \Omega^{\otimes N} B(u_i) \Big] \\
 &\equiv \left[\left(\prod_{k=1}^{\ell} f(v - u_k) \right) \alpha^N(v) + \left(\prod_{k=1}^{\ell} h(v - u_k) \right) \delta^N(v) \right] |u_1, \dots, u_M\rangle + 0 \\
 &\Rightarrow \sum_{k=1}^{\ell} \left[\left(g(v - u_k) \left[\prod_{j \neq k}^{\ell} f(u_k - u_j) \right] \alpha^N(u_k) \right) + h(v - u_k) \left(\prod_{j \neq k}^{\ell} h(u_k - u_j) \right) \delta^N(u_j) \right] = 0 \\
 &\quad \sum_{k=1}^{\ell} \left[\left(g(v - u_k) \left[\prod_{j \neq k}^{\ell} f(u_k - u_j) \right] \alpha^N(u_k) \right) - g(v - u_k) \left(\prod_{j \neq k}^{\ell} h(u_k - u_j) \right) \delta^N(u_j) \right] = 0 \\
 &\quad \sum_{k=1}^{\ell} g(v - u_k) \left[\prod_{j \neq k}^{\ell} f(u_k - u_j) \right] \alpha^N(u_k) + \prod_{j \neq k}^{\ell} h(u_k - u_j) \delta^N(u_j) = 0,
 \end{aligned}$$

which must be true for all k ,

$$(67) \quad \left[\prod_{j \neq k}^{\ell} f(v - u_k) \right] \alpha^N(u_j) = \left[\prod_{j \neq k}^{\ell} h(v - u_k) \right] \delta^N(u_j).$$

Now using [theorem 1](#) functions and the that $\alpha(u) = u + \frac{i}{2}$ and $\delta(u) = u - \frac{i}{2}$, the previous equation can be rewritten as

$$\begin{aligned}
(68) \quad & \left[\prod_{j \neq k}^{\ell} f(u_k - u_j) \right] \alpha^N(u_j) = \left[\prod_{j \neq k}^{\ell} h(u_k - u_j) \right] \delta^N(u_j) \\
& \prod_{j \neq k}^{\ell} \frac{u_k - u_j - i}{u_k - u_j} \left(u_j + \frac{i}{2} \right) = \prod_{j \neq k}^{\ell} \frac{u_k - u_j + i}{u_k - u_j} \left(u_j - \frac{i}{2} \right) \\
& 0 = \prod_{j \neq k}^{\ell} \frac{u_k - u_j - i}{u_k - u_j} \left(u_j + \frac{i}{2} \right) - \prod_{j \neq k}^{\ell} \frac{u_k - u_j + i}{u_k - u_j} \left(u_j - \frac{i}{2} \right),
\end{aligned}$$

which can be rearranged to yield the Bethe ansatz equations,

$$(69) \quad \left(\frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} \right)^N = \prod_{j \neq k}^{\ell} \frac{u_k - u_j + i}{u_k - u_j - i},$$

which is a set of ℓ non linear equations on the coefficients $u_k \in \mathbb{C}$.

Momentum and Energy. Let $\{u_k\}_{k=1}^{\ell < N} \subset \mathbb{C}$ satisfy the Bethe equations. Then,

$$\begin{aligned}
(70) \quad & (A(v) + D(v)) |u_1, \dots, u_M\rangle = \left[\left(\prod_{k=1}^{\ell} f(v - u_k) \right) \alpha^N(v) + \left(\prod_{k=1}^{\ell} h(v - u_k) \right) \delta^N(v) \right] |u_1, \dots, u_M\rangle \\
& (A(v) + D(v)) |u_1, \dots, u_M\rangle = \left[\left(v + \frac{i}{2} \right)^N \prod_{k=1}^{\ell} \frac{v - u_k - i}{v - u_k} + \left(v - \frac{i}{2} \right)^N \prod_{k=1}^{\ell} \frac{v - u_k + i}{v - u_k} \right] |u_1, \dots, u_M\rangle \\
& \left(A \left(\frac{i}{2} \right) + D \left(\frac{i}{2} \right) \right) |u_1, \dots, u_M\rangle = i^N \prod_{k=1}^{\ell} \frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} |u_1, \dots, u_M\rangle \equiv \lambda(\{u_k\}_{k=1}^{\ell < N}) |u_1, \dots, u_M\rangle
\end{aligned}$$

Therefore, via the transfer matrix, it holds that $U = e^{iP} = i^{-N} \mathfrak{t} \left(\frac{i}{2} \right) = i^{-N} \left(A \left(\frac{i}{2} \right) + D \left(\frac{i}{2} \right) \right)$, from which

$$(71) \quad P |u_1, \dots, u_{\ell}\rangle = \sum_{k=1}^{\ell} p(u_k) |u_1, \dots, u_{\ell}\rangle \text{ where } p(u_k) = \frac{1}{i} \log \frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}}.$$

Now, [equation \(60\)](#) establishes that the XXX Hamiltonian belongs to the monodromy matrices' family. Furthermore, it provides a link between the $A + D$'s eigenvalues and the eigenenergies. In effect, said eigenenergies may be found by differentiating the $A + D$'s eigenvalues over u and setting $u = z_0 = \frac{i}{2}$. Let the former

$$\Lambda(v) = \left(v + \frac{i}{2} \right)^N \prod_{k=1}^{\ell} \frac{v - u_k - i}{v - u_k} + \left(v - \frac{i}{2} \right)^N \prod_{k=1}^{\ell} \frac{v - u_k + i}{v - u_k},$$

then,

$$\begin{aligned}
\frac{d \log \Lambda(v)}{dv} \Big|_{v=z_0} &= \frac{1}{\Lambda(v)} \frac{d\Lambda(v)}{dv} \Big|_{v=z_0} \\
&= \frac{1}{\Lambda(v) \Big|_{v=z_0}} \left[N i^{N-1} \prod_{k=1}^{\ell} \frac{-u_k - \frac{i}{2}}{\frac{i}{2} - u_k} + i^N \sum_k \frac{i}{(\frac{i}{2} - u_k)^2} \prod_{\substack{j=1 \\ j \neq k}}^{\ell} \frac{-u_j - \frac{i}{2}}{\frac{i}{2} - u_j} \right] \\
&= i \left(\sum_{k=1}^{\ell} \frac{1}{u_k^2 + \frac{1}{4}} - N \right).
\end{aligned}$$

Then, using [equation \(60\)](#), yields the eigenenergies

$$(72) \quad H |u_1, \dots u_M\rangle = \left[\frac{i}{2} i \left(\sum_{k=1}^{\ell} \frac{1}{u_k^2 + \frac{1}{4}} - N \right) - \frac{N}{2} \right] |u_1, \dots u_M\rangle$$

or, equivalently,

$$(73) \quad \mathbf{H} |u_1, \dots u_M\rangle = \sum_{k=1}^{\ell} \epsilon(u_k) |u_1, \dots u_M\rangle \text{ where } \epsilon(u_k) = -\frac{1}{2} \frac{1}{u_k^2 + \frac{1}{4}},$$

which concludes the Bethe solution for the XXX Heisenberg chain.

6. .