

## **Abstract**

White matter hyperintensities (WMHs) are foci of abnormal signal intensity in white matter regions seen with magnetic resonance imaging (MRI). WMHs are associated with normal aging and have shown prognostic value in neurological conditions such as traumatic brain injury (TBI). The impracticality of manually quantifying these lesions limits their clinical utility and motivates the utilization of machine learning techniques for automated segmentation workflows. Herein, we develop a concatenated random forest framework with image features for segmenting WMHs in a TBI cohort. The framework is built upon the Advanced Normalization Tools (ANTs) and ANTsR toolkits. MR (3D FLAIR, T2-, and T1-weighted) images from 24 service members and veterans scanned in the Chronic Effects of Neurotrauma Consortium's (CENC) observational study were acquired. Manual annotations were employed for both training and evaluation using a leave-one-out strategy. Lesion load and overlap evaluative comparisons are complimented by feature rankings which showcase the utility of the concatenated approach. Our findings suggest supervised learning methods may be applied to quantify WMHs on routine brain imaging. Paired with correlative outcome data, supervised learning methods may allow for identification of imaging features predictive of diagnosis and prognosis in individual TBI patients.

# Introduction

## White matter hyperintensities in TBI

White matter hyperintensities (WMHs) are foci of abnormally increased signal intensity seen within white matter regions within the cerebrum and brainstem on fluid attenuation inversion recovery (FLAIR) magnetic resonance imaging (MRI) sequences. WMHs are a frequent finding following traumatic brain injury (TBI) and have been correlated with functional outcome and injury severity in both pediatric<sup>1,2</sup> and adult<sup>3–6</sup> populations. Further research involving WMHs has shown that regional distribution and volume of WMHs have been shown to possess prognostic value in the TBI patient<sup>2,6–8</sup>. Specifically, lesion volume in corpus callosum correlates with functional scores in the acute phase following injury, while lesion volume in frontal lobes correlates with scores at 1 year following injury<sup>6</sup>. Further, volume of FLAIR lesions within the corpus callosum, brainstem, and thalamus in patients with severe TBI correlates with Glasgow Outcome-Extended (GOS-E) scores—a numeric groupwise assessment used to classify “outcome” in TBI patients where “outcome” refers to the spectrum of possible prognoses from death to disability to recovery.<sup>4</sup> Additionally, in patients who are comatose following severe TBI the regional distribution of FLAIR lesions within the pons, midbrain, hypothalamus, basal forebrain, parietal, temporal, occipital lobes, and insula along with the observation of grasping or chewing behavior are associated with poor outcome<sup>7</sup>.

Despite the above findings which demonstrate that WMHs have potential prognostic value, they are not routinely employed as a diagnostic measure in clinical practice. Performing a comprehensive manual counting of number and distribution of lesions in the clinical setting is simply not practical. The development of such lesion quantification approaches may allow for the practical inclusion of this type of information within routine radiological practice. In this work, we present an automated framework for quantification of WMHs in multi-modal MRI using the random forest machine learning technique.

## Random forests for WMH segmentation

The random forests framework<sup>9</sup> is a popular machine learning technique that has demonstrated significant utility for supervised segmentation tasks (e.g., normal human brain segmentation<sup>10</sup>) and other computer vision applications (e.g.,<sup>11</sup>). Random forest-based paradigms have been successfully employed in the delineation of other neuropathologies<sup>12–17</sup> for both single and multi-modal acquisition protocols.

Random forests are conceptually straightforward<sup>9</sup>. The basic component of the random forest paradigm is the “decision tree” often represented by a flowchart or graph where internal nodes represent “tests”, or decisions, and the edges represent the outcome of those tests. The final, or end, nodes represent the various classifications produced by traversal through the decision tree. For the proposed application, individual voxels (and their corresponding feature values) are introduced at the root of a particular decision tree and traverse the edges and internal nodes ultimately ending up at one of the classification nodes according to the tests at each internal node. A single random forest model will consist of many such trees (often referred to as an “ensemble”). Although decision trees had been extensively studied, the success of employing collections of such weak learners for boosting machine learning performance (e.g., AdaBoost)<sup>18,19</sup> influenced the similarly styled conglomeration of decision trees into “forests” with randomized node optimization<sup>20,21</sup>. Finally, Breiman<sup>9</sup> improved accuracy by random sampling of training data (i.e., “bagging”) resulting in the current random forest technique applied here. As voxels and their feature values are “pushed” through each decision tree in the forest, votes for each label are accumulated and converted to probability values for all classification possibilities at each voxel location.

In this work, we develop a concatenated random forest framework with a feature image set (both spatial and intensity-based) for segmenting WMHs in a large TBI cohort. The entire framework is built on the well-known open-source Advanced Normalization Tools (ANTs)<sup>1</sup> and ANTsR<sup>2</sup> toolkits. Further motivating this research

---

<sup>1</sup><https://github.com/stnava/ANTs>

<sup>2</sup><https://github.com/stnava/ANTsR>

is the availability of several large publicly available imaging data sets that permits testing reproducibility of this automated routine for WMH segmentation and quantification.

## Materials and Methods

### Imaging

MR images utilized for this initial report were acquired from a single scanner involved in the Chronic Effects of Neurotrauma Consortium's (CENC) observational study (see Walker et al., this issue). Briefly, participants were Operation Iraqi Freedom/Operation Enduring Freedom (OIF/OEF) era Service Members and Veterans between the ages of 18-60 years with prior combat exposure and deployment(s). The feature images were derived from MR acquisitions of 26 subjects aged  $39.6 \pm 8.1$  years (range 28–58 years). Within this cohort, 24 (92%) were considered positive for TBI based upon the potential concussive events (PCE) interview process described in detail in Walker et al., this issue). All lesions were isolated in the white matter of the cerebrum. Table 1 provides a descriptive statistical summary of the variation in lesion load across the selected cohort.

Images were acquired on a Philips 3.0T Ingenia system with an 8-channel SENSE head coil (Philips Medical Systems, Best, Netherlands). 3D FLAIR sequences were acquired with a turbo spin echo inversion recovery sequence with the following parameters: repetition time (TR) = 4800 ms, echo time (TE) = 325 ms, inversion time (TI) = 1650 ms; 170 sagittal slices with a 1.2 mm slice thickness,  $256 \times 256$  acquisition matrix, and  $256 \times 256$  mm FOV. 3D T1-weighted sequences were acquired with a fast field echo (FFE) sequence with the following parameters: TR = 6.8 ms, TE = 3.2 ms, echo train length (ETL) = 240; Flip angle =  $9^\circ$ , 170 sagittal slices with a 1.2 mm slice thickness,  $256 \times 240$  acquisition matrix, and  $256 \times 256$  mm FOV. In addition, 3D T2-weighted images were acquired with a turbo spin echo sequence with the following parameters: TR = 2500 ms, TE = 245 ms, ET = 133; 170 sagittal slices with a 1.2 mm slice thickness,  $256 \times 256$  acquisition matrix, and  $256 \times 256$  mm FOV.

The first author (J. R. S.) performed the manual WMH tracings for all 26 subjects. J. R. S. is a radiologist

certified by the American Board of Radiology, with a certificate of advanced qualification in vascular and interventional radiology, over 18 years of research experience in TBI, and 6 years of clinical imaging experience. All multi-modal MR dicom image slices were converted to the nifti file format.<sup>3</sup> Given the utility of the FLAIR sequence in detecting white matter lesions in TBI<sup>3</sup>, all nifti image volumes for each subject were rigidly aligned to the FLAIR image of that subject using the ANTs software<sup>22</sup>. The normalized MRI volumes were then provided to J. R. S. who traced each lesion using the ITK-SNAP tool<sup>23</sup> which has multi-image overlay capabilities for visualizing all modalities in all three canonical views.

## **Quantitative analysis**

Figure 1 provides a graphical overview of the proposed workflow. The major components include offline generation of symmetric multimodal templates, the creation of feature images from the training data which are then employed for statistical prediction using a concatenated random forest framework. This framework involves the use of two random forest models where the outcome (i.e., the tissue probability estimates) of the first RF model application, which we denote as “Stage1”, is used as input (along with the original set of feature images) to a refinement RF model segmentation which we denote as “Stage 2”. Once these offline steps are performed, a new, unsegmented subject can then be processed using the proposed pipeline.

## **Symmetric multi-modal templates**

Following<sup>24</sup> and<sup>17</sup>, optimally derived templates serve for both brain tissue segmentation (for the derivation of tissue prior probability maps) and generation of asymmetry feature images. For this work we use the multi-modal data available from the public MMRR data set<sup>25</sup>. We used all 46 multi-modal acquisitions of that study to produce a multi-modal template according to the procedure described in<sup>26</sup> which results in a mean (in terms of both shape and intensity) multivariate template representing the entire cohort. Mid-canonical slices of the FLAIR, T1, and T2 components are illustrated in Figure 2.

---

<sup>3</sup><http://nifti.nimh.nih.gov/nifti-1>

## Feature images for WMH segmentation

Crucial to these supervised segmentation approaches are the creation and selection of “features” as input (i.e., feature images constructed from the training data) in conjunction with expertly identified structures of interest (i.e., WMHs) for model construction. For the targeted application in this work, tissue classification is performed at the voxelwise level. In other words, each voxel within the region of interest is sent through the ensemble of decision trees and receives a set of classification votes from each tree thus permitting a regression or classification solution. Since this procedure is performed at the voxelwise level, intensity information alone is insufficient for good segmentation performance due to the lack of spatial context. For example, as pointed out in<sup>27</sup>, higher intensities can be found at the periventricular caps in normal subjects which often confounds automated lesion detection algorithms. Other potential confounds include MR signal inhomogeneity and noise. Therefore, even though machine learning and pattern recognition techniques are extremely powerful and have significant potential, just as crucial to outcome is the creative construction and deployment of salient feature images which we detail below.

Supervised methodologies are uniquely characterized, in part, by the feature images that are used to identify the regions of interest. In Table 2, we provide a list and basic categorization of the feature images used for the initial (i.e., Stage 1—more on the use of multiple random forest stages below) segmentation of the WMHs. In addition Figure 3 provides a representation of a set of feature images for a single subject analyzed in this work.

Note that in this work we categorize the brain parenchyma with seven labels:

- cerebrospinal fluid (label 1),
- gray matter (label 2),
- white matter (label 3),
- deep gray matter (label 4),
- brain stem (label 5),

- cerebellum (label 6), and
- white matter hyperintensities (label 7).

As mentioned previously, input for each subject comprises FLAIR, T1-, and T2-weighted acquisitions. The T1 and T2 images are rigidly registered to the FLAIR image using the open-source Advanced Normalization Tools (ANTs)<sup>22</sup>. The aligned images are then preprocessed using the denoising algorithm of<sup>28</sup> followed by N4 bias correction<sup>29</sup> which are then normalized to the intensity range  $[0, 1]$ . Although we could have used an alternative intensity standardization algorithm (e.g.,<sup>30</sup>), we found that a simple linear rescaling produced better results similar to previous work<sup>17</sup>.

The T1 image is then processed via the ANTs brain extraction and normal tissue segmentation pipelines<sup>24</sup>. The result is a mask delineating the brain parenchyma and probabilistic estimates of the CSF, gray matter, white matter, deep gray matter, brain stem, and cerebellum<sup>31</sup>. These provide the expertly annotated labels for the first six tissue labels given above. Tissue prior probability maps for segmentation are from multi-model optimal symmetric shape/intensity templates<sup>17</sup> created from the public MMRR data set<sup>25</sup> (cf Figure 2).

Feature values include the preprocessed FLAIR, T1, and T2 image voxel intensities. We also calculate a set of neighborhood statistics (mean, standard deviation, and skewness) feature images using a Manhattan radius of one voxel given the typical size of individual WMHs. For each of the preprocessed images, we calculate the difference in intensities with the corresponding warped template component. Previous success in the international brain tumor segmentation competition<sup>32</sup> was based on an important set of intensity features that were created from multi-modal templates mentioned previously<sup>17</sup> and listed in Table 2. We employ the same strategy here.

To take advantage of the gross bilateral symmetry of the normal brain (in terms of both shape and intensity), and the fact that WMHs do not generally manifest symmetrically across hemispheres, we use the symmetric templates to compute the contralateral intensity differences as an additional intensity feature.

The segmentation probability images described above are used as feature images to provide a spatial context for the random forest model prediction step. Additional spatial contextual feature images include the distance maps<sup>33</sup> based on the csf, gray matter, and deep gray matter images. These latter images are intended to help distinguish white matter hyperintensities from false positives induced by the partial voluming at the gray/white matter interface. A third set of images are based on the voxel location within the space of the template. Similar feature images were used in<sup>34</sup> although, unlike the proposed framework, this previous work lacks normalization to the standard coordinate system provided by the template to dramatically improve spatial specificity across all subjects. To generate these images, the T1 image of each subject is registered to the T1 template component using a B-spline variant<sup>35</sup> of the well-known ANTs Symmetric Normalization (SyN) algorithm<sup>36</sup>. Using the derived transforms, the template coordinate images are warped back to the space of the individual subject.

### **Stacked (concatenated) random forests for improved segmentation performance**

In previous brain tumor segmentation work<sup>17</sup>, it was demonstrated that a concatenated supervised approach, whereby the prediction output from the first random forest model serves as partial input for a second random forest model, can significantly improve segmentation performance. We do the same thing for the work described here where we employ two stacked random forests (or two “stages”). The Stage 1 feature images of the training data (as described previously) are used to construct the Stage 1 model. The training data Stage 1 features are then used to produce the voxelwise “voting maps” (i.e., the classification count of each decision tree for each tissue label) via the Stage 1 random forest model. All the Stage 1 features plus the Stage 1 voting maps are used as input to the Stage 2 model. In addition, we use the Stage 1 voting maps as tissue priors (i.e., probabilistic estimates of the tissue spatial locations) for a second application of the 6-tissue segmentation algorithm with an additional Markov Random Field spatial prior (MAP-MRF)<sup>31</sup>. In order to maximize the spatial information for the  $n$ -tissue segmentation process following the voxelwise RF classification of Stage 1, we use all three aligned preprocessed images for multivariate segmentation during the second stage. The



resulting seven posterior probability images constitute a third additional feature image set for Stage 2.

## Implementation

As pointed out in a recent comprehensive lesion segmentation review<sup>37</sup>, although the number of algorithms reported in the literature is quite extensive, there were only four publicly available segmentation algorithms at the time of writing this article. In contrast to the current work, none are based on supervised learning. As we did for our brain tumor segmentation algorithm<sup>17</sup>, all of the code described in this work is publicly available through the open-source ANTs/ANTsR toolkits. Through ANTsR (an add-on toolkit which, in part, bridges ANTs and the R statistical project) we use the *randomForest* package<sup>38</sup> using the default settings with 2000 trees per model and 500 randomly selected samples per label per image. Note that we saw little variation in performance when these parameters were changed (i.e. up to 1000 random samples and as little as 1000 trees) which is consistent with our previous experience.

In addition, similar to our previous offering,<sup>4</sup> we plan on creating a self-encapsulated example to showcase the proposed methodology [which will also be available on github](#).<sup>5</sup> The fact that the data will also be made available through the Federal Interagency Traumatic Brain Injury Research (FITBIR) repository along with the manual labelings will facilitate reproducibility on the part of the reader as well as any interest in extending the proposed framework to other data sets.

## Evaluation protocol overview

In order to evaluate the protocol described, we performed a leave-one-out evaluation using the data acquired from the 24 subjects described above. Initial processing included the creation of all Stage 1 feature images for all subjects. The initial brain segmentation of each T1 image and the manual white matter hyperintensity tracings were combined to provide the truth labels for the training data. The “truth” labels are the seven anatomical regions given above.

---

<sup>4</sup><https://github.com/ntustison/ANTsAndArboles>

<sup>5</sup><https://github.com/ntustison/WatchMeHyperventilate>

The leave-one-out procedure is as follows:

- Create Stage 1 feature images for all 24 subjects.
- For each of the 24 subjects:
  - sequester the current subject and corresponding feature images.
  - construct the Stage 1 random forest model from the remaining 23 subjects.
  - apply the Stage 1 random forest model to the feature images of the 23 training subjects.
  - the previous step produces the Stage 1 voting maps for all seven labels.
  - for each of the 23 subjects, perform a Bayesian-based segmentation with an MRF spatial prior using the seven voting maps as additional tissue priors.
  - construct the Stage 2 random forest model from all the Stage 1 feature images, seven voting maps, and seven posterior probability maps from the previous step.
  - send the sequestered subject through the random forest models for both stages.
  - compare the final results with the manually-defined white matter hyperintensity regions.

Several measures have been employed in the literature for evaluating automated white matter lesion segmentation involving such quantities as lesion load, voxel-based overlap measures (such as the Dice similarity coefficient), and lesion-based measures.<sup>37</sup> For this work, due to the relatively small-size distribution<sup>6</sup> of the lesion load in our data cohort (see Table 1), we used four lesion-based measures: sensitivity, positive predictive value (i.e., precision),  $F_1$  score, and relative volume difference. The first three quantities are based on the number of false positives ( $TN$ ), false negatives ( $FN$ ), and true positives ( $TP$ ) in terms of identified lesions. It should be noted that the number of true negatives is not readily incorporated into measures of accuracy as the quantity of “true negatives” (i.e., correctly identified normal brain tissue) would severely skew accuracy assessments.

---

<sup>6</sup>For comparison, the training data set of the MS Lesion Segmentation challenge associated with the international MICCAI 2008 conference has a mean lesion load of  $204 (\pm 752) \text{ mm}^3$  per lesion and the resolution is almost twice what is used in this study (i.e.,  $0.5 \times 0.5 \times 0.5$ ).

The  $F_1$  score is an assessment of accuracy which takes into account both the sensitivity,

$$sensitivity = \frac{TP}{TP + FN},$$

and the positive predictive value (PPV),

$$PPV = \frac{TP}{TP + FP},$$

such that  $F_1$  is given by

$$F_1 = \frac{2 \cdot TP}{2 \cdot TP + FP + FN}.$$

The relative volume difference is calculated for each of the true positive lesions using the manual and predicted lesion volumes:

$$Relative\ volume\ difference = \frac{V_{manual} - V_{predicted}}{V_{manual}}.$$

To illustrate how the performance of our framework varies with lesion size, we calculated the above measures based on evenly split quantiles of the manual estimates of lesion volumes into 3 groups. These three size ranges (in terms of the number of voxels) were:  $[3 - 12]$ ,  $[12 - 28]$ , and  $[28 - 551]$ .

## Results

### White matter hyperintensity segmentation evaluation

In Figure 8 we provide the segmentation evaluations derived from the leave-one-out evaluation of the previously described TBI data. These performance measures include sensitivity, positive predictive value,  $F_1$  score, and relative volume difference. The three lesion size ranges over which these measures are computed to illustrate the variation in performance with lesion size. Smaller lesions ( $< 12$  voxels) are more difficult to identify

which is why the sensitivity is widely varied over the cohort vs. the largest set of voxels ( $> 28$  voxels). The first three measures are based on the identification of entire lesions. The relative volume difference provides a direct assessment of how accurate the volumetric estimate is when comparing the manually identified lesions versus the automatically predicted lesions.

### **Ranking feature importance**

After performing the leave-one-out evaluation, we calculated the *MeanDecreaseAccuracy* feature values for each of the  $24 \text{ subjects} \times 2 \text{ models per subject} = 48$  total models. This measure (per feature, per model) is calculated during the out-of-bag phase of the random forest model construction and quantifies the decrease in prediction accuracy from omitting the specified feature. In other words, this quantity helps determine the importance of a particular feature and, although we save such efforts for future work, this information provides us with guidance for future feature pruning and/or additions.

The resulting rankings for both Stages are given in Figures 5 and 6 where the values for the separate stages are averaged over the entire corresponding model set. In addition, we track the variance for each feature over all models to illustrate the stability of the chosen features during the evaluation. This latter information is illustrated as horizontal errors bars providing the 95<sup>th</sup> percentile. Note that the reader can cross reference Table 1 for identifying corresponding feature types and names.

Additionally, it is interesting to note some of the other top performing features for Stage 1. The contralateral difference FLAIR image is highly discriminative over the set of evaluation random forest models (see Figure 7). This accords with the known clinical relevance of FLAIR images for identifying white matter hyperintensities and the fact that such pathology does not typically manifest symmetrically in both hemispheres. Interestingly, the posterior maps for the deep gray matter are extremely important for accurate white matter hyperintensity segmentation. Perhaps the spatial specification of deep gray matter aids in the removal of false positives. Inspection of the bottom of the plots demonstrates the lack of discriminating features associated with the T1 image which is also well-known in the clinical literature.

As described earlier, for Stage 2, we used the output random forest voting maps from Stage 1 as both features themselves and as priors for input to a Bayesian-based segmentation with an additional MRF spatial prior. In Figure 6, the voting maps are labeled as “*RFStage1VotingMaps*” where the final numeral is associated with the brain parenchymal labeling given previously. Similarly, the additional RF prior segmentation feature probability maps are labeled as “*RFBrainSegmentationPosteriors*”. The Stage 2 feature importance plot follows similar trends as that for Stage 1 with the T1 images not contributing much to the identification of white matter hyperintensity voxels. The initial voting maps from Stage 1 are extremely important with the top 3 being the estimated locations of the 1) gray matter, 2) white matter, and 3) white matter hyperintensities. Since these tissue type can be conflated based on intensity alone it is intuitive that such features would be important.

## Discussion

The current communications describes a supervised statistical learning methodology for identifying WHMs within multimodal MR brain imaging. This effort utilized information acquired from the manual segmentation of WMHs from FLAIR images to help build two-stage ensembles of decision trees for the automated identification of these lesions. Although only a single expert was used to produce the manual labelings, our intent is to further refine the proposed paradigm by crowdsourcing with feedback from other experts who interact with both the data and methodology. Also, we recognize that only a single site was used for evaluating the proposed framework. However, we are currently processing other site data with the models developed for this work and the results look promising since the developed features are site-agnostic.

As far as we know, this is the first report utilizing a novel random forest approach to identify WMHs in a cohort of TBI patients. TBI WMHs tend to be more difficult to segment than MS lesions as the former tend to be smaller with an overall smaller lesion load. Also, enhancement protocols with the former tend to be less successful than with the latter. As mentioned previously, the work in MS lesion segmentation is extensive with a handful of techniques being publicly available.

Two major meta-analyses of WMHs have been published covering the periods prior to<sup>39</sup> and after 2010<sup>40</sup>. Debbete & Markus<sup>39</sup> found that the presence of WMHs was related to subsequent cognitive decline, a higher risk of developing dementia, stroke, and of mortality. Lesion volume at baseline was also predictive of cognitive decline. Kloppenborg et al.<sup>40</sup> of 23 cross-sectional studies reporting MRI and concurrent neuropsychological results in patients with heterogeneous diagnoses but without previously diagnosed cognitive impairment, found that WMHs were associated with cognitive deficit (effect size of -0.10, 95% CI: -0.13 to -0.08) after controlling for age.

Despite the potential clinical significance of WMHs these lesions receive little attention in current clinical workflows. When reported in a standard neuroradiologist interpretation, they are typically handled as incidental findings and are assigned little clinical significance. This likely reflects the impracticality of performing a detailed assessment of number, volume, and distribution within a qualitative neuroradiologist interpretation as well as the lack of correlative information on how the presence and distribution of these lesions may inform a diagnosis and prognosis in the appropriate clinical setting. To date, automated or semi-automated tools for the detection of WMHs have lacked the specificity and efficiency for the mining of large-scale datasets to generate highly granular data on whether these lesions possess any true diagnostic or prognostic value in the setting of a specific disease process. The present communication describes a supervised statistical learning tool that is appropriate for the application to such large-scale datasets.

## **Acknowledgements**

The authors wish to acknowledge all other members of the CENC Neuroimaging Steering Committee and CENC leadership (Drs. David X. Cifu, Ramon Diaz-Arrastia, and Rick Williams) for their support. We also gratefully acknowledge the assistance of Tracy Nolen, Chris Siegel and Kevin Wilson. We would also like to thank the study participants and their family members. This project was jointly supported by the Department of Defense (W81XWH-13-2-0095), the U.S. Department of Veterans Affairs (Io1 CX001135 and Io1 RX 002174), as well as USUHS Grant HU 0001-08-0001.

## **Declaration of Interest/Disclaimer**

The authors report no financial disclosures or conflicts of interest. The views expressed here are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government. This work was prepared as a part of official duties; Title 17 USC §105 provides that Copyright protection under this title is not available for any work of the U.S. Government. Title 17 USC §101 defines a US Government work as a work prepared by a military service member or employee of the US Government as part of that person's official duties.

## References

### Figure Captions

**Figure 1:** Workflow illustration for the proposed pipeline. Processing of the multi-modal input MRI for a single subject, using the multi-modal symmetric template, results in the generation of the feature images. These feature images are used as input to the Stage 1 RF model producing the initial RF probability map estimates. The Stage 1 voting maps, the original feature images, and the Stage 2 RF model result in the final voting maps which includes the WMH probability estimate. Note that the RF models are constructed once from a set of training data which are processed using the same feature-construction pipeline as the single-subject input MRI.

**Figure 2:** Canonical views of the multivariate, bilaterally symmetric template constructed from the MMRR data set<sup>25</sup> (only shown are the FLAIR, T1, and T2 modalities— the components relevant for this work). Template construction is detailed in<sup>17</sup>. These images are important for asymmetry-based features.

**Figure 3:** Representation of Stage 1 feature images for subject 01C1019. The FLAIR, T1-, and T2-weighted images are rigidly pre-aligned<sup>22</sup> to the space of the T1 image. The three modality images are then preprocessed (N4 bias correction<sup>29</sup> and adaptive denoising<sup>28</sup>) followed by application of standard ANTs brain extraction and  $n$ -tissue segmentation protocols using the MMRR symmetric template and corresponding priors<sup>24</sup> applied to the T1 image. The feature images are then generated for voxelwise input to the RF model which results in the voting maps illustrated on the right. This gives a probabilistic classification of tissue type. Not shown are the probability and voting images for the brain stem and cerebellum.

**Figure 4:** Sample FLAIR acquisition image slices showing both manual and random forest segmentations for both stages obtained during the leave-one-out evaluation. Manual segmentations were performed by one of the authors and provided the ground truth WMH labels for training the random forest models.

**Figure 5:** Average *MeanDecreaseAccuracy* plots generated from the creation of all 24 random forest models



for Stage 1 during the leave-one-out evaluation. These plots are useful in providing a quantitative assessment of the predictive importance of each feature. Features are ranked in descending order of importance. The horizontal error bars provide the 95<sup>th</sup> percentile and illustrate the stability of the feature importance across the leave-one-out models. At this initial stage only 31 feature images are used.

**Figure 6:** Average *MeanDecreaseAccuracy* plots generated from the creation of all 24 random forest models for Stage 2 during the leave-one-out evaluation. These plots are useful in providing a quantitative assessment of the predictive importance of each feature. Features are ranked in descending order of importance. The horizontal error bars provide the 95<sup>th</sup> percentile and illustrate the stability of the feature importance across the leave-one-out models. We augment the 31 feature images from the first stage by adding an additional seven voting maps and 7 segmentation posteriors from application of the Bayesian-based segmentation for a total of 45 images for the second stage.

**Figure 7:** (a) FLAIR image slice illustrating WMHs which have been manually delineated. The region around the WMHs is enlarged (b) in the original FLAIR and the (c) contralateral FLAIR difference image.

**Figure 8:** Voxelwise comparison with manual delineation of white matter hyperintensities. On the left are the calculated Dice values over all white matter hyperintensities. Note the improvement in the Dice metric from the employment of the Stage 2 component of the processing pipeline. (Right) Similar results can be seen by comparing the total lesion load volume between manual and automated detection strategies. Although some outliers are found after the Stage 2 processing in a couple subjects, the number of outliers caused by false positives is decreased significantly with the second stage processing.

1. Bigler ED, Abildskov TJ, Petrie J, Farrer TJ, Dennis M, Simic N, Taylor HG, Rubin KH, Vannatta K, Gerhardt CA, et al. Heterogeneity of brain lesions in pediatric traumatic brain injury. *Neuropsychology*. 2013;27(4):438–51.
2. Smitherman E, Hernandez A, Stavinocha PL, Huang R, Kernie SG, Diaz-Arrastia R, Miles DK. Predicting outcome after pediatric traumatic brain injury by early magnetic resonance imaging lesion location and volume. *J Neurotrauma*. 2016;33(1):35–48.
3. Marquez de la Plata C, Ardelean A, Koovakkattu D, Srinivasan P, Miller A, Phuong V, Harper C, Moore C, Whittemore A, Madden C, et al. Magnetic resonance imaging of diffuse axonal injury: Quantitative assessment of white matter lesion volume. *J Neurotrauma*. 2007;24(4):591–8.
4. Moen KG, Brezova V, Skandsen T, Håberg AK, Folvik M, Vik A. Traumatic axonal injury: The prognostic value of lesion load in corpus callosum, brain stem, and thalamus in different magnetic resonance imaging sequences. *J Neurotrauma*. 2014;31(17):1486–96.
5. Ding K, Marquez de la Plata C, Wang JY, Mumphrey M, Moore C, Harper C, Madden CJ, McColl R, Whittemore A, Devous MD, et al. Cerebral atrophy after traumatic white matter injury: Correlation with acute neuroimaging and outcome. *J Neurotrauma*. 2008;25(12):1433–40.
6. Pierallini A, Pantano P, Fantozzi LM, Bonamini M, Vichi R, Zylberman R, Pisarri F, Colonnese C, Bozzao L. Correlation between mRI findings and long-term outcome in patients with severe brain trauma. *Neuroradiology*. 2000;42(12):860–7.
7. Weiss N, Galanaud D, Carpentier A, Tezenas de Montcel S, Naccache L, Coriat P, Puybasset L. A combined clinical and mRI approach for outcome assessment of traumatic head injured comatose patients. *J Neurol*. 2008;255(2):217–23.
8. Levin HS, Williams D, Crofford MJ, High WM Jr, Eisenberg HM, Amparo EG, Guinto FC Jr, Kalisky Z, Handel SF, Goldman AM. Relationship of depth of brain lesions to consciousness and outcome after closed

head injury. *J Neurosurg.* 1988;69(6):861–6.

9. Breiman L. Random forests. In: *Machine learning.* 2001. pp. 5–32.

10. Yi Z, Criminisi A, Shotton J, Blake A. Discriminative, semantic segmentation of brain tissue in MR images. *Med Image Comput Comput Assist Interv.* 2009;12(Pt 2):558–65.

11. Viola P, Jones M, Snow D. Detecting pedestrians using patterns of motion and appearance. *International Journal of Computer Vision.* 2005;63:153–161.

12. Geremia E, Clatz O, Menze BH, Konukoglu E, Criminisi A, Ayache N. Spatial decision forests for MS lesion segmentation in multi-channel magnetic resonance images. *Neuroimage.* 2011;57(2):378–90.

13. Pustina D, Coslett HB, Turkeltaub PE, Tustison N, Schwartz MF, Avants B. Automated segmentation of chronic stroke lesions using lINDA: Lesion identification with neighborhood data analysis. *Hum Brain Mapp.* 2016 Jan.

14. Geremia E, Menze BH, Ayache N. Spatial decision forests for glioma segmentation in multi-channel MR images. In: *Proceedings of MICCAI-BRATS 2012.* 2012.

15. Bauer S, Fejes T, Slotboom J, Wiest R, Nolte L-P, Reyes M. Segmentation of brain tumor images based on integrated hierarchical classification and regularization. In: *Proceedings of MICCAI-BRATS 2012.* 2012. pp. 10–13.

16. Zikic D, Glocker B, Konukoglu E, Shotton J, Criminisi A, Ye DH, Demiralp C, Thomas OM, Das T, Jena R, et al. Context-sensitive classification forests for segmentation of brain tumor tissues. In: *Proceedings of MICCAI-BRATS 2012.* 2012. pp. 1–9.

17. Tustison NJ, Shrinidhi KL, Wintermark M, Durst CR, Kandel BM, Gee JC, Grossman MC, Avants BB. Optimal symmetric multimodal templates and concatenated random forests for supervised brain tumor seg-

mentation (simplified) with aNTsR. *Neuroinformatics*. 2015;13(2):209–25.

18. Schapire R. The strength of weak learnability. *Machine Learning*. 1990;5:197–227.

19. Freund Y, Schapire R. A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of Computer and System Sciences*. 1997;55:119–139.

20. Ho TK. Random decision forests. In: *Document analysis and recognition, 1995., proceedings of the third international conference on*. Vol. 1. 1995. pp. 278–282 vol.1.

21. Amit Y, Geman D. Shape quantization and recognition with randomized trees. *Neural Computation*. 1997;9:1545–1588.

22. Avants BB, Tustison NJ, Stauffer M, Song G, Wu B, Gee JC. The Insight ToolKit image registration framework. *Front Neuroinform*. 2014;8:44.

23. Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, Gerig G. User-guided 3D active contour segmentation of anatomical structures: Significantly improved efficiency and reliability. *Neuroimage*. 2006;31(3):1116–28.

24. Tustison NJ, Cook PA, Klein A, Song G, Das SR, Duda JT, Kandel BM, Strien N van, Stone JR, Gee JC, et al. Large-scale evaluation of aNTs and freeSurfer cortical thickness measurements. *Neuroimage*. 2014;99:166–79.

25. Landman BA, Huang AJ, Gifford A, Vikram DS, Lim IAL, Farrell JAD, Bogovic JA, Hua J, Chen M, Jarso S, et al. Multi-parametric neuroimaging reproducibility: A 3-T resource study. *Neuroimage*. 2011;54(4):2854–66.

26. Avants BB, Yushkevich P, Pluta J, Minkoff D, Korczykowski M, Detre J, Gee JC. The optimal template effect in hippocampus studies of diseased populations. *Neuroimage*. 2010;49(3):2457–66.

27. Neema M, Guss ZD, Stankiewicz JM, Arora A, Healy BC, Bakshi R. Normal findings on brain fluid-

attenuated inversion recovery mR images at 3T. *AJNR Am J Neuroradiol*. 2009;30(5):911–6.

28. Manjón JV, Coupé P, Martí-Bonmatí L, Collins DL, Robles M. Adaptive non-local means denoising of mR images with spatially varying noise levels. *J Magn Reson Imaging*. 2010;31(1):192–203.

29. Tustison NJ, Avants BB, Cook PA, Zheng Y, Egan A, Yushkevich PA, Gee JC. N4ITK: Improved N<sub>3</sub> bias correction. *IEEE Trans Med Imaging*. 2010;29(6):1310–20.

30. Nyúl LG, Udupa JK, Zhang X. New variants of a method of MRI scale standardization. *IEEE Trans Med Imaging*. 2000;19(2):143–50.

31. Avants BB, Tustison NJ, Wu J, Cook PA, Gee JC. An open source multivariate framework for  $n$ -tissue segmentation with evaluation on public data. *Neuroinformatics*. 2011;9(4):381–400.

32. Menze BH, Jakab A, Bauer S, Kalpathy-Cramer J, Farahani K, Kirby J, Burren Y, Porz N, Slotboom J, Wiest R, et al. The multimodal brain tumor image segmentation benchmark (bRATS). *IEEE Trans Med Imaging*. 2015;34(10):1993–2024.

33. Maurer CR, Rensheng Q, Raghavan V. A linear time algorithm for computing exact Euclidean distance transforms of binary images in arbitrary dimensions. *Pattern Analysis and Machine Intelligence, IEEE Transactions on*. 2003;25(2):265–270.

34. Anbeek P, Vincken KL, Osch MJP van, Bisschops RHC, Grond J van der. Probabilistic segmentation of white matter lesions in mR imaging. *Neuroimage*. 2004;21(3):1037–44.

35. Tustison NJ, Avants BB. Explicit B-spline regularization in diffeomorphic image registration. *Front Neuroinform*. 2013;7:39.

36. Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage*. 2011;54(3):2033–44.

37. García-Lorenzo D, Francis S, Narayanan S, Arnold DL, Collins DL. Review of automatic segmentation methods of multiple sclerosis white matter lesions on conventional magnetic resonance imaging. *Med Image*

Anal. 2013;17(1):1–18.

38. Liaw A, Wiener M. Classification and regression by randomForest. R News. 2002;2/3:18–22.

39. Debette S, Markus HS. The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: Systematic review and meta-analysis. BMJ. 2010;341:c3666.

40. Kloppenborg RP, Nederkoorn PJ, Geerlings MI, Berg E van den. Presence and progression of white matter hyperintensities and cognition: A meta-analysis. Neurology. 2014;82(23):2127–38.