

The High Time Resolution Universe Pulsar Survey – I. System configuration and initial discoveries

M. J. Keith,^{1*} A. Jameson,² W. van Straten,² M. Bailes,² S. Johnston,¹ M. Kramer,^{3,4}
A. Possenti,⁵ S. D. Bates,⁴ N. D. R. Bhat,² M. Burgay,⁵ S. Burke-Spolaor,^{1,2}
N. D’Amico,⁵ L. Levin,^{1,2} Peter L. McMahon,^{6,7} S. Milia^{5,8} and B. W. Stappers⁴

¹*Australia Telescope National Facility, CSIRO, PO Box 76, Epping, NSW 1710, Australia*

²*Swinburne University of Technology, Centre for Astrophysics and Supercomputing, Mail H39, PO Box 218, VIC 3122, Australia*

³*Max Planck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany*

⁴*University of Manchester, Jodrell Bank Centre for Astrophysics, Alan Turing Building, Manchester M13 9PL*

⁵*INAF – Osservatorio Astronomico di Cagliari, località Poggio dei Pini, strada 54, I-09012 Capoterra, Italy*

⁶*Berkeley Wireless Research Center, University of California, Berkeley, CA 94704, USA*

⁷*Department of Electrical Engineering, University of Cape Town, Rondebosch, 7701, South Africa*

⁸*Dipartimento di Fisica, Università degli Studi di Cagliari, Cittadella Universitaria, 09042 Monserrato (CA), Italy*

Accepted 2010 July 7. Received 2010 June 28; in original form 2010 May 11

ABSTRACT

We have embarked on a survey for pulsars and fast transients using the 13-beam multibeam receiver on the Parkes Radio Telescope. Installation of a digital backend allows us to record 400 MHz of bandwidth for each beam, split into 1024 channels and sampled every 64 μ s. Limits of the receiver package restrict us to a 340 MHz observing band centred at 1352 MHz. The factor of 8 improvement in frequency resolution over previous multibeam surveys allows us to probe deeper into the Galactic plane for short-duration signals such as the pulses from millisecond pulsars. We plan to survey the entire southern sky in 42 641 pointings, split into low, mid and high Galactic latitude regions, with integration times of 4200, 540 and 270 s, respectively. Simulations suggest that we will discover 400 pulsars, of which 75 will be millisecond pulsars. With ~ 30 per cent of the mid-latitude survey complete, we have redetected 223 previously known pulsars and discovered 27 pulsars, five of which are millisecond pulsars. The newly discovered millisecond pulsars tend to have larger dispersion measures than those discovered in previous surveys, as expected from the improved time and frequency resolution of our instrument.

Key words: pulsars: general.

1 INTRODUCTION

Pulsars are fascinating celestial objects, with spin periods ranging from ~ 1.6 ms to ~ 8.5 s and magnetic fields that span more than five orders of magnitude. Their study yields a better understanding of a variety of physics problems, from acceleration of particles in the ultrastrong magnetic field, to tests of gravity in the strong field regime. Pulsars provide useful probes of their environments whether inside pulsar wind nebulae, the centres of globular clusters or the Galactic Centre. Discovery of these unique objects has continued unabated for more than 40 yr, and the diversity in their properties means that, even though nearly 2000 pulsars, including ~ 170 millisecond pulsars (MSPs), are currently known, further discoveries are warranted.

In the past decade alone, surveys have uncovered a number of intriguing objects. These include the pulsar with the fastest rotation

period (in the globular cluster Terzan 5; Hessels et al. 2006), the double pulsar system (Burgay et al. 2003; Lyne et al. 2004) and its impressive tests of general relativity (Kramer et al. 2006), the ‘missing link’ between the low-mass X-ray binaries and the MSPs (Archibald et al. 2009), rotating radio transients (McLaughlin et al. 2006) and young, energetic pulsars such as PSR J1028–5819 (Keith et al. 2008) with their subsequent tie to gamma-ray pulsars and their emission properties (Weltevrede et al. 2010). Furthermore, timing of MSPs provides highly accurate parameters (Verbiest et al. 2009) possibly leading to the detection of gravitational waves in the near future (Hobbs et al. 2009). Finally, the discovery of a burst of radio emission of unknown origin but seemingly at cosmological distances (Lorimer et al. 2007) has motivated searches for other such events.

Since the discovery of pulsars (Hewish et al. 1968), searches have been split into targeted and untargeted surveys. Targets of interest include, for example, globular clusters, supernova remnants, the Galactic Centre and other regions known to be rich in pulsars. Although targeted searches are highly efficient in their use of

*E-mail: mkeith@pulsarastronomy.net

telescope time, it is impossible to predict the location of the majority of pulsars. Therefore, untargeted searches over large areas of sky are the only way to significantly increase the number of known pulsars.

Since the early 1990s, large-area pulsar surveys with the Parkes telescope, such as the low-frequency Parkes Southern Pulsar Survey (PSPS; Manchester et al. 1996) and the higher frequency Parkes Multibeam Pulsar Survey (PMPS; Manchester et al. 2001), Swinburne Intermediate Latitude Survey (SILS; Edwards et al. 2001) and its extensions (Jacoby et al. 2009) have led to the discovery of nearly 1000 pulsars. The all-sky PSPS was highly successful in finding MSPs (Lyne et al. 1998), but the low observing frequency limited the survey volume, especially towards the Galactic plane. The success of the later surveys was due to a combination of the higher observing frequency, the low system temperature and the large field of view provided by the 13-beam ‘multibeam’ receiver (Staveley-Smith et al. 1996). These surveys had relatively coarse frequency resolution however, and this limited the searchable volume for short-duration pulses from MSPs and/or transient bursts. For the PMPS this limit was due to the costs associated with replicating complex sets of analogue filters over 13 beams and the processing power available in the late 1990s.

In the 14 yr since the commissioning of the multibeam receiver, there has been a great advance in the development of backend recording technology. As pen chart recorders gave way to digital data analysis in the early 1970s, so analogue signal processing equipment is now being replaced by digital systems that offer a more affordable way to obtain higher time and frequency resolution. Increased frequency resolution allows for better correction of the dispersive effects of the interstellar medium, which, in combination with the higher time resolution, gives increased sensitivity to short-duration pulses, such as those from MSPs.

Encouraged by these developments, we conceived the ‘High Time Resolution Universe’ (HTRU) survey, motivated by two main drivers: to increase our understanding of the population of MSPs and to characterize the transient sky on time-scales down to tens of microseconds. These drivers led us to build a new digital backend system connected to the Parkes Multibeam Receiver. We use this new backend to give the HTRU survey four times the time resolution and eight times the frequency resolution of the PMPS. This allows us to detect MSPs for which interstellar dispersion had hidden from previous surveys (Bates et al., in preparation). With the discovery of the radio magnetar PSR J1622–4950 (Levin et al. 2010), we have also shown that the transient nature of some exotic pulsars gives us good reason to resurvey areas of sky that are already well studied.

The HTRU survey intends to be an all-sky survey for pulsars and short-duration radio transients, with a strong focus on the lower Galactic latitudes, where we make most use of the higher frequency resolution for negation of interstellar dispersion. A parallel effort is taking place at the Effelsberg Radio Telescope, covering the northern sky to a similar sensitivity, so as to provide a true all-sky survey at high time resolution.

In Section 2 of this paper we describe the survey strategy and the predicted results. Section 3 outlines the observing set-up in hardware and software. Section 4 describes the analysis software that is being used to process the survey data. In Section 5 we compute and measure the effective sensitivity limits of the survey. Finally, in Section 6 we give the basic parameters of the first pulsars discovered in the HTRU survey.

Table 1. Survey parameters for the three parts of the HTRU survey.

Survey	High	Mid	Low
Region	$\delta < +10^\circ$	$-120^\circ < l$ $l < 30^\circ$ $ b < 15^\circ$	$-80^\circ < l$ $l < 30^\circ$ $ b < 3.5^\circ$
τ_{obs} (s)	270	540	4300
N_{beams}	443 287	95 056	15 990
τ_{samp} (μs)	64	64	64
B (MHz)	340	340	340
$\Delta\nu_{\text{chan}}$ (kHz)	390.625	390.625	390.625
N_{chans}	870	870	870
Data length (samples)	$\sim 2^{22}$	$\sim 2^{23}$	$\sim 2^{26}$
Data/beam (GB)	1.0	2.0	16.0
Data/total (TB)	435	190	250

2 SIMULATION AND SURVEY STRATEGY

The HTRU survey of the southern sky is composed of three parts, outlined in Table 1. Other than the observation time and sky coverage, the observing parameters are identical for all three parts. The low-latitude component covers a thin strip of the Galactic plane, with Galactic latitude $|b| < 3.5^\circ$ and longitude $-80^\circ < l < 30^\circ$. The 4200-s observation time is twice that of the PMPS, providing the most thorough survey of the inner Galactic plane to date. The mid-latitude component covers $|b| < 15^\circ$ and $-120^\circ < l < 30^\circ$. Here the observation time is 540 s, allowing for a large area of sky to be covered quickly. Finally, the high-latitude component covers the entire sky south of declination $+10^\circ$ with 270-s integrations, excluding the region covered by the mid-latitude survey. Broadly speaking, the aim of the low-latitude part is to discover faint pulsars deep in the Galactic plane, whereas the mid-latitude survey will find bright MSPs suitable for timing array projects. Finally, the high-latitude survey will give us a snapshot of the transient sky at 64- μs resolution.

Detection of MSPs is limited by a combination of dispersion measure (DM) broadening in individual frequency channels, scattering in the interstellar medium and luminosity. For sufficiently luminous MSPs, the PMPS was predominantly limited by dispersion broadening in the 3-MHz channels, rather than by scattering. The increased frequency resolution of the HTRU survey means that the broadening is now dominated by scattering in essentially all directions. Fig. 1 shows contours of constant pulse broadening for the PMPS and HTRU surveys, taking into account both DM and scattering. Even in the worst case, towards the Galactic Centre, the limiting distance of the HTRU survey is greater than twice that of the PMPS, and significantly more in all other directions. This implies that the mid-latitude survey, with an integration time only one quarter that of the PMPS, should still discover pulsars that lie in previously surveyed regions. The deeper low-latitude survey will be able to penetrate further into the Galactic plane for both MSPs and longer period pulsars.

In order to determine the discovery potential of the HTRU survey, we ran simulations of the pulsar population based on the model presented in Lorimer et al. (2006) using the PSRPOP¹ software.

For the normal pulsar population, the input parameters to the model were as follows.

- (i) A log-normal spin period distribution, where the log of period (in ms) has mean 2.71 and standard deviation 0.34.

¹<http://psrpop.sourceforge.net>




Figure 1. Contours of constant pulse broadening time-scale for lines of sight at $b = 0$ for a frequency of 1352 MHz. The values for the PMPS are shown in dashed lines and for the HTRU survey in solid lines, with contours at 0.5, 1, 5 and 10 ms. Scattering time-scales and DMs were computed using the model of Cordes & Lazio (2002). The Earth and the approximate location of the Galactic Centre (GC) are shown with crosses.

- (ii) A simple power law for the luminosity distribution, with index -0.59 and low-luminosity cut-off at 0.1 mJy kpc^2 .
- (iii) An exponential function for the height above the Galactic plane, with scale 0.33 kpc .
- (iv) The radial distribution of Yusifov & Küçük (2004).
- (v) A total of 28 000 pulsars, such that the simulated detection of the PMPS matched closely to the true discovery rate.

The number of known Galactic MSPs is small and many previous surveys have been restricted by interstellar dispersion, making simulation of the Galactic population of MSPs more uncertain, and indeed this population was not considered by Lorimer et al. (2006). To produce simulations consistent with observed detection rates, we made the following modifications to the model.

- (i) The period distribution was based on that of the known Galactic plane MSPs.
- (ii) A larger scaleheight of 0.5 kpc was used to better match the known MSP population (Cordes & Chernoff 1997).

Along with the three parts of the HTRU survey, we also simulated the PSPS, PMPS and SILS surveys. The results are summarized in Table 2. In total, the HTRU surveys should discover nearly 400 pulsars, of which 75 will be MSPs, which would represent a doubling of the known population of MSPs. Final results from the survey will also allow us to obtain a better description of the radial and velocity distributions of MSPs.

3 OBSERVING SYSTEM

Observational data are acquired using the Parkes 21-cm Multibeam Receiver (Staveley-Smith et al. 1996) and a new digital backend system, the details of which are described in the following sections.

3.1 Analogue systems

The Parkes Multibeam Receiver consists of 13 feeds at the prime focus of the Parkes 64-m antenna, organized as a central feed sur-

Table 2. Simulated results for the mean number of total pulsar detections and new discoveries for three previous surveys and the three components of the HTRU survey. The numbers are split the simulation of ‘normal’ pulsars and ‘MSPs’ as described in the text. The numbers in parentheses are the actual numbers of pulsars found in each of the complete surveys, where we defined a pulsar as an MSP if it had a period of less than 30 ms and a period derivative less than 10^{-17} .

Survey	Normal		MSPs	
	Total	New	Total	New
PSPS	281 (279)	– (89)	20 (19)	– (17)
PMPS	1008 (1047)	876 (721)	19 (20)	18 (21)
SILS	137 (158)	57 (61)	10 (11)	6 (8)
HTRU				
Low-lat	957	260	51	33
Mid-lat	831	52	48	28
High-lat	783	11	65	13




Figure 2. The beam pattern of the multibeam receiver. The black circles indicate the 14-arcmin FWHM of each of the 13 beams of the receiver. The separation D_1 is also equal to this 14-arcmin FWHM and $D_2 = \sqrt{3}D_1$. The filled circles show three further pointings interleaved to make up the shape from which the entire survey is tessellated.

Table 3. Details of the multibeam receiver for the central feed, the inner ring and outer ring of feeds, taken from Manchester et al. (2001).

Beam	Centre	Inner	Outer
Telescope gain (K Jy^{-1})	0.735	0.690	0.581
Half-power beamwidth (arcmin)	14.0	14.1	14.5
Beam ellipticity	0.0	0.03	0.06
Coma lobe (dB)	none	17	14

rounded by two hexagonal rings, with beam pattern on the sky as shown in Fig. 2. Our measurement of the system temperature of the central beam – 23 K at 1400 MHz – is slightly higher than the value quoted in Manchester et al. (2001), which may be due to the refurbishment of the analogue electronics in 2006 (Reynolds, private communication). The central feed has a symmetric beam pattern, and the feeds further from the prime focus have a slight ellipticity and gain degradation. The details of the receiver and telescope set-up are listed in Table 3. For each beam of the receiver,

both polarizations are down-converted from 1182–1582 MHz to 0–400 MHz.

Unfortunately, the high-frequency part of this band is badly affected by interference from the *Thuraya* 3 geostationary communications satellite, which transmits a space-to-earth signal in the 1525–1559 MHz band. We have installed low-pass hardware filters after the receiver that restrict the upper end of our observing band to 1522 MHz; therefore, the effective observing band is 340-MHz wide, centred on 1352 MHz.

3.2 The BPSR backend system

The Berkeley–Parkes–Swinburne Recorder (BPSR) system consists of 26 digital spectrometers connected to 13 server-class workstations that format the data and write it to disk. The digital spectrometers are based on the IBOB² platform developed by the CASPER³ group at the University of California, Berkeley, described in detail in McMahon (2008).

Each IBOB consists of two analogue-to-digital converters and two polyphase filters implemented on an FPGA.⁴ Each pair samples orthogonal polarizations at 800 MHz using 8 bits per sample, then divides each signal into 1024 spectral channels of width 390 kHz. Each spectral channel is detected and integrated over 25 samples to yield the output time resolution of 64 μ s. The 32-bit output of the spectrometer is linearly scaled before decimation to 8 bits; the 8-bit data are written to a single UDP⁵ packet with a 2048-byte payload containing 1024 channels times two polarizations. Each packet is sent to a 10 Gb s⁻¹ Ethernet (10 GbE) interface connected directly to a Dell PowerEdge 1950 server via a CX4 cable.

On the server, data are collected into 10-s blocks and further processed. Both polarization streams of each channel are summed and then normalized by subtracting the mean and dividing by the standard deviation (as measured in the first 10-s block of each observation). The resultant data are then decimated to 2 bits per sample and written to disk at a rate of 32 Mb s⁻¹.

3.3 Data collection

Each server in the BPSR cluster has sufficient disk space to buffer ~ 3 d of observations. For the HTRU survey, we have chosen to archive the data to magnetic tapes before removing the data from the buffer, allowing for continuous observing. One copy is written to tape at Parkes and another is streamed via a dedicated 1 Gb s⁻¹ fibre link to the supercomputer located at Swinburne University. Here the data are also written to magnetic tape and, if there is sufficient free capacity on the supercomputer, run through the survey data processing pipeline in close to real time.

In addition to the astronomical signal, the BPSR backend also records a header file and a collection of auxiliary data files for each beam. The header file contains information about the observation in an XML⁶ format as described in Section 3.4. The auxiliary files contain bandpass and low-resolution time series snapshots, as described in Section 3.5. The raw time series is stored in the SIGPROC⁷

‘filterbank’ data format, which consists of a binary header followed by continuous raw data.

For the HTRU survey, all data are recorded using 2 bits of precision and with four samples packed per byte, with the first sample in the lowest significant bits of the byte. The data are ordered first by frequency and then by time, each time the sample being stored as a consecutive 256-byte word. The raw data are preceded by a 345-byte header, used for backwards compatibility with the SIGPROC software; however, more information is available in the XML header file. These data are then packed with the header and auxiliary files into a single GNU tarball per beam.

3.4 Psrxml header format

The psrxml header is a format-independent XML header file that describes both the observational properties of the recorded data and the format of the actual data itself. The psrxml header file is stored separately to the raw data, allowing trivial indexing of header data for very large data files, such as those produced by the BPSR backend.

The psrxml data format is intended to be suitable for long-term archival of pulsar data for future uses. Features of note are as follows.

(i) Checksumming of data blocks: as the data are written to disk, each block of data (usually a few hundred megabytes) is passed through the SHA-1⁸ hash algorithm and the hash value stored in the header file. This allows for the data to be checked for errors and consistency at a later time. Archival data are valuable: this checksumming ensures that data extracted from an archive are identical with the data recorded at the telescope.

(ii) Format and machine independence: the psrxml header file is written in a well-defined format, typically encoded in standard 7-bit ASCII.⁹ This means that the header will be readable on almost any computer that exists or is likely to exist. Since the header completely describes the data format, in terms of bit and byte ordering, and which bytes to read, one can therefore read the raw data on any computer with the appropriate software.

(iii) Human readable: XML documents can be easily read using a standard text editor. With the use of XSL,¹⁰ viewing a psrxml header file in a web browser displays the data in a pleasant format without the use of any specialist software. This does not change the file in any way, so the file is readable in ideal formats by both machine and human.

(iv) Extendable: since the header is written in XML, it is trivial to add new XML tags to the file in a safe manner, storing either fields specific to a certain project, or extending the data format to new generic uses.

3.5 Auxiliary files

In addition to the header and data files, data generated for online monitoring of the system are stored for future use. This consists of the passband and time series measured in the pre-normalized, dual-polarization, 8-bit data samples, completely averaged in frequency.

² Internet Break-Out Board.

³ Center for Astronomy Signal Processing and Electronics Research.

⁴ Field Programmable Gate Array.

⁵ User Datagram Protocol.

⁶ eXtensible Markup Language, <http://w3.org/TR/REC-xml/>.

⁷ <http://sigproc.sourceforge.net>

⁸ Secure Hash Standard, Federal Information Processing Standards Publications, 180-1, 1995.

⁹ American Standard Code for Information Interchange, American National Standards Institute, X3.4-1986, 1986.

¹⁰ eXtensible Stylesheet Language.

3.6 Data tracking

At the completion of every observation, a copy of the psrxml header file is kept on disk and indexed in a book keeping data base. The data base links the observation against the planned observations and marks that observation as completed, removing that pointing from the pool of pointings from which observing schedules are selected.

4 PROCESSING

Processing of the survey is currently being carried out at the Jodrell Bank Centre for Astrophysics and the Swinburne Centre for Astrophysics and Supercomputing.

The analysis of our data is limited by the available computational power, therefore reprocessing of the data with future computer hardware will likely yield additional results. We follow a typical pulsar search workflow as described, for example, in Lorimer & Kramer (2005).

4.1 Software pipeline

For our survey processing we have developed the HTRUN processing pipeline. HTRUN is designed to be a flexible system that can process both BPSR data and legacy 1-bit data formats without modifying the software. This flexibility, as well as being a useful tool, allows us to easily test the software on legacy data and against legacy codes.

The HTRUN pipeline is composed of several stages: radio frequency interference (RFI) removal, dedispersion, time series analysis, candidate sorting, candidate folding and optimization. Here, we outline the algorithms used for each step of the process.

4.1.1 RFI removal

RFI removal consists of two tasks, removal of ‘bad’ spectral channels and removal of ‘bad’ time samples.

First, we remove spectral channels that are affected by periodic RFI. Initially, we have a list of channels that are known to have poor data quality, and these are always removed. Then, we Fourier analyse each of the channels individually, using the Fourier time series analysis described below. If any signals are detected with a signal-to-noise ratio (S/N) of greater than 15, we reject that channel.

Next, we sum the data in frequency, producing a single time series at $DM = 0$. This time series is then searched for bursts of emission, and if found, those samples are removed from the time series. To remove longer duration RFI bursts, we sum together 2, 4, 8, 16 and 32 adjacent samples and repeat the search. Any time samples that are removed in this process are then removed from the original data file, replaced by samples that are drawn from a random population statistically indistinguishable from the data. An example of this RFI removal is shown in Fig. 3, where a series of short-duration wide-band signals have been removed from the data. It should be noted that this process will remove low-DM astrophysical signals of sufficient intensity and width. With our 5σ threshold for removal, astrophysical signals of maximum recordable intensity will be removed if their DM is less than $0.12(W/64 \mu s) \text{ cm}^{-3} \text{ pc}$, where W is the approximate width of the pulse.

4.1.2 Dedispersion

Since the DM of any signal is not known a priori, we have to search many trial DM values. We must therefore transform the single




Figure 3. The intensity of recorded signals over a short time as a function of frequency, before (left-hand panel) and after (right-hand panel) RFI removal. The RFI can be seen as a series of vertical stripes, which are replaced with random noise by the RFI excision routine.

time series with many frequency channels into many frequency-summed time series with different dispersion delays. For our survey, the processing time is largely dominated by the dedispersion time, making it important to choose the most efficient algorithm possible.

The most efficient dedispersion algorithm known is the ‘tree’ dedispersion approximation of Taylor (1974). This, however, assumes a linear dependence of delay on frequency, which is a good approximation for small fractional bandwidths. For many recent surveys with large fractional bandwidths this approximation is inappropriate, and work-arounds have been used. In the PSPS, a two-step approach was used where smaller subbands were dedispersed using the tree algorithm, before a final dedispersion step to combine the subbands (Manchester et al. 1996). In addition to this, the PMPS used a linearization approach where the data were padded with empty channels in a manner such that the dispersion sweep could be considered linear (Manchester et al. 2001). For the HTRU survey we have decided not to use the tree algorithm as the cost of linearizing the data outweighed the gain from the more efficient algorithm. Instead, we have chosen to implement an efficient ν^{-2} dedispersion algorithm.

Much of the cost of dedispersion is due to the large volume of data that must pass through the system; therefore we compact the data as much as possible. Since we sum 1024 2-bit values per output sample, there are 4096 possible output levels. Our processing systems perform 64-bit integer arithmetic, so we can pack four samples per 64-bit word, giving a factor of 4 processing throughput increase. By dividing the output by 16, the number of levels is reduced to 256, with little sensitivity loss. We therefore only need to write out 8 bits per sample, keeping the input/output costs to a minimum. Additionally, we make use of modern multicore processors with multithreaded code performing four dedispersion threads simultaneously. In future, it will likely be possible to gain additional performance benefits with technologies such as graphics processing unit coprocessing.

4.1.3 Time series analysis

By this stage of the processing we have a number of dedispersed time series on disk, which have to be searched for pulsar signals. HTRUN provides both time-domain and frequency-domain methods.

In the frequency domain we take the power spectrum of the data using a fast Fourier transform. We then remove the strong red-noise component of the spectrum at frequencies below about 10 Hz by subtraction of a running mean and division by a running variance. This also has the effect of normalizing the spectrum. The resulting power spectrum is then searched for signals with a S/N of greater than 6. To reclaim power in the harmonics of narrow pulses, we also search spectra which have had two, four, eight and 16 harmonics

summed. Potential signals from each DM trial are written out for later sorting, as described below.

In the time domain we conduct a search with similar methodology to that reported in Burke-Spolaor & Bailes (2010). In brief, we search for bright, single pulses by collating all points above 6σ from the mean, after which the detections from a single pointing are compared across all the beams to identify sky-localized events. Details of the single-pulse-specific analysis and the results of the time-domain search will be reported in a future paper.

4.1.4 Candidate sorting

Candidate sorting is the process of collecting the Fourier detections from each DM step and combining those into a manageable number of candidate pulsars for further analysis. This is done by grouping detections with the same frequency into a single candidate, and by combining candidates that are harmonically related. In *HITRAN*, this is done using software from *PULSARHUNTER*.¹¹

4.1.5 Candidate folding and optimization

We then fold the data, producing an averaged pulse profile for each candidate. Folding is a coherent technique unlike the incoherent harmonic summing in the Fourier domain. *HITRAN* utilizes the *DSPSR*¹² software to fold each candidate, producing profiles for each of the 16 frequency bands and 32 subintegrations.

Then the *PSRCHIVE* package (Hotan, van Straten & Manchester 2004) is used to sum the folded subintegrations and periods with a series of trial periods and DMs around the values returned from the Fourier domain search. The optimized period and DM, i.e. that which give highest folded S/N, are then used to make the final diagnostic plots for the candidate.

4.2 Selection of candidates

The goal of the data processing is to select a number of likely pulsar signals for reobservation. To achieve this, we take the optimized candidates and the associated diagnostic plots and view them with graphical candidate selection tools, such as *JReaper* (Keith et al. 2009). These tools reduce the large number of candidates to a series of points on a scatter chart, usually with period and S/N on the axes. Selecting these points presents the user with the full candidate details and diagnostic plots, at which point the user must exercise judgement on whether the candidate should be flagged for further follow-up.

Since manual candidate selection is a time-consuming and potentially error-prone process, we are investigating the potential for using artificial neural network (ANN) algorithms to pre-select promising candidates for viewing (Eatough et al. 2010). An ANN is a mathematical construct, consisting of a set of nodes that are connected to form a decision tree that, based on a given input vector, derives a conclusion about the validity of a hypothesis. In our case, we provide the ANN with a vector of ‘scores’, formed from analysis of the diagnostic plots, e.g. the agreement of the observed S/N–DM curve of a candidate with theoretical predictions as measured by a χ^2 estimator. The more types of scores that are used, the larger (usually) the chance to differentiate between pulsar and non-pulsar signals; however, the more known pulsar signals are required to

‘train’ the ANN (Eatough et al. 2010). Indeed, our ANN is poor at selecting binary or millisecond pulsars, because of the limited size of training set available for these classes of pulsar. The effectiveness of this technique can be seen in the discovery of PSR J1701–44, which was identified as a candidate by the ANN. The S/N of the candidate was rather low, at only 7, and would quite probably have been missed by a graphical interface reliant upon a human to select the candidate manually.

Although the ANN is limited in the types of pulsars it will select, it does provide a thorough analysis of all the candidates free from human error. We find that adding the ANN to our existing candidate selection effort gives a beneficial additional way to select candidates, with little cost in terms of the required computing power. Direct inspection of candidates is vital, however, and remains the primary way that pulsars have been discovered in the *HTRU* survey.

5 SURVEY SENSITIVITY

Knowledge of the minimum detectable flux density of the survey is vital for determining the survey effectiveness. Determining this limiting flux density is not trivial, since the survey is not uniformly sensitive, even over a small region of sky, nor to all pulsars. One option is to compute a theoretical mean sensitivity to some standard ‘typical’ pulsar, as done by other large pulsar surveys. This gives a value that can be easily compared to other similar surveys.

To measure the actual sensitivity of the survey, we can look at redetections of known pulsars. Since published values exist for the position and flux density of thousands of pulsars, we can cross-match those positions with our survey pointings. We then compute the theoretical S/N value for that pulsar given the offset in the beam position and compare that to the output of the search pipeline. We consider each of these in turn below.

5.1 Theoretical sensitivity

The radiometer equation can be used to compute the fundamental limiting flux density for the centre of the central beam of a given survey observation.

$$S_{\min} = \frac{\sigma(T_{\text{sys}} + T_{\text{sky}})}{G\sqrt{2B\tau_{\text{obs}}}}, \quad (1)$$

where σ is the minimum acceptable S/N, T_{sys} is the system noise temperature, T_{sky} is the sky noise temperature, G is the system gain, B is the observing bandwidth and τ_{obs} is the integration time. T_{sky} is only a significant contribution to the system noise for low Galactic latitudes and has a strong dependence on the area of sky that is observed. The value of T_{sky} at 1352 MHz for the survey region is extrapolated from the Haslam et al. (1981) sky map at 408 MHz. For equation (1) we use the mean values listed in Table 4; however, it should be noted that T_{sky} , and therefore S_{\min} , varies considerably across the sky. Here we use $\sigma = 8$, $T_{\text{sys}} = 23$ K and $G = 0.735$ to compute a S_{\min} of 0.20, 0.47 and 0.61 mJy for the low-,

Table 4. The maximum, minimum and mean sky noise temperature over the three survey regions. Note that the values for the mid-latitude survey exclude the low-latitude survey region.

Survey	Mean (K)	Max (K)	Min (K)
Low	7.6	36.0	1.6
Mid	2.5	9.1	0.6
High	1.0	2.4	0.6

¹¹ <http://www.pulsarastronomy.net/wiki/Software/PulsarHunter>

¹² <http://dsp.sr.sourceforge.net>




Figure 4. The mean limiting flux density for the mid-latitude survey shown as a function of pulsar period for various values of DM. Values shown here assume a pulse duty cycle of 5 per cent.

mid- and high-latitude portions of the survey, respectively. However as described below, the practical limiting flux density is lower due to the narrow pulse width of most pulsars.

Our Fourier-based detection algorithm is not uniformly sensitive to all DMs, pulse periods and pulse shapes. This is somewhat difficult to quantify; however, we follow the approach of Manchester et al. (2001) by modelling the frequency response of the pulsar as a uniform train of impulses separated by frequency $f_p = 1/P$. This train of pulses has amplitude $1/S_{\min}$, to which we multiply a series of functions representing the various transformations that the data undertake.

We can model each pulse as a Gaussian with width $W_{50} = 0.05P$, so that the frequency response is multiplied by the Gaussian

$$F(f) = \exp\left(-\frac{(\pi f W_{50})^2}{4 \ln(2)}\right). \quad (2)$$

The pulses are also convolved with the effect of interstellar dispersion, which can be modelled as another Gaussian with width τ_{DM} , the dispersion delay. Finally we must account for the finite sampling interval, τ_s , which multiplies the amplitudes by

$$F(f) = \left| \frac{\sin(\pi f \tau_s)}{\pi f \tau_s} \right|. \quad (3)$$

The final S/N is computed after summing N harmonics, where $N \in \{1, 2, 4, 8, 16\}$. Fig. 4 shows the minimum detectable flux density as a function of pulsar period and DM for the mid-latitude survey. It can be seen that the HTRU survey is sensitive to MSPs out to DMs of a few hundred $\text{cm}^{-3} \text{pc}$. The effects of T_{sky} and the differing observing time for the three surveys can be seen in Fig. 5, which shows the sensitivity curves for the three surveys at the minimum and maximum T_{sky} in that survey region.

The mean sensitivity of the survey is reduced by a number of factors. First, the outer beams of the receiver are less sensitive than the centre beam, and the sensitivity of each beam reduces towards the edges. This means that the sky coverage is not uniform, with the mean sensitivity over the mid-latitude survey of about 0.25 mJy.

5.2 Redetection of known pulsars

A total of 290 previously known pulsars lie within one beamwidth of one of the HTRU survey pointings collected to date. For each pulsar we obtained the flux density and its error from the pulsar




Figure 5. Comparison of the sensitivity of the three survey regions for a pulsar with a DM of $100 \text{ cm}^{-3} \text{pc}$. Each region has two curves, showing the values for the minimum and maximum value of T_{sky} in the survey region.

catalogue,¹³ a value of T_{sky} at the pulsar position, and the offset between its position and the centre of the relevant beam. We then computed the expected S/N for each pulsar following the process outlined in Section 5.1.

In all, we detected 223 of the 290 pulsars. For the 67 pulsars we failed to detect, five observations were ruined by broad-band RFI, 54 pulsars were not detected above a S/N of 10 even after we folded observations at the known pulsar ephemeris. This leaves eight pulsars that are seen in folded observations with a S/N of greater than 10 but were not detected in our search procedure. Each of the eight pulsars had a spin period above 500 ms, and so the Fourier S/N may well have been reduced by a larger red-noise component in the data, or the pulsar period falling between Fourier bins. The loss of pulsar detections due to RFI is an uncontrollable aspect of search observations. Although we have applied the techniques described in Section 4.1.1, they are unable to remove this type of broad-band RFI. Future reprocessing using the technique of Eatough, Keane & Lyne (2009) may be possible; however, this is currently unfeasible as it would increase the time for processing the survey data by more than a factor of 10.

Fig. 6 shows the expected versus actual S/N for the 223 detected pulsars. Although the correlation between the predicted and observed values is good, the predicted values are, on average, somewhat higher than the observed values. This can be caused by either sensitivity loss in our observing system, or error in the catalogued flux densities. First, we checked that the backend was not introducing any loss by collecting data simultaneously with BPSR and the Parkes Digital Filterbank 3 (PDFB3). These observations show that the BPSR and PDFB3 performance are equivalent and we therefore consider it unlikely that the backend is introducing any losses. Hobbs (private communication) has shown that the catalogue flux densities appear to be systematically higher than those obtained in interferometric observations. This bias may be due to the tendency to publish only the data with the highest S/N, for example when scintillation boosts the apparent flux density. Given our ability to detect the majority of the pulsars that are in our search area, we are relatively confident that the system is performing to specification.

¹³ <http://www.atnf.csiro.au/research/pulsar/psrcat/>




Figure 6. The predicted S/N for 223 known pulsars redetected in the HTRU survey, shown as a function of the observed S/N. Errors shown are from the published error in the pulsar flux density. The line shows the expected 1:1 correlation.

6 NEW PULSAR DISCOVERIES

In this paper we present the results of the analysis of the first ~ 30 per cent of the mid-latitude part of the survey. Results so far include:

- (i) Discovery of five MSPs, all of which are in binary systems. Full details of these pulsars will be reported in Bates et al. (in preparation).
- (ii) Discovery of the radio pulsar with the highest known magnetic field which appears to be a radio-loud magnetar, as reported in Levin et al. (2010).
- (iii) Discovery of 21 pulsars with periods between 0.2 and 3.5 s and moderate to high DMs. These are likely ‘standard’ solitary radio pulsars but full timing solutions still need to be obtained.

The Galactic coordinates, spin period and DM for each of these pulsars are given in Table 5.

6.1 Comparison to PMPS

Fig. 7 shows a comparison between the sensitivities of the mid- and low-latitude HTRU survey and the PMPS for a DM of $100 \text{ cm}^{-3} \text{ pc}$. Although the mid-latitude HTRU survey has only one quarter the observing time of the PMPS, it is significantly more sensitive to short-period pulsars at large DMs. This is due to a combination of the faster sampling rate, narrower frequency channels, slightly increased total bandwidth and multibit sampling of our survey.

Of the new discoveries, nine pulsars lie within the sky coverage of the PMPS. We have reanalysed archival data for each of these pulsars and the results are presented in Table 6. Four of these pulsars are MSPs with high DMs and are included in Fig. 7. It can be seen that all bar one fall below the (nominal) PMPS detection threshold. In fact, re-examination of the PMPS archival data shows that both PSR J1708–35 and J1801–32 are visible but that the pulse broadening from the 3 MHz channel bandwidth makes them difficult to distinguish from sinusoidal interference in a blind search.

Table 5. Basic parameters for the 27 pulsars discovered in the HTRU survey to date. Pulsars for which no full timing solution is published have been assigned a temporary name containing only two digits of declination.

PSR J	l ($^{\circ}$)	b ($^{\circ}$)	Period (s)	DM ($\text{cm}^{-3} \text{ pc}$)
1125–58 ^a	291.8	2.6	0.003 102	125
1330–52	308.8	9.6	0.648 104	149
1433–50	318.9	9.1	1.017 495	98
1442–51	320.1	7.7	0.732 061	97
1517–46	327.38	9.22	0.886 61	139
1612–58	327.0	−5.0	0.615 520	172
1623–4950 ^b	333.9	−0.1	4.325 800	881
1625–49	334.6	0.0	0.355 856	717
1627–59	327.3	−7.3	0.354 239	92
1648–36	347.1	5.8	0.212 316	222
1650–60	328.2	−10.2	0.583 771	106
1701–44	342.3	−1.3	0.755 535	413
1705–43	343.3	−1.5	0.222 561	184
1705–61	328.7	−12.2	0.808 546	87
1708–35 ^a	350.4	3.2	0.004 505	146
1731–18 ^a	6.9	8.2	0.002 345	106
1753–38	352.3	−6.4	0.666 804	168
1754–24	4.9	0.8	2.090 263	789
1801–32 ^a	358.9	−4.5	0.007 454	177
1802–33	357.6	−5.5	2.461 052	254
1803–33	358.0	−5.6	0.633 412	171
1810–01	27.0	8.6	0.744 976	135
1811–24 ^a	7.1	−2.5	0.002 661	60
1811–49	344.2	−14.2	1.432 704	49
1812–27	4.0	−4.6	0.236 983	105
1814–05	23.8	5.8	1.014 405	119
1854–15	19.0	−8.0	3.453 121	157

^aTiming parameters in Bates et al. (in preparation).

^bTiming parameters in Levin et al. (2010).




Figure 7. Comparison of the mid- and low-latitude parts of the HTRU survey and the PMPS. The lines give the sensitivity limit for a DM of $100 \text{ cm}^{-3} \text{ pc}$. The shaded areas show the range of sensitivities for DMs between 50 and $150 \text{ cm}^{-3} \text{ pc}$. Note that the limiting flux density for the mid-latitude survey used for this curve is computed using the mean T_{sky} for the PMPS survey area. Also shown as diamonds are the periods and estimated flux densities for the four MSPs discovered in HTRU survey that lie in the PMPS region.

Table 6. Pulsars discovered in the HTRU survey for which archival data exists in the PMPS at the position of the pulsar. The unique PMPS grid identification number (see Manchester et al. 2001 for details) for the processed beam, the MJD of the observation and the S/N and periodicity if the pulsar was detected.

PSR J	Grid ID	MJD	S/N	Period (s)
1125–58	G4707513	51296.42	–	–
1622–4950	G4887499	50686.25	–	–
1625–49	G4891500	50686.23	7.9	0.355 8530
1701–44	G4924493	51152.94	6.4	0.755 5360
1705–43	G4928493	51457.10	10.4	0.222 5614
1708–35	G4959516	51718.68	17.4	0.004 5052
1754–24	G5021504	50843.07	–	–
1801–32	G4995477	52080.69	14.4	0.007 5344
1811–24	G5030488	51245.84	–	–

Of the remaining five pulsars, three are seen in the archival data with low S/N and none was selected for re-observation in the PMPS. Although the pulsar positions are not well known at this time, the nominal position of each falls more than 5 arcmin from the beam centre of the PMPS observation. Therefore we suggest that these pulsars fall in the regions between beam centres where the PMPS sensitivity is the lowest. Two pulsars appear to be below the detection limit of the PMPS including the newly discovered magnetar PSR J1622–4950, which shows extreme time variability (Levin et al. 2010).

7 CONCLUSION

We have described the system configuration and initial discoveries of a new pulsar survey using the Parkes Radio Telescope. At the time of writing, a total of 250 pulsars have been detected, of which 27 are new discoveries. The high time and frequency resolution of our digital backend system leads to increased sensitivity to short-period, high-DM pulsars compared to previous surveys. Indeed, five of our new discoveries are MSPs at high DM, all of which lie within previously surveyed regions of sky.

ACKNOWLEDGMENTS

The Parkes Observatory is part of the Australia Telescope which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. The HYDRA supercomputer at the JBCA was supported by a grant from the UK Science and Technology Facilities Council. SB-S gratefully acknowledges the support of STFC in his PhD studentship.

REFERENCES

- Archibald A. M. et al., 2009, *Sci*, 324, 1411
 Burgay M. et al., 2003, *Nat*, 426, 531
 Burke-Spolaor S., Bailes M., 2010, *MNRAS*, 402, 855
 Cordes J. M., Chernoff D. F., 1997, *ApJ*, 482, 971
 Cordes J. M., Lazio T. J. W., 2002, preprint (astro-ph/0207156)
 Eatough R. P., Keane E. F., Lyne A. G., 2009, *MNRAS*, 395, 410
 Eatough R. P., Molkenthin N., Kramer M., Noutsos A., Keith M. J., Stappers B. W., Lyne A. G., 2010, *MNRAS*, 407, 2443
 Edwards R. T., Bailes M., van Straten W., Britton M. C., 2001, *MNRAS*, 326, 358
 Haslam C. G. T., Klein U., Salter C. J., Stoffel H., Wilson W. E., Cleary M. N., Cooke D. J., Thomasson P., 1981, *A&A*, 100, 209
 Hessels J. W. T., Ransom S. M., Stairs I. H., Freire P. C. C., Kaspi V. M., Camilo F., 2006, *Sci*, 311, 1901
 Hewish A., Bell S. J., Pilkington J. D. H., Scott P. F., Collins R. A., 1968, *Nat*, 217, 709
 Hobbs G. B. et al., 2009, *Publ. Astron. Soc. Australia*, 26, 103
 Hotan A. W., van Straten W., Manchester R. N., 2004, *Publ. Astron. Soc. Australia*, 21, 302
 Jacoby B. A., Bailes M., Ord S. M., Edwards R. T., Kulkarni S. R., 2009, *ApJ*, 699, 2009
 Keith M. J., Johnston S., Kramer M., Weltevrede P., Watters K. P., Stappers B. W., 2008, *MNRAS*, 389, 1881
 Keith M. J., Eatough R. P., Lyne A. G., Kramer M., Possenti A., Camilo F., Manchester R. N., 2009, *MNRAS*, 395, 837
 Kramer M. et al., 2006, *Sci*, 314, 97
 Levin L. et al., 2010, *ApJL*, in press (arXiv:1007.1052)
 Lorimer D. R., Kramer M., 2005, *Handbook of Pulsar Astronomy*. Cambridge Univ. Press, Cambridge
 Lorimer D. R. et al., 2006, *MNRAS*, 372, 777
 Lorimer D. R., Bailes M., McLaughlin M. A., Narkevic D. J., Crawford F., 2007, *Sci*, 318, 777
 Lyne A. G. et al., 1998, *MNRAS*, 295, 743
 Lyne A. G. et al., 2004, *Sci*, 303, 1153
 McLaughlin M. A. et al., 2006, *Nat*, 439, 817
 McMahon P., 2008, Master's thesis, Univ. Cape Town
 Manchester R. N. et al., 1996, *MNRAS*, 279, 1235
 Manchester R. N. et al., 2001, *MNRAS*, 328, 17
 Staveley-Smith L. et al., 1996, *Publ. Astron. Soc. Australia*, 13, 243
 Taylor J. H., 1974, *A&AS*, 15, 367
 Verbiest J. P. W. et al., 2009, *MNRAS*, 400, 951
 Weltevrede P. et al., 2010, *ApJ*, 708, 1426
 Yusifov I., Küçük I., 2004, *A&A*, 422, 545

This paper has been typeset from a \LaTeX file prepared by the author.