A Mechanism-Informed Gene Signature for Regenerative vs. Fibrotic Tendon Healing

Mapping the transcription-factor program behind the fate decision in tendon progenitors — and compressing it into a validated 23-gene Healing Index.

Fangyi Li[1] · **Wilder Scott**[2] · **Yan Yan**[3]

[1]Department of Integrative Biology, College of Biological Science, University of Guelph · [2]Sunnybrook Research Institute · [3]School of Computer Science, College of Computational, Mathematical and Physical Sciences, University of Guelph

1 Introduction

Tendon heals badly: repair tissue is scar, not aligned tendon. Two progenitor populations inside the same PDGFRα⁺ pool decide which one forms — tendon stem/progenitor cells (TSPC) rebuild tendon, tendon fibro-adipogenic progenitors (T-FAP) lay down scar (Harvey et al., 2019).



3,900 PDGFRα⁺ cells from injured mouse tendon.
Sub-clustering resolves four populations (Cherief et al., 2023; day 14 post-injury).



Blocking sensory-nerve signalling shifts the pool.
TSPC share falls by more than half; T-FAP becomes the largest population. Nerve signalling is blocked chemical-genetically (TrkA-F592A mice + 1NMPP1), not surgically — "denervated" is shorthand throughout.

OBJECTIVE

Identify the transcription-factor program that carries out this innervation-dependent fate decision, then compress it into a minimal, validated gene signature that scores how regeneratively a tendon is healing.

2 Methods

Two independent public mouse scRNA-seq datasets, analysed through one integrated pipeline. No new animal work was performed.

Harvey et al. (2019)
Patellar tendon, uninjured · 4,069 cells after QC · SRA PRJNA506218
Ground-truth TSPC / T-FAP labels. Training set.

Cherief et al. (2023)
Achilles tendon, day 14 post-injury · 22,615 cells · GEO GSE244921
Innervated vs. TrkA-blocked. Held-out test set.

Analysis pipeline

1. **QC, clustering, annotation**
   PDGFRα⁺ pool (3,900 cells) sub-clustered into four populations
2. **pySCENIC — TF regulon inference**
   Run independently per dataset, then compared (Van de Sande et al., 2020)
3. **CellRank 2 — fate drivers**
   Genes correlated with commitment to each terminal population (Weiler et al., 2024)
4. **LIANA — cell–cell communication**
   Ligand–receptor signalling, innervated vs. blocked (Dimitrov et al., 2022)
5. **Feature selection**
   LASSO, Boruta and XGBoost run separately on 124 mechanism-derived candidates
6. **Classifier and Healing Index**
   Logistic regression trained on Harvey, applied unchanged to Cherief

Robustness checks

Every headline claim was re-tested. Two did not survive.

"Efna1→Epha3 is the strongest signal"
LIANA permutation test, n = 100
RETRACTED · p = 1.0

"T-FAP regulons feed back on Pdgfra"
Hypergeometric, BH-corrected
RETRACTED · nothing at p < 0.05

"Zero TSPC regulons replicate"
200× power-matched stability selection
CONFIRMED · still zero at matched n

3 Results

Which transcription-factor regulons separate the two fates, and do they replicate in a second, independent dataset?



Fifteen regulons replicate as fibrotic. None replicate as regenerative.
A 200-iteration power-matched audit reproduced the empty quadrant, so it is not a sample-size artefact.



bar length = CellRank fate-driver genes contributed · italic = upstream signal inferred from literature, not tested here

The surviving program is an AP-1 / KLF / NF-κB network.
Junb, Jund and Klf9 contribute the most fate-driver genes; 14 of the 23 final panel genes are their targets. 12 of the 15 regulons are shown — three contribute none.

KEY FINDING

The fate decision is asymmetric. Innervation appears to suppress a default fibrotic program rather than to activate a regenerative one — the mirror image of our starting hypothesis.

From mechanism to a gene signature



from 124 mechanism-derived candidates · 41 selected by none

Three methods agree on 23 genes.
LASSO, Boruta and XGBoost ran independently on 124 mechanism-derived candidates.



The panel generalises to a held-out dataset.
Trained on Harvey et al. (2019), then applied unchanged to Cherief et al. (2023) — independently labelled, never seen during selection or training.



The Healing Index tracks innervation status.
It also places two cell identities it never saw in training (Stromal, Tenogenic-progenitor) between the two fates.

4 Conclusions

A mechanism-derived 23-gene Healing Index now tracks the regenerative-to-fibrotic shift across two independent tendon datasets.

- Innervation suppresses a fibrotic program rather than activating a regenerative one: 15 T-FAP regulons replicate across datasets, 0 TSPC regulons do.
- A 23-gene panel separates the two fates at 0.989 ROC-AUC on a held-out, independently labelled dataset.
- The Healing Index falls from 25.2 to 15.5 when nerve signalling is blocked, and orders unseen cell identities sensibly.

LIMITATIONS · WHAT'S NEXT

- No reproducible regenerative regulon is not proof none exists — both datasets are small.
- Neither dataset has biological replicates — validation is cross-dataset, not cross-animal.
- TSPC recall is 77–78% vs. 98–99% for T-FAP: the index flags fibrotic drift better than it confirms repair.
- Next: score newly generated lab tendon scRNA-seq — the first true biological-replicate test.

Data & code availability
All data are publicly available (GEO GSE244921; SRA PRJNA506218). No unpublished or confidential data were used. Computation was performed on the Digital Research Alliance of Canada (NIBI). Code and analysis notebooks: github.com/lifangy6/sunnybrook-tendon-fate-mapping

Acknowledgements
We thank Dr. Karl Cottenie for course instruction and for feedback at every stage of this project, and Sabrina Saiphoo for her generosity with help and shared resources during our time in the same lab.

References

1. Cherief, M., et al. (2023). TrkA-mediated sensory innervation of injured mouse tendon supports tendon sheath progenitor cell expansion and tendon repair. Science Translational Medicine, 15(727), eade4619.
2. Dimitrov, D., et al. (2022). Comparison of methods and resources for cell-cell communication inference from single-cell RNA-Seq data. Nature Communications, 13, 3224.
3. Harvey, T., Flamenco, S., & Fan, C.-M. (2019). A Tppp3+Pdgfrα+ tendon stem cell population contributes to regeneration and reveals a shared role for PDGF signalling in regeneration and fibrosis. Nature Cell Biology, 21, 1490–1503.
4. Van de Sande, B., et al. (2020). A scalable SCENIC workflow for single-cell gene regulatory network analysis. Nature Protocols, 15, 2247–2276.
5. Weiler, P., Lange, M., Klein, M., Pe'er, D., & Theis, F. J. (2024). CellRank 2: Unified fate mapping in multiview single-cell data. Nature Methods, 21, 1196–1205.