

Sermorelin Acetate

Drug Class

Growth Hormone-Releasing Hormone (GHRH)

Mechanism of Action

Sermorelin is a synthetic peptide analog of GHRH consisting of the first 29 amino acids of the endogenous hormone. It stimulates the anterior pituitary to release endogenous growth hormone (GH), which in turn increases insulin-like growth factor 1 (IGF-1) levels.

Indications

- Diagnostic evaluation of pituitary function
- Off-label: Age-related GH decline, body composition improvement, sleep enhancement, and recovery support (under medical supervision)

Dosage and Administration

♦ Initial Titration Dosing

Phase	Dose Range	Frequency	Notes
Initiation	100–300 mcg	Daily at bedtime	Start low to assess tolerance
Adjustment	Increase by 50–100 mcg	Every 1–2 weeks	Based on IGF-1 levels and clinical response
Mainten.	300–500 mcg	5 nights/week	Common long-term protocol

- Route: Subcutaneous injection
- Timing: Administer at least 1–2 hours after last meal, ideally 30 minutes before bedtime to align with natural GH secretion

Monitoring Parameters

- IGF-1 levels (baseline and periodic)
- Sleep quality, energy levels, body composition
- Injection site reactions

Adverse Effects

- Common: Flushing, headache, nausea, injection site irritation
- Rare: Rash, chest tightness, breathing difficulty

Storage and Handling

- Reconstitute with bacteriostatic water
- Store refrigerated at 2–8°C (36–46°F)
- Use within 28 days of reconstitution