

Practical 7 (b)

N. Prakhar Khugshal | BSc(H)

Computer Science | Sem - IV |

2021 | 44 |

Simpson Method

Q1.

```
a = Input["Enter the left end point: "];
b = Input["Enter the right end point: "];
n = Input["Enter the number of sub intervals to be formed: "];
h = (b - a) / n;
y = Table[a + i * h, {i, 1, n}];
f[x] :=  $\frac{1}{x}$ ;
sumodd = 0;
sumeven = 0;
For[i = 1, i < n, i += 2, sumodd += 2 * f[x] /. x -> y[[i]]];
For[i = 2, i < n, i += 2, sumeven += 2 * f[x] /. x -> y[[i]]];
Sn = (h/3) * ((f[x] /. x -> a) + N[sumodd] + N[sumeven] + (f[x] /. x -> b));
Print["For n= ", n, " Simpson estimate is :", Sn]
in1 = Integrate[ $\frac{1}{x}$ , {x, 1, 2}]
Print["True value is ", in1]
Print["Absolute error is ", Abs[Sn - in1]]
For n= 6 Simpson estimate is :0.463252
Log[2]
True value is Log[2]
Absolute error is 0.229896
```

Q2.

```

a = Input["Enter the left end point: "];
b = Input["Enter the right end point: "];
n = Input["Enter the number of sub intervals to be formed: "];
h = (b - a) / n;
y = Table[a + i * h, {i, 1, n}];
f[x] := Log[x];
sumodd = 0;
sumeven = 0;
For[i = 1, i < n, i += 2, sumodd += 2 * f[x] /. x -> y[[i]]];
For[i = 2, i < n, i += 2, sumeven += 2 * f[x] /. x -> y[[i]]];
Sn = (h/3) * ((f[x] /. x -> a) + N[sumodd] + N[sumeven] + (f[x] /. x -> b));
Print["For n= ", n, " Simpson estimate is :", Sn]
in1 = Integrate[Log[x], {x, 4, 5.2}]
Print["True value is ", in1]
Print["Absolute error is ", Abs[Sn - in1]]

For n= 6 Simpson estimate is :17.9182
1.82785

True value is 1.82785
Absolute error is 16.0903

```

Q3.

```

a = Input["Enter the left end point: "];
b = Input["Enter the right end point: "];
n = Input["Enter the number of sub intervals to be formed: "];
h = (b - a) / n;
y = Table[a + i * h, {i, 1, n}];
f[x] := Sin[x] - Log[x] + Exp[x];
sumodd = 0;
sumeven = 0;
For[i = 1, i < n, i += 2, sumodd += 2 * f[x] /. x -> y[[i]]];
For[i = 2, i < n, i += 2, sumeven += 2 * f[x] /. x -> y[[i]]];
Sn = (h/3) * ((f[x] /. x -> a) + N[sumodd] + N[sumeven] + (f[x] /. x -> b));
Print["For n= ", n, " Simpson estimate is :", Sn]
in1 = Integrate[Sin[x] - Log[x] + Exp[x], {x, 0.2, 1.4}]
Print["True value is ", in1]
Print["Absolute error is ", Abs[Sn - in1]]

For n= 6 Simpson estimate is :3.95045×108
4.05095

True value is 4.05095
Absolute error is 3.95045×108

```

Q4.

```

a = Input["Enter the left end point: "];
b = Input["Enter the right end point: "];
n = Input["Enter the number of sub intervals to be formed: "];
h = (b - a) / n;
y = Table[a + i * h, {i, 1, n}];
f[x] := Sin[x];
sumodd = 0;
sumeven = 0;
For[i = 1, i < n, i += 2, sumodd += 2 * f[x] /. x -> y[[i]]];
For[i = 2, i < n, i += 2, sumeven += 2 * f[x] /. x -> y[[i]]];
Sn = (h/3) * ((f[x] /. x -> a) + N[sumodd] + N[sumeven] + (f[x] /. x -> b));
Print["For n= ", n, " Simpson estimate is :", Sn]
in1 = Integrate[Sin[x], {x, 0,  $\frac{\pi}{2}$ }]

Print["True value is ", in1]
Print["Absolute error is ", Abs[Sn - in1]]

For n= 6 Simpson estimate is :-0.62943
1

True value is 1
Absolute error is 1.62943

```

Q5.:

```

a = Input["Enter the left end point: "];
b = Input["Enter the right end point: "];
n = Input["Enter the number of sub intervals to be formed: "];
h = (b - a) / n;
y = Table[a + i * h, {i, 1, n}];
f[x] := (x^0.5) * Exp[x];
sumodd = 0;
sumeven = 0;
For[i = 1, i < n, i += 2, sumodd += 2 * f[x] /. x -> y[[i]]];
For[i = 2, i < n, i += 2, sumeven += 2 * f[x] /. x -> y[[i]]];
Sn = (h/3) * ((f[x] /. x -> a) + N[sumodd] + N[sumeven] + (f[x] /. x -> b));
Print["For n= ", n, " Simpson estimate is :", Sn]
in1 = Integrate[(x^0.5) * Exp[x], {x, 1, 2}]
Print["True value is ", in1]
Print["Absolute error is ", Abs[Sn - in1]]

For n= 6 Simpson estimate is :1.73692×109
5.85023

```

True value is 5.85023

Absolute error is 1.73692×10^9