

CUDA-based Geant4 Monte Carlo Simulation for Radiation Therapy

N. Henderson & K. Murakami


GTC 2013

The collaboration

Geant4 @ **SLAC**



INSTITUTE for COMPUTATIONAL &
MATHEMATICAL ENGINEERING
at STANFORD UNIVERSITY



Makoto Asai, SLAC
Joseph Perl, SLAC
Koichi Murakami, KEK-SLAC
Takashi Sasaki, KEK
Margot Gerritsen, ICME
Nick Henderson, ICME


Special thanks to the
CUDA Center of
Excellence Program

20,000,000

radiotherapy treatments per year in US



Elekta Synergy Platform



Simulation methods

Analytic

- time: seconds to minutes
- accurate within 3-5%
- used in treatment planning

Monte Carlo


- time: several hours to days of CPU time
- accurate within 1-2%
- used to verify treatment plans in certain cases

Geant4 Toolkit

- Enables Monte Carlo simulation of particles travelling through and interacting with matter
- Allows modeling of complex geometries
- Covers all elementary particles and nuclei for a wide energy range


Geant4 Applications

High Energy Physics




ATLAS

Space & Radiation



LISA


Medical Physics



gMocren

Geant4 101

- Geant4 simulates particles travelling through and interacting with matter
- Example: photoelectric effect



Parallelization challenges in Geant4

- Large and complex code base
- Sophisticated geometry framework
- Elaborate physics models
- Branching, look-up tables, single-thread optimizations

Thank goodness

The simulation is embarrassingly parallel!
(the particles are independent)

Requirements for X-ray Radiotherapy

- Geometry is a voxelized box
- Physics is limited to low-energy electromagnetics
- Material is modeled as water with different densities

Low energy electromagnetics

Gamma

- Compton scattering
- Photoelectric effect
- Gamma conversion




Image: CCSA by Manticorp on [Wikipedia](#)

Electron/positron

- Ionization
- Bremsstrahlung
- Positron annihilation
- **Todo: Multiple scattering**





Image: public domain

What makes a physics process?

- Sample for angular and energy distributions of secondary particles
- Deposit energy to material
- Produce secondary particles that must be tracked at a later point


Tracking algorithm

- Particles are tracked through space
- Each discrete move is called a step
- Physics process may occur along step, after the step, or both




How are processes selected?

- Each process has an “interaction length” (dx)
- Process with shortest dx is selected
- After the step dx is decreased or resampled



Energy deposition

- Happens along the step for the ionization process
- May happen at the end of the step
- Secondary particles with too low energy are not generated, but treated as point-like energy depositions



G4CU: CUDA-base MC for RT


- Important data structures
- Algorithm summary
- Details
 - Parallel stack
 - Look-up tables

Struct-of-arrays data pattern

```
struct ParticleArray {  
    // length of arrays  
    int length;  
    // kind of particle  
    ParticleKind *kind;  
    // position  
    float *x, *y, *z;  
    // direction  
    float *dx, *dy, *dz;  
    // particle energy  
    float *energy;  
    // voxel index  
    int *vx, *vy, *vz, *vid;  
};
```

- Common pattern in CUDA to allow for coalesced memory access
- Experiments with transport showed this to be **3-4x faster** than AOS


G4CU: algorithm



Some details

- A process does a few things:
 1. Changes direction and momentum of primary particle
 2. Generates secondary particles
 3. Deposits energy to the material
- We use thread local stacks to handle 2 and 3


Parallel stacks



```
template <typename T>
struct pstack {
    // number of stacks
    int stack_num;
    // size of each stack
    int stack_size;
    // starts and ends of stacks
    int *start, *end;
    // stack positions
    int *pos;
    // stack data array
    T *data;
};
```


Dose reduction & storage

- Energy dose is stored in a thread local variable
- When particle moves to new voxel, dose is pushed onto stack
- Dose reduction is performed periodically
- We use Thrust



Look-up tables

- Used for interaction length computation
- 40 bins, log spaced
- Linear and spline interpolation
- Bremsstrahlung also uses 2D interpolation



Benchmarks

Configuration:

- Geometry: 512 x 512 x 256 voxels
- Dose reduction frequency: every 200 iterations
- 128 blocks with 256 threads
- Primary particle is a 6 MeV gamma
- Tesla C2070

Simulation time

- 100 million primary particles
- Time: 72 minutes
- ~ 23.1 primary particles per ms
- ~ 50 - 60 x speedup over Geant4 on 1 CPU


Profile

Component	Percentage of overall time
Physics processes	50
Energy dose reduction	30
Interaction length	18
Run management	2

Physics process breakdown

Process	Particle	Percentage of overall time
Bremsstrahlung	e^- / e^+	23
Pair production	γ	7.5
Transport	all	7
Photoelectric effect	γ	7
Ionization	e^- / e^+	3
Compton scattering	γ	1.5
Positron annihilation	e^+	1

Dose distribution



Acknowledgements

Geant4 Collaboration, see

- **Geant4—a simulation toolkit**, *Nuclear Instruments and Methods in Physics Research A* **506** (2003) 250-303
- **Geant4 developments and Applications**, *IEEE Transactions on Nuclear Science* **53** No. 1 (2006) 270-278

NVIDIA & CCOE Program

People:

- Koichi Murakami, KEK
- Makoto Asai, SLAC
- Joseph Perl, SLAC
- Takashi Sasaki, KEK
- Margot Gerritsen, ICME