

ISKRA – 2017

**ПРОНИКНОВЕНИЕ КОСМИЧЕСКИХ
ЛУЧЕЙ В ДИНАМИЧЕСКУЮ
МАГНИТОСФЕРУ И ИХ ВЛИЯНИЕ НА
АТМОСФЕРУ**


by Lev I. Dorman (1, 2)

**1. Cosmic Ray Department of N.V. Pushkov's IZMIRAN
Russian Academy of Science**

**2. Israel Cosmic Ray and Space Weather Center with
Emilio Ségre Observatory, affiliated to Tel Aviv
University, Shamir Research Institute, and Israel Space
Agency**

E-mail: lid010529@gmail.com

Changes in the distribution of vertical planetary geomagnetic cutoff rigidity of the era of 2015 with



Digital and graphical results of calculations can be found at

ftp Server [Magnetosphere Effect](#). (user name : anonymous)

Accounting for the penumbra region



Excluding the effects of the atmosphere. For the primary spectrum, for example, $J(R) = aR^\gamma$, the effective stiffness of geomagnetic cutoff is determined from the equation:

$$\int_{R_{eff}}^{R_H} J(R) dR = \int_{R_S}^{R_H} g(R) J(R) dR \cdots \Rightarrow R_{eff}^{\gamma+1} = R_H^{\gamma+1} - (\gamma + 1) \int_{R_S}^{R_H} g(R) \cdot R^\gamma dR$$

In the case of the primary spectrum is not dependent on energy ("white" or "flat")

$$R_{eff} = R_H - \int_{R_S}^{R_H} g(R) dR$$

Into account the atmosphere. In this case the communication functions in the penumbra $W(R) = cR^\eta$, the effective stiffness of geomagnetic cutoff is determined from the equation:

$$\int_{R_{eff}}^{R_H} W(R, h) \cdot \delta J / J(R) \cdot dR = \int_{R_S}^{R_H} g(R) \cdot W(R, h) \cdot \delta J / J(R) \cdot dR$$

$$R_{eff}^{\gamma+\eta+1} = R_H^{\gamma+\eta+1} - (\gamma + \eta + 1) \int_{R_S}^{R_H} g(R) \cdot R^{\eta+\gamma} dR$$

Accounting for the penumbra region








Illustration methodology for assessing the effective stiffness of circumcission to account penumbra of cosmic rays.

Changes of vertical geomagnetic cut off rigidity



Changes of vertical geomagnetic cut off rigidity for 4 groups of the CR stations relatively to the epoch 1950 (in the approach IGRF and flat spectra of CR variations)

We consider how galactic and solar cosmic rays (CR) influence on the physics and chemistry of the Earth's atmosphere and atmospheric processes:

1. Through nuclear reactions of primary and secondary CR with air and aerosol matter accompanied by formation of many unstable and stable cosmogenic nuclides, especially C-14 (radiocarbon) and Be-10.
2. Through generation in the atmosphere of secondary relativistic electrons and EAS (Extensive Atmospheric Showers) playing a crucial role in atmospheric electric field phenomena: thunder-storms, discharges, the Earth's electric charge balance.
3. Through air ionization influences on the low ionosphere and radio wave propagation, as well as on the formation of clouds and influence on long-term global climate change, and wheat production and prices.
4. Through chemical reactions induced by interactions of CR with air atoms including production of nitrates and influence on ozone layer.

Contents

- **1. Nuclear reactions of cosmic rays in the Earth's atmosphere and production of cosmogenic nuclides**
- **2. Cosmic ray Influence on atmospheric electric field and thunderstorms, Global Earth's charge and global electric current**
- **3. Cosmic ray Influence on the chemical processes in the atmosphere and formation of nitrates and influence on ozone layer**
- **4. Cosmic ray influence on planetary cloud-covering and long-term climate change**

1. Nuclear reactions of cosmic rays in the Earth's atmosphere and production of cosmogenic nuclides

- 1.1. Cosmogenic nuclides and vertical mixing of elements in the Earth's atmosphere; local cosmogenic coupling functions
- 1.2. The planetary mixing in the atmosphere, variations in planetary cosmogenic nuclides production rate and planetary coupling functions

$$q_i(R_c, h, t) = \sum_Z \int_{R_c}^{\infty} D_Z(R, t) dR \sum_l \sum_k \int_0^{E(R)} M_l(R, Z_l, E_l, h) \sigma_{ilk}(E_l) N_k(h) dE_l$$

$$M_i(R, h, t) = \sum_Z \sum_l \sum_k \int_0^{E(R)} M_l(R, Z_l, E_l, h) \sigma_{ilk}(E_l) B_Z N_k(h, t) dE_l$$

$$Q_i(R_c, h_o, t) = T_{iv}^{-1} \sum_Z \int_{R_c}^{\infty} dR \int_{t-T_{iv}}^t D_Z(R, t) M_{Zi}(R, h_o, t) dt$$

$$M_{Zi}(R, h_o, t) = \int_0^{h_o} dh \sum_l \sum_k \int_0^{E(R)} M_{Zl}(R, Z_l, E_l, h) \sigma_{ilk}(E_l) N_k(h, t) dE_l$$

$$W_{iZR_{co}}(R, h_o) = D_{Zo}(R) M_{iZo}(R, h_o) / Q_{io}(R_{co}, h_o)$$

$$Q_{ip}(R_{cp},h_{op},t)=\left(4\pi T_{ip}\right)^{-1}\int\limits_{t-T_{ip}}^td\tau\int\limits_{-\pi/2}^{\pi/2}\sin\theta d\theta\int\limits_0^{2\pi}d\varphi Q_i(R_c(\theta,\varphi,\tau),h_o(\theta,\varphi,\tau),\tau)$$

$$R_c(\theta,t)=R_c(\theta,0)M_p(t)/M_p(0)$$

$$W_{ip}(R,h_{opo})=D_o(R)M_{ip}(R,h_{opo})/Q_{ipo}$$

$$d\,N_{ip1}(t)/dt=Q_{ip}\big(R_{cp}(t),h_{op}(t),t\big)-\big(\lambda_i+\lambda_{12}\big)N_{ip1}(t)+\lambda_{21}N_{ip2}(t),$$

$$d\,N_{ip2}(t)/dt=-\big(\lambda_i+\lambda_{21}\big)N_{ip2}(t)+\lambda_{12}N_{ip1}(t).$$

H-bombs explosions, generation of radiocarbon, and estimation of parameters of the elements exchange model; influence on global environment

$$Q_p(t) = Q_{av} (1 + A\Delta t \delta(t - t_o))$$

$$N_A(t) = N_{Ast} \left[1 + \frac{\lambda \lambda_1 A \Delta t \exp(-\lambda(t - t_o))}{\lambda_2(\lambda + \lambda_{FA})} (\lambda_{AF} \exp(-\lambda_2(t - t_o)) + \lambda_{FA}) \right]$$

$$N_F(t) = N_{Fst} \left[1 + \frac{\lambda_1 A \Delta t \exp(-\lambda(t - t_o))}{\lambda_2} (1 - \exp(-\lambda_1(t - t_o))) \right]$$

$$N_{A\max} = N_{Ast} (1 + \lambda A \Delta t (N_{st} / N_{Ast})) = N_{Ast} (1 + 9.98 \times 10^{-3} A \Delta t)$$

Application to USSR and USA H-bomb explosions in 1962

- $9.98 \times 10^{-3} A \Delta t \approx 0.9, \quad A = 90.2$

$$6.83 \times 10^9 \text{ atom } ^{14}\text{C.cm}^{-2} \quad 7.48 \times 10^7$$

$$\lambda_{AF} = 0.145 \text{ year}^{-1}, \quad \lambda_{FA} = 1.66 \times 10^{-3} \text{ year}^{-1}$$

$$t_{\max F} = t_o + 48.3 \text{ years} = 2010.3 \text{ years}$$

$$N_{F \max} = N_{Fst} \left(1 + 5.31 \times 10^{-3} A \Delta t \right) = N_{Fst} (1 + 0.479)$$

The basic equations for the 5-reservoir model

$$dN_A(t)/dt = Q_p(t) - (\lambda + \lambda_{AM} + \lambda_{AB})N_A(t) + \lambda_{HA}N_H(t) + \lambda_{MA}N_M(t)$$

$$dN_B(t)/dt = -(\lambda + \lambda_{BH})N_B(t) + \lambda_{AB}N_A(t)$$

$$dN_H(t)/dt = -(\lambda + \lambda_{HA})N_H(t) + \lambda_{BH}N_B(t)$$


$$dN_M(t)/dt = -(\lambda + \lambda_{MO} + \lambda_{MA})N_M(t) + \lambda_{AM}N_A(t)$$

$$dN_O(t)/dt = -(\lambda + \lambda_{OM})N_O(t) + \lambda_{MO}N_M(t)$$


2. Cosmic Ray Influence on Atmospheric Electric Field and Thunderstorms, Global Earth's Charge and Global Electric Current

- 1. On two mechanisms of CR connection with thunderstorm discharges
- 2. Necessary conditions for atmospheric electric field discharges in the atmosphere
- 3. Measurements of atmospheric electric field, critical electric field, lightnings, and sprites


18 June 1987, Hinton, OK




30 May 1988, Dalhart, TX




2 August 1984, Langmuir Lab., NM



6 August 1984, Langmuir Lab., NM




Time variations for 17 years observations of discharged electric current J (light circles) according to Roble (1985) and of CR intensity $I(h)$ (black circles) according to Stozhkov et al. (2001).



The annual number of thunderstorm discharges N (open circles) over the territory of the USA in 1989–1998 according to Orville and Huffines (1999) and time variation of yearly average ion production rate q (black circles) in the column of atmosphere between 2 and 10 km (calculated on the basis of balloon CR measurements).

According to Stozhkov et al. (2001).



On the CR role in the equilibrium between charged and discharged global atmospheric electric currents, and in the supporting the stability of the Earth's Charge

$$J_{CR} = eI_{CR} \approx 1.6 \times 10^{-19} \text{ C} \times 10^4 \text{ m}^{-2} \text{ sec}^{-1} \approx 2 \times 10^{-15} \text{ A/m}^2$$


is negligible in comparison with currents caused by the global thunderstorm activity

$$J \approx 10^{-12} \text{ A/m}^2$$

From other hand, the increase of CR intensity I (increase of ion production rate q) leads to an increase of discharged global electric currents. This will mostly compensate the increasing of charged electric currents caused by increasing thunderstorms discharges frequency (caused by the same increasing of CR intensity).


Air Ionization by CR, Influence on the Ionosphere and Radio Wave Propagation

Total ionization during GLE in October 1989, July 2000, and April 2001. According to Quack et al. (2001).




3. Cosmic Ray Influence on the Chemical Processes in the Atmosphere and Formation of Ozone Layer

- NO production by CR (Crutzen et al., 1975).




McCracken et al. (2001)




CR influence on stratospheric chemistry

Modeled of NO_x during the ground level event on April 14, 2001.

According to Quack et al (2001).




Percentage decrease of the O₃ partial pressure versus air pressure derived from the average of the 7 days before 4 August 1972 and 7 day periods centered on 8 and 19 days after the GLE (solid lines). According to Heath et al. (1977).





Variations compared to background level in O₃ in % (down panel modeled and upper panel observed) during the event in July 2000.


According to Quack et al (2001).






4. Cosmic Ray Influence on Planetary Cloud-Covering and Long-Term Climate Change












FIGURE 6 Wheat prices in England during 1259–1702 with a price transition at 1530–1630. From Refs. [51,52].








FIGURE 7 Histograms of the interval distribution for price bursts for the period, 1249–1702, and of minimum–minimum intervals of sunspots during 1700–2000. From Refs. [51,52].










TABLE 2 Vertical cutoff rigidities (in GV) for various epochs 1600, 1700, 1800, 1900 and 2000, as well as change from 1900 to 2000 owed to changes of geomagnetic field. According to Shea and Smart [73]

Lat.	Long. (E)	Epoch 2000	Epoch 1900	Epoch 1800	Epoch 1700	Epoch 1600	Change 1900–2000	Region
55	30	2.30	2.84	2.31	1.49	1.31	−0.54	Europe
50	0	3.36	2.94	2.01	1.33	1.81	+0.42	Europe
50	15	3.52	3.83	2.85	1.69	1.76	−0.31	Europe
40	15	7.22	7.62	5.86	3.98	3.97	−0.40	Europe
45	285	1.45	1.20	1.52	2.36	4.1	+0.25	North America
40	255	2.55	3.18	4.08	4.88	5.89	−0.63	North America
20	255	8.67	12.02	14.11	15.05	16.85	−3.35	North America
20	300	10.01	7.36	9.24	12.31	15.41	+2.65	North America
50	105	4.25	4.65	5.08	5.79	8.60	−0.40	Asia
40	120	9.25	9.48	10.24	11.28	13.88	−0.23	Asia
35	135	11.79	11.68	12.40	13.13	14.39	+0.11	Japan
−25	150	8.56	9.75	10.41	11.54	11.35	−1.19	Australia
−35	15	4.40	5.93	8.41	11.29	12.19	−1.53	South Africa
−35	300	8.94	12.07	13.09	10.84	8.10	−3.13	South America


Conclusions

- **1. The atmosphere influenced very strong on CR: meteorological effects, cascade processes, generation of new particles, secondary components, albedo radiation.**
- **2. From other hand, CR influenced on the atmosphere and atmospheric processes.**
- **3. This influence is determined by nuclear reactions of primary and secondary CR (mostly, neutrons) with air and aerosol matter accompanied by the formation of many unstable and stable cosmogenic nuclides Be-10, C-14, and many others.**
- **4. It determined by generation in the atmosphere by CR of secondary relativistic electrons and EAS (Extensive Atmospheric Showers) playing a crucial role in atmospheric electric field phenomena.**
- **5. It determined by CR air ionization influences on the low ionosphere and radio wave propagation.**
- **6. It determined by induced chemical reactions, influences on the chemistry of the atmosphere and the ozone layer.**
- **7. It determined by possible influence on long-term global climate change through influence on the formation of clouds (mostly, low clouds).**

- 8. We take into account big variability of CR intensity caused by modulation in the Heliosphere, generation of solar energetic particles in flare events, and by influence of changing geomagnetic cutoff rigidities.
- 9. Planetary observations on balloons of different components of primary and secondary components of CR as well as air isotopic and chemical contents in dependence of latitude and altitude are crucial for understanding of the nature of CR influence on physical and chemical processes in the Earth's atmosphere as well as for space weather and global climate change problems.

Main references

1. Dorman L.I., *Cosmic Rays in the Earth's Atmosphere and Underground*, Kluwer Academic Publishers, Dordrecht/London/Boston, pp 862 (2004).
- 2. Dorman L.I., *Cosmic Ray Interactions, Propagation, and Acceleration in Space Plasmas*, Springer, Netherlands, pp 847 (2006).
- 3. Dorman L.I., *Cosmic Rays in Magnetospheres of the Earth and other Planets*, Springer, Netherlands, pp 770 (2009).
- 4. Dorman L.I. "Chapter 3. The Role of Space Weather and Cosmic Ray Effects in Climate Change", in book *Climate Change: Observed Impacts on Planet Earth*, Edited by Trevor M. Letcher, Elsevier, 2009



20% conference discount

Lev I. Dorman, Israel Cosmic Ray Center, Space Weather Center, and Emilio Segrè Observatory, affiliated to Tel Aviv University, Israel Space Agency, and Technion, Qazrin, Israel; and Cosmic Ray Dept. of IZMIRAN, Russian Academy of Science, Troitsk, Russia

Cosmic Rays in the Earth's Atmosphere and Underground

This book consists of four parts. In the first part (Chapters 1-4) a full overview is given of the theoretical and experimental basis of Cosmic Ray (CR) research in the atmosphere and underground for Geophysics and Space Physics; the development of CR research and a short history of many fundamental discoveries, main properties of primary and secondary CR, methods of transformation of CR observation data in the atmosphere and underground to space, and the experimental basis of CR research underground and on the ground, on balloons and on satellites and space probes.

The second part (Chapters 5-9) is devoted to the influence of atmospheric properties on CR, so called CR meteorological effects; pressure, temperature, humidity, snow, wind, gravitation, and atmospheric electric field effects. The inverse problem - the influence of CR properties on the atmosphere and atmospheric processes is considered in the third part (Chapters 10-14); influence on atmospheric, nuclear and chemical compositions, ionization and radio-wave propagation, formation of thunderstorms and lightning, clouds and climate change.


The fourth part (Chapters 15-18) describes many realized and potential applications of CR research in different branches of Science and Technology; Meteorology and Aerodrome Service, Geology and Geophysical Prospecting, Hydrology and Agricultural Applications, Archaeology and Medicine, Seismology and Big Earthquakes Forecasting, Space Weather and Environment Monitoring/Forecasting.

The book ends with a list providing more than 1,500 full references, a discussion on future developments and unsolved problems, as well as object and author indices. This book will be useful for experts in different branches of Science and Technology, and for students to be used as additional literature to text-books.

Contents: 1: Cosmic Rays As An Object Of Research And As A Research Instrument.- 2: Influence Of The Changing Atmosphere On Cosmic Rays (Meteorological Effects).- 3: Cosmic Ray Influence On The Atmosphere And Atmospheric Processes.- 4: Applications Of Cosmic Ray Research.- Conclusion and Problems.

2004, 862 p. Hardcover

• € 175.00 | £ 121.00 • Conference price: € 140.00 | £ 96.80 Valid until 31 December 2009. This applies to personal orders by conference delegates using this order form.



20% conference discount

Lev I. Dorman, *Israel Cosmic Ray Center, Space Weather Center, and Emilio Segrè Observatory, affiliated to Tel Aviv University, Israel Space Agency, and Technion, Qazrin, Israel; and Cosmic Ray Dept. of IZMIRAN, Russian Academy of Science, Troitsk, Russia*

Cosmic Ray Interactions, Propagation, and Acceleration in Space Plasmas

The book consists of four Chapters. Chapter 1 shortly describes main properties of space plasmas and primary CR, different types of CR interactions with space plasmas components (matter, photons, and frozen in magnetic fields). Chapter 2 considers the problem of CR propagation in space plasmas described by the kinetic equation and different types of diffusion approximations (diffusion in momentum space and in pitch-angle space, anisotropic diffusion, anomaly CR diffusion and compound diffusion, the influence of magnetic clouds on CR propagation, non-diffusive CR particle pulse transport). Chapter 3 is devoted to CR non-linear effects in space plasmas caused by CR pressure and CR kinetic stream instabilities with the generation of Alfvén turbulence (these effects are important in galaxies, in the Heliosphere, in CR and gamma-ray sources and in the processes of CR acceleration). In Chapter 4 different processes of CR acceleration in space plasmas are considered: the development of the Fermi statistical mechanism, acceleration in the turbulent plasma, Alfvén mechanism of magnetic pumping, induction mechanisms, acceleration during magnetic collapse and compression, cumulative acceleration mechanism near the zero lines of a magnetic field, acceleration in shear flows, shock-wave diffusion (regular) acceleration. The book ends with a list providing more than 1,300 full references, a discussion on future developments and unsolved problems, as well as Object and Author indexes. This book will be useful for experts and students in CR research, Astrophysics and Geophysics, and in Space Physics.

Contents: Preface. Acknowledgments. Frequently used Abbreviations and Notations.- 1. Cosmic Ray Interactions in Space Plasmas.- 2. Cosmic Ray Propagation in Space Plasmas.- 3. Cosmic Ray Nonlinear Effects in Space Plasmas.- 4. Cosmic Ray Acceleration in Space Plasmas.- Conclusion and Problems.- References. Object Index. Author Index.


2006, 847 p. Hardcover
Astrophysics and Space Science Library, Volume 339

• € 139.95 | £ 107.50 • Conference price: € 111.96 | £ 86.00

L. Dorman, Cosmic Rays Research Center, Tel Aviv, Israel

Cosmic Rays in Magnetospheres of the Earth and other Planets


This monograph describes the behaviour of cosmic rays in the magnetosphere of the Earth and of some other planets. Recently this has become an important topic both theoretically, because it is closely connected with the physics of the Earth's magnetosphere, and practically, since cosmic rays determine a significant part of space weather effects on satellites and aircraft. The book contains eight chapters, dealing with – The history of the discovery of geomagnetic effects caused by cosmic rays and their importance for the determination of the nature of cosmic rays or gamma rays – The first explanations of geomagnetic effects within the framework of the dipole approximation of the Earth's magnetic field – Trajectory computations of cutoff rigidities, transmittance functions, asymptotic directions, and acceptance cones in the real geomagnetic field taking into account higher harmonics – Cosmic ray latitude-longitude surveys on ships, trains, tracks, planes, balloons and satellites for determining the planetary distribution of the intensity of cosmic rays and how it changes with time – Geomagnetic time variations of cosmic rays caused by the changing of internal sources of Earth's main magnetic field and variable... *more on* <http://springer.com/978-1-4020-9238-1>



2009. XXXIV, 770 p. (Astrophysics and Space Science Library, Vol. 358) Hardcover

► **179,95 €**

ISBN 978-1-4020-9238-1



All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are prohibited. For exceptions, permission may be sought for such use through Elsevier's permissions site at:

<http://www.elsevier.com/locate/permissionusematerial>

From: Lev I. Dorman, The Role of Space Weather and Cosmic Ray Effects in Climate Change. In Trevor M. Letcher, editor: Climate Change: Observed impacts on Planet Earth, The Netherlands, 2009, pp. 43-76.

ISBN: 978-0-444-53301-2

The Role of Space Weather and Cosmic Ray Effects in Climate Change

Lev I. Dorman

*Head of Cosmic Ray and Space Weather Center with Emilio Sègre Observatory, affiliated to Tel Aviv University, TECHNION and Israel Space Agency, P.O. Box 2217, Qazrin 12900, Israel
Chief Scientist of Cosmic Ray Department of IZMIRAN Russian Academy of Science, Troitsk 142090, Moscow Region, Russia*

1. **Introduction**
2. **Solar Activity, Cosmic Rays and Climate Change**
 - 2.1. Long-Term Cosmic Ray Intensity Variations and Climate Change
 - 2.2. The Possible Role of Solar Activity and Solar Irradiance in Climate Change
 - 2.3. Cosmic Rays as an Important Link between Solar Activity and Climate Change
 - 2.4. The Connection between Galactic Cosmic Ray Solar Cycles and the Earth's Cloud Coverage
 - 2.5. The Influence of Cosmic Rays on the Earth's Temperature
 - 2.6. Cosmic Ray Influence on Weather during Maunder Minimum
 - 2.7. The Influence of Long-Term Variations of Cosmic Ray Intensity on Wheat Prices (Related to Climate Change) in Medieval England and Modern USA
 - 2.8. The Connection between Ion Generation in the Atmosphere by Cosmic Rays and Total Surface of Clouds
 - 2.9. The Influence of Big Magnetic Storms (Forbush Decreases) and Solar Cosmic Ray Events on Rainfall
 - 2.10. The Influence of Geomagnetic Disturbances and Solar Activity on the Climate through Energetic Particle Precipitation from Inner Radiation Belt

- 2.11. On the Possible Influence of Galactic Cosmic Rays on Formation of Cirrus Hole and Global Warming
- 2.12. Description of Long-Term Galactic Cosmic Ray Variation by both Convection-Diffusion and Drift Mechanisms with Possibility of Forecasting of Some Part of Climate Change in Near Future Caused by Cosmic Rays
- 2.13. Influence of Long-Term Variation of Main Geomagnetic Field on Global Climate Change through Cosmic Ray Cutoff Rigidity Variation
- 2.14. Atmospheric Ionisation by Cosmic Rays: The Altitude Dependence and Planetary Distribution
- 2.15. Project 'Cloud' as an Important Step in Understanding the Link between Cosmic Rays and Cloud Formation
- 3. The Influence on the Earth's Climate of the Solar System Moving Around the Galactic Centre and Crossing Galaxy Arms**
- 4. The Influence of Molecular-dust Galactic Clouds on the Earth's Climate**
- 5. The Influence of Interplanetary Dust Sources on the Earth's Climate**
- 6. Space Factors and Global Warming**
- 7. The Influence of Asteroids on the Earth's Climate**
- 8. The Influence of Nearby Supernova on the Earth's Climate**
- 9. Discussion and Conclusions
Acknowledgments
References**