

MATH240 Summer 2022

Haoran Li

Contents

1	Worksheets	2
1.1	Worksheet for Section 6.1	2
1.2	Worksheet for Section 6.4	4
1.3	Worksheet for Section 6.5	6
1.4	Worksheet for Sections 6.7, 6.2, and 6.8	7
1.5	Sample Exam 1 MATH 141 - Calculus II	9
1.6	Worksheet for Section 7.1	11
1.7	Worksheet for Section 7.2	13
1.8	Worksheet for Section 7.3 and 7.5	15
1.9	Worksheet for Section 7.6	17
1.10	Worksheet for Section 7.6 and 8.1	19
1.11	Sample Exam 2 MATH 141 - Calculus II	21
1.12	Worksheet for Section 8.2	23
1.13	Worksheet for Section 8.3	24
1.14	Worksheet for Section 8.4 and 8.6	26
1.15	Worksheet for Section 8.6 and 8.7	28
2	Exams	31
2.1	Exam 1	31
2.2	Exam 2	32

1 Worksheets

1.1 Worksheet for Section 6.1

1. A cylindrical tank standing upright, with a volume of 40π cubic meters, is to be built. Materials for the sides cost 10 dollars per square meter, and for the top and bottom the materials cost 25 dollar per square meter. Find the dimension that will minimize the cost.
2. Let $g(x) = \int_x^{x+\pi} \sin^{2/3} t dt$, for any given value of $x \geq 0$. Show that g is a constant function.
3. Consider the solid region whose base R is bounded by the negative x -axis, the positive y -axis, and the curve $y = 4 - x^2$ for $-2 \leq x \leq 0$.
 - (a) Let D_1 be the solid with base R , and assume that the cross sections of D_1 perpendicular to the x -axis are squares. Draw a picture of the base, and then draw a representative cross section perpendicular to the x -axis at some arbitrary x in the interval $(-2, 0)$. Finally, find the cross-sectional area A_1 of the cross section.
 - (b) Now let D_2 be the solid with base R , and assume that the cross sections of D_2 perpendicular to the x -axis are semi-circles. Draw a second picture of the base, and then draw a representative cross section perpendicular to the x -axis at some arbitrary x in the interval $(-2, 0)$. Finally, find the cross-sectional area A_2 of the cross section.
 - (c) Suppose that the solid D_3 has the same base R , but has equilateral triangles for cross sections perpendicular to the x -axis. Without necessarily calculating the volumes, order the volumes V_1, V_2, V_3 of D_1, D_2, D_3 , respectively, from smallest to largest.
4. Let f and g be continuous, and $0 \leq g(x) \leq f(x)$ on $[a, b]$. The washer method yields the volume formula $V = \int_a^b \pi[(f(x))^2 - (g(x))^2]dx$ when the region between the graphs of f and g are revolved about the x -axis. Now suppose that $g(x) < 0 < f(x)$ for x in $[a, b]$, and $|g(x)| \leq |f(x)|$ for x in $[a, b]$. What should be the resulting formula for V when the bounded region between the graphs of f and g on $[a, b]$ is revolved about the x -axis? Justify your answer.
5. (This problem is related to problem 44 in Section 6.1.) Consider the sphere of radius r that is centered at the origin. Take a vertical slice through the sphere at a distance h to the right of the center.
 - (a) Draw a picture of the situation, and then show that the volume V of the smaller piece cut off from the ellipse (i.e., the piece to the right of $x = h$) satisfies the formula
$$V = \frac{\pi(r-h)^2}{3}(2r+h)$$
 - (b) Suppose that $r = 2$. Approximate the value of h such that $V =$ half the volume of the right hemisphere of the whole sphere, that is, so that $V = 8\pi/3$. (Hint: You likely will need to use the Newton-Raphson method to solve the equation.)

Solution.


1. Denote the radius, height and cost as r, h, f , we have the volume is $\pi r^2 h = 40\pi$, so the cost

$$f = 10 \cdot 2\pi \frac{40}{r} + 25 \cdot 2\pi r^2$$

Solve

$$\frac{df}{dr} = -\frac{800\pi}{r^2} + 100\pi r = 0$$

we get $r = 2$, so $h = 10$. These minimizes the cost.




2.

$$g'(x) = \sin^{2/3}(x + \pi) - \sin^{2/3}(x) = [-\sin(x)]^{2/3} - [\sin(x)]^{2/3} = 0$$


3. (a)

$$A_1(x) = (4 - x^2)^2$$




(b)

$$A_2(x) = \frac{\pi}{2} \left(\frac{(4 - x^2)^2}{2} \right)$$



(c)

$$A_3(x) = \frac{\sqrt{3}(4-x^2)^2}{4}$$



It is not hard to see that

$$A_1(x) > A_3(x) > A_2(x), \quad V_1 > V_3 > V_2$$

The bigger sectional area, the bigger volume.

4. $V = \int_a^b \pi[f(x)]^2 dx$

5. (a)

$$V = \int_h^r \pi(r^2 - x^2) dx = \pi \left(\frac{2r^3}{3} - r^2 h + \frac{h^3}{3} \right)$$

(b)

□

1.2 Worksheet for Section 6.4

1. In calculating the work, W necessary to stretch a spring from its natural length of 8 centimeters to a length of 12 centimeters, we need some more information about the spring. There are two distinct kinds of information, each of which separately would provide the necessary assistance so we could find W . Describe what the two additional kinds of information are.

2. We wish to empty a hemispherical tank full of water out the top. Which should entail the larger amount of work: when the tank is like a cup full of water, or when the tank is upside down with a hole at the top? Explain your answer, and include a picture of the situation.
3. In Example 4 (page 393), a hemispherical tank with radius 10 feet and open at the top is filled with water (Figure 6.43). In determining the work W required to pump all the water to 6 feet above the top of the tank, we assumed that the origin was at the top of the tank, and we derived the formula

$$W = \int_{-10}^0 62.5(6 - x)\pi(100 - x^2)dx$$

Now let us assume that the origin is at the *bottom* of the tank. Find a formula for the radius $r(x)$ of the cross section x units above the origin, and use this formula to help write down the integral for the work W . (If you have time, you should check that the two formulas for W yield the same number of foot-pounds.)

4. A painter weighing 120 pounds is hoisted by means of a rope windlass from the ground to a window on the tenth story of a hotel, 80 feet above ground level.
 - (a) If the weight of the entire rope is considered as negligible, then find the work W required to lift the person the 80 feet above ground level.
 - (b) Assume that instead of a rope, a substantial chain weighing 2 pounds per foot is used in the windlass. Find the work W_1 required to lift the person the 80 feet from ground level. (Hint: Note that as the person is lifted, the length of the chain that must be lifted decreases.)
5. (a) At a party, the punch bowl is hemispherical, with 2 foot radius and open at the top. The fruit punch weighs 66 pounds per cubic foot. At the beginning of the party the bowl is full to the brim. The rules for serving punch require each person to fill the ladle, raise it exactly 1 foot above the top of the bowl (without spilling!), and then pour the punch into a cup at that level. After 10 minutes, 30 people have filled 30 cups, and the punch remaining in the bowl is 4 inches deep. Draw a picture of the situation, and then set up the integral for the work W involved in raising the punch until the punch remaining in the bowl has a depth of only 4 inches. Include all pertinent information in the integral, but do not evaluate the integral.
- (b) Suppose that each person is to raise the ladle exactly 1 foot (*not* 1 foot above the top of the bowl). Set up the integral for the work W_1 involved until the punch is 4 inches deep. Does this activity make any sense for the partygoers? Explain why or why not.

Solution.

1. We need either the spring constant k or the resultant force F .
2. The tank upside down requires more work since more water are concentrated at the bottom
3. The radius is $r(x) = \sqrt{100 - (10 - x)^2}$, and the work needed is

$$W = \int_0^{10} 62.5(16 - x)(100 - (10 - x)^2)dx$$

4. (a) The work need is

$$W = 120 * 80 = 9600 \text{ pound-foot}$$

- (b) The work should be

$$W = 9600 + \int_0^{80} 2(80 - x)dx = 9600 + 6400 = 16000$$

5. (a)

$$W = \int_{-5/3}^0 66(1-x)(4-x^2)dx$$

(b)

$$W_1 = \int_{-5/3}^0 66(4-x^2)dx$$

It doesn't make sense raise the ladle by exactly 1 foot, some punch will stay in the bowl forever.

□

1.3 Worksheet for Section 6.5

1. Let $f(x) = 3\sin x$ for $0 \leq x \leq \pi$, and consider the region R bounded by the x -axis and the graph of f .

(a) Explain in a complete sentence why you don't need to evaluate any integrals in order to determine the x -coordinate \bar{x} of the center of gravity.

(b) Find the center of gravity (\bar{x}, \bar{y}) of R .

2. Let $r > 0$, and let R be the semicircular region bounded below by the x -axis and above by the circle $x^2 + y^2 = r^2$, that is, $x^2 + y^2 = r^2$ with $y \geq 0$.

(a) Find the center of gravity (\bar{x}, \bar{y}) of R .

(b) Find the radius r for which $(\bar{x}, \bar{y}) = (0, \pi)$.

(c) Suppose S is the quarter circular region in the first quadrant, bounded by the x -axis, the y -axis, and the quarter circle $x^2 + y^2 = r^2$. Without evaluating any integrals or making further computations, use symmetry considerations and computations in (a) to determine (\bar{x}, \bar{y}) .

3. Let $a > 0$ and $h > 0$. Consider the triangular region R whose vertices are $(0, 0)$, $(a, 0)$, and $(0, h)$.

(a) Find a formula for the line that represents the hypotenuse of the triangle.

(b) Show that the center of gravity of R is $(a/3, h/3)$. (The center of gravity of R turns out to be precisely the centroid of the triangle, which is by definition the intersection of the 3 medians of the triangle, that is, the lines from the vertices to the midpoints of the opposite sides.)

4. Let f and g be continuous on $[a, b]$, with $a > 0$ and $0 \leq g(x) \leq f(x)$ for $a \leq x \leq b$. Also let R be the region between the graphs of f and g on $[a, b]$, and A the area of R .

(a) Write down a formula for the volume V of the solid obtained by revolving the region R about the y -axis. (Note: If you are in doubt of the formula, please consult Section 6.1.)

(b) Write down a formula for the moment M_y of R about the y -axis.

(c) Use the formulas in (a) and (b) to derive the formula $V = 2\pi\bar{x}A$. (This is the result of Pappus and Guldin in Theorem 6.6, where b is substituted for \bar{x} .)

(d) Suppose someone says that $\bar{x} = 0$. What hypothesis in the problem helps to prevent $\bar{x} = 0$?

5. Find the center of gravity (\bar{x}, \bar{y}) of the region R in Exercise 32(d) in Section 6.5, which region can be split into two rectangles. Then use the formulas at the beginning of Exercise 32 to help find (\bar{x}, \bar{y}) .

Solution.

1. (a) $\bar{x} = \pi/2$ since R is symmetric over $x = \pi/2$.

(b)

$$\bar{y} = \frac{\int_0^\pi \frac{(3 \sin x)^2}{2} dx}{\int_0^\pi 3 \sin x dx} = \frac{3\pi}{16}$$

2. $\bar{x} = 0$ since R is symmetric over $x = 0$, $A = \pi r^2/2$

$$\bar{y} = \frac{1}{A} \int_{-r}^r \frac{r^2 - x^2}{2} dx = \frac{4r}{3\pi}$$

3. (a) $y = -\frac{h}{a}x + h$

(b)

$$M_x = \int_0^a \frac{\left(-\frac{h}{a}x + h\right)^2}{2} dx \stackrel{u = -\frac{h}{a}x + h}{=} -\frac{a}{h} \int_h^0 \frac{u^2}{2} du = \frac{ah^2}{6}, \quad \bar{y} = \frac{h}{3}$$

$$M_y = \int_0^a x \left(-\frac{h}{a}x + h\right) dx = \frac{ha^2}{6}, \quad \bar{x} = \frac{a}{3}$$


4. (a) $V = \int_a^b 2\pi x(f(x) - g(x)) dx$

(b) $M_y = \int_a^b x(f(x) - g(x)) dx$

(c) From (a) and (b) we know that $V = 2\pi M_y = 2\pi \bar{x} A$.

(d) $a > 0$

5. $(\bar{x}, \bar{y}) = (39.5/19, 50.5/19)$



□

1.4 Worksheet for Sections 6.7, 6.2, and 6.8

- Consider the curves C_1 and C_2 parametrized by $x = \cos t, y = \sin t$, and $x = \sin 2t, y = \cos 2t$, respectively, for $0 \leq t \leq 2\pi$. Draw the graphs of C_1 and C_2 , and indicate all values of t in $[0, 2\pi]$, any, for which $C_1(t) = C_2(t)$.
- Under what conditions is it possible to consider a parametrized curve as the graph of a function? (Hint: If $x = f(t)$ and $y = g(t)$ for $a < t < b$, what might one let $f(t)$ be?)
 - Write down parametric equations for the line l defined by $y = ax + b$, where a and b are arbitrary constants with $a \neq 0$ and b .

- (c) We are given a differentiable function f on $[a, b]$, whose graph is C . Derive the formulas for the length L of C from the formula for the length of C when f is given parametrically.
3. Let $f(x) = \sin x$ for $0 \leq x \leq \pi/2$.
- (a) Write down a formula for the length L of the graph of f .
- (b) By using the Comparison Property for Integrals, show that $\pi/2 \leq L \leq \pi$.
4. (a) Let $f(t) = t - \sin t$ for all t . Show that f is a strictly increasing function of t , and determine the inflection points of the graph of f .
- (b) Consider the cycloid C parametrized by $x = t - \sin t$ and $y = 1 - \cos t$, for all real t . Find $C(0)$, $C(\pi)$, and $C(2\pi)$, and show that the highest point on C occurs for $t = \pi$. Tell why this shows that one arch of the cycloid is not a semicircle.
- (c) Let $P(t) = (t - \sin t, 1 - \cos t)$ for all real t . Use (a) to show that if $t_1 \neq t_2$, then $x(t_1) \neq x(t_2)$. (This means that the cycloid C is the graph of a function!)
5. This problem is taken from Project 1 on page 383, which is related to Archimedes' approximation of the circumference of a circle by finding lengths of inscribed (regular) polygons such as those in Figure 6.26. Here we will assume that the point $(1, 0)$ is one vertex of each of the inscribed polygons.
- (a) Consider the inscribed regular polygon of n sides. Show that a vertex adjacent to $(1, 0)$ is $(\cos(2\pi/n), \sin(2\pi/n))$.
- (b) For the inscribed polygon of n sides in (a), show that the length of the side S determined by the 2 points in (a) is $\sqrt{2 - 2\cos(2\pi/n)}$, and show that this expression equals $2\sin(\pi/n)$.
- (c) Using the result of (b), find the length L_n of the inscribed regular polygon of n sides.
- (d) By letting n be large, and using a limit involving $(\sin t)/t$, show why the limit L of L_n as n increases without bound should be 2π .
6. Let $r > 0$. The equations $x = r \cos^3 t$ and $y = r \sin^3 t$ parametrize an astroid (Figure 6.84 in the book). Draw the graph, and find the length L of the astroid. (Hint: Square roots are nonnegative!)

Solution.

1. $C_1(t) = C_2(t)$ reads $\begin{cases} \cos t = \sin 2t \\ \sin t = \cos 2t = 1 - 2\sin^2 t \end{cases}$, solve the second equation we get $\sin t = -1$ or $\sin t = \frac{1}{2}$, so t could be $\frac{3\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}$, all three satisfies the first equation as well.
2. (a)
- (b)
- (c)
3. (a)

$$L = \int_0^{\frac{\pi}{2}} \sqrt{1 + \cos^2 x} dx$$

- (b) Note that $1 \leq \sqrt{1 + \cos^2 x} \leq 2$, so we have

$$\int_0^{\frac{\pi}{2}} 1 dx \leq \int_0^{\frac{\pi}{2}} \sqrt{1 + \cos^2 x} dx \leq \int_0^{\frac{\pi}{2}} 2 dx$$

which is precisely $\pi/2 \leq L \leq \pi$

4. (a) $f'(x) = 1 - \cos t \geq 0$, so f is increasing. $f''(x) = \sin t = 0 \Rightarrow x = k\pi$ for all $k \in \mathbb{Z}$.
- (b) $C(0) = (0, 0)$, $C(\pi) = (\pi, 2)$, $C(2\pi) = (2\pi, 0)$, it cannot be a semicircle since the radius would not be defined!
- (c) Since $x(t) = f(t)$ is strictly increasing.

5. (a) Since it is a regular polygon, we just divide $2n$ into n equal angles, hence the vertex adjacent to $(1, 0)$ will be $(\cos(2\pi/n), \sin(2\pi/n))$.

(b)

$$\sqrt{(1 - \cos(2\pi/n))^2 + \sin^2(2\pi/n)} = \sqrt{2 - 2\cos(2\pi/n)} = 2\sin(\pi/n)$$

(c)

$$L_n = 2n \sin(\pi/n)$$

(d)

$$L = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} 2\pi \frac{\sin(\pi/n)}{\pi/n} = 2\pi$$

6.

$$\begin{aligned} L &= \int_0^{\pi/2} \sqrt{(-3r \cos^2 t \sin t)^2 + (3r \sin^2 t \cos t)^2} dt = \int_0^{\pi/2} \sqrt{9r^2 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t)} dt \\ &= \int_0^{\pi/2} 3r \sin t \cos t dt \stackrel{u=\sin t}{=} \int_0^1 3r u du = \frac{3r}{2} \end{aligned}$$

□

1.5 Sample Exam 1 MATH 141 - Calculus II

Instructions: Please put everything away and no calculators are permitted on this exam. Legibly print your full name on each answer sheet, and write and sign the honor pledge on the first page. Write your solutions neatly on the answer sheets, doing problem 1 on the first page, problem 2 on the second page, and so on. Continue your work onto the back of the page, if necessary. When you have finished, arrange the answer sheets in order and turn them in to your TA.

- (13 points) Let R be the region between the graphs of $f(x) = 1 + \sqrt{x}$ and $g(x) = e^{-x}$ over the interval $[0, 1]$. Find the volume of the solid of revolution obtained by revolving the region R around the x -axis.
- (15 points) Suppose the base of a solid figure S is the triangular region with vertices $(0, 0)$, $(2, 0)$, and $(0, 1)$. Cross-sections perpendicular to the x -axis are semicircles. Calculate the volume of S .
- (15 points) A hemispherical swimming pool has a radius of 9 feet. It is filled with water to a depth of 6 feet. Set up, but DO NOT EVALUATE, the integral for the work required to pump all but 3 feet of water to a platform 1 foot above the top of the pool. Assume the weight of water is 62.5 pounds per cubic foot.
 - (12 points) A spring is expanded 5 inches from its natural length and 20 pounds of force is required to hold it there. Find the work done in stretching the spring an additional 5 inches.
- (15 points) Find the center of mass (centroid) of the region bounded by the curves $y = \sqrt{x}$ and $y = x$.
- (15 points) Find the length of the curve

$$y = \frac{x^3}{6} + \frac{1}{2x}$$


for $1 \leq x \leq 2$.

- (15 points) Calculate the length of the curve described parametrically by $x = \cos^3 t$ and $y = \sin^3 t$ for $0 \leq t \leq \pi/2$.

Solution.

1. First we sketch the situation


1



Note that $0 < g(x) \leq f(x)$. So

$$\begin{aligned} V &= \int_0^1 \pi [f(x)^2 - g(x)^2] dx = \int_0^1 \pi ((1 + \sqrt{x})^2 - e^{-2x}) dx = \int_0^1 \pi (1 + 2\sqrt{x} + x - e^{-2x}) dx \\ &= \left(x + \frac{4}{3}x^{3/2} + \frac{x^2}{2} + \frac{e^{-2x}}{2} \right) \Big|_0^1 = \frac{7}{3} + \frac{e^{-2}}{2} \end{aligned}$$

2. We first sketch the solid



From this we can see that $r(x) = \frac{1 - \frac{1}{2}x}{2}$, and $A(x) = \frac{\pi r(x)^2}{2} = \frac{1}{2} \left(\frac{-\frac{1}{2}x + 1}{2} \right)^2$

$$V = \int_0^2 \frac{1}{2} \left(\frac{-\frac{1}{2}x + 1}{2} \right)^2 dx = \int_0^2 A(x) dx = \frac{1}{12}$$

3. (a)

$$W = \int_{-6}^{-3} 62.5(1-x)\pi(9^2 - x^2)dx$$

(b) We know $20 = 5k \Rightarrow k = 4$, so

$$W = \int_5^{10} kx dx = \int_5^{10} 4x dx = 150$$

4.

$$M_x = \int_0^1 \frac{(\sqrt{x})^2 - x^2}{2} dx = \frac{1}{12}, \quad M_y = \int_0^1 x(\sqrt{x} - x) dx = \frac{1}{15}, \quad A = \int_0^1 (\sqrt{x} - x) dx = \frac{1}{6}$$

$$\bar{x} = \frac{M_y}{A} = \frac{2}{5}, \quad \bar{y} = \frac{M_x}{A} = \frac{1}{2}$$

5. (a)

$$L = \int_1^2 \sqrt{1 + \left(\frac{x^2}{2} - \frac{1}{2x^2}\right)^2} dx = \int_1^2 \sqrt{\left(\frac{x^2}{2} + \frac{1}{2x^2}\right)^2} dx = \int_1^2 \left(\frac{x^2}{2} + \frac{1}{2x^2}\right) dx = \frac{17}{12}$$

(b)

$$L = \int_0^{\frac{\pi}{2}} \sqrt{(-3 \cos^2 t \sin t)^2 + (3 \sin^2 t \cos t)^2} dt = \int_0^{\frac{\pi}{2}} \sqrt{9 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t)} dt$$

$$= \int_0^{\frac{\pi}{2}} 3 \sin t \cos t dt \stackrel{u=\sin t}{=} \int_0^1 3u du = \frac{3}{2}$$

□

1.6 Worksheet for Section 7.1

1. Simplify the following:

a) $(\ln 2 + \ln 3)/\ln 36$

b) $e^{-2 \ln x}$

c) $\ln(e^{-x^2})$

2. Let $f(x) = mx + b$, where $m \neq 0$. Show that f has an inverse, and find a formula for f^{-1} . Finally, explain why f does *not* have an inverse if $m = 0$.

3. (a) Suppose that the formula $y = f(x)$ defines a function f that has an inverse on $[a, b]$. Describe the process you would use in order to find a formula for the inverse f^{-1} .

(b) Let $f(x) = 3/x$. Draw the graph of f , then find a formula for f^{-1} . Does it appear that $f^{-1} = f$? Explain. Do the same for $g(x) = -3/x$.

(c) What property of the graph of a function h implies that $h^{-1} = h$?

4. (a) In a complete sentence, tell why an increasing function f defined on an interval I must have an inverse. Must the inverse be an increasing function? Explain your answer.

(b) Draw the graph of a function f defined on the interval $[0, 1]$ that has an inverse but is *neither* increasing on $[0, 1]$ *nor* decreasing on $[0, 1]$.

(c) Suppose that g is defined and differentiable on $[0, 2]$, with $g(1) = 3$ and $g'(1) = 0$. Assume that g has an inverse. Can $(g^{-1})'(3)$ exist? Explain why or why not.

(d) Suppose that $h(x) = x^5 + x^3 + 2x$. Show that h has an inverse h^{-1} , and that although h is a differentiable function, h^{-1} is *not* a differentiable function.

5. The function $f(x) = x^2$ has no inverse. However, if we restrict the domain to, say, $[0, \infty)$ or $[1, 100)$, then the new function f_1 *does* have an inverse. In the following, find the largest interval containing 0 for which the restricted function f_1 has an inverse:

a) $f(x) = \sin x$

b) $f(x) = \tan x$


c) $f(x) = x^3 - x$

d) $f(x) = x + \sin x$


6. Suppose a ball is thrown vertically (either upward or downward) from a balcony h_0 meters above the ground, with an initial velocity v_0 and constant acceleration due to gravity (-9.8 meters per second squared). Assume further that the ball hits the ground after c seconds.
- Write down a formula for the height h as a function of time t , for $0 \leq t \leq c$. (*Hint:* See p. 323.)
 - Under what physical conditions does h have an inverse h^{-1} ?
 - In the event that h does have an inverse, what does h^{-1} tell us in physical terms?

Solution.


- $\frac{\ln(2 \cdot 3)}{\ln(6^2)} = \frac{\ln 6}{2 \ln 6} = 1/2$.
 - $e^{-2 \ln x} = e^{\ln(x^{-2})} = x^{-2}$.
 - $\ln(e^{-x^2}) = -x^2$.
- $f^{-1}(x) = \frac{x-b}{m}$, f has no inverse since m as the denominator cannot be zero, also because $f(x) = b$ is horizontal line, which evidently fails the horizontal line test.
- First solve x in terms of y , and then change y into x , we get an expression of $f^{-1}(x)$
 - Yes.
 - The graph of h is symmetric over the line $y = x$.
- Because it is one-to-one, the inverse function is also increasing since suppose $b_1 < b_2$, we have unique $a_1 < a_2$ such that $f(a_1) = b_1, f(a_2) = b_2$, i.e. $f^{-1}(b_1) < f^{-1}(b_2)$.
 - We can consider a discontinuous function which increases on $[0, 0.5)$, drops at $x = 0.5$ and decreases on $[0.5, 1]$.




- No. Since otherwise it will be $(g^{-1})'(3) = \frac{1}{g'(1)}$ which is undefined.
 - $h'(x) = 5x^4 + 3x^2 + 2 \geq 2$, hence h is a strictly increasing function.
5. a) $[-\pi/2, \pi/2]$



b) $(-\pi/2, \pi/2)$



c) Solve $f'(x) = 3x^2 - 1 = 0$ we get $x = \pm 1/\sqrt{3}$, so the interval should be $[-1/\sqrt{3}, 1/\sqrt{3}]$



d) Since $f'(x) = 1 + \cos x \geq 0$, f is increasing, so the interval should be $\mathbb{R} = (-\infty, \infty)$.

6. (a) $h(t) = h_0 + v_0 t - 4.9t^2$
 (b) Thrown downwards.
 (c) The time needed to reach certain height.

□

1.7 Worksheet for Section 7.2

- Let $f(x) = e^{(x^2)}$ and $g(x) = (e^x)^2$.
 - Find $f'(x)$ and $g'(x)$.
 - One of the two integrals $\int e^{(x^2)} dx$ and $\int (e^x)^2 dx$ can be evaluated by the methods of Chapter 5. Determine which integral it is, and evaluate the integral.
- Iodine 131, which has been used for treating cancer of the thyroid gland, is also used in order to detect leaks in water pipes. It has a half-life of (approximately) 8.14 days. Suppose that at noon today you have a bottle with 5 grams of Iodine 131.
 - Find a formula for the amount $f(t)$ of Iodine 131 in the bottle t hours after noon today, that is, for each $t \geq 0$. (Hint: Section 4.4 might be helpful.)
 - Show that f has an inverse f^{-1} , and find a formula for f^{-1} .
 - In a complete sentence, indicate what the function f^{-1} tells us physically.
- Let b be any real number. Use the product rule for derivatives (and *not* the Law of Exponents) to show that $\frac{d}{dx}(e^{-x}e^{b+x}) = 0$. Consequently by Theorem 4.6, $e^{-x}e^{b+x}$ is a constant function.

- (b) By using (a) and letting $x = 0$, show that the constant function in (a) is e^b , so that $e^{-x}e^{b+x} = e^b$ for all x .
- (c) Use (b) with $b = 0$ to prove that $e^{-x} = 1/e^x$ for all real x .
- (d) Use (b) and (c) to prove that $e^{b+c} = e^b e^c$ for all real numbers b and c . (You have just proved the Law of Exponents!)
4. Let $f(x) = \frac{1}{\sigma\sqrt{2\pi}}e^{-(x-\mu)^2/2\sigma^2}$, where μ and σ are constants, and $\sigma > 0$.
- (a) Show that $f'(x) = 0$ if $x = \mu$. Then, without taking the second derivative, show that $f(\mu) = \frac{1}{\sigma\sqrt{2\pi}}$ is the maximum value of f .
- (b) Show that $\lim_{x \rightarrow \infty} f(x) = 0$ and $\lim_{x \rightarrow -\infty} f(x) = 0$.
- (c) Let $\sigma = 1$ and $\mu = 0$. Use the left sum formula with $n = 50$ to get an approximate value of $\int_{-3}^3 f(x)dx$. From your answer, guess the value of $\lim_{k \rightarrow \infty} \int_{-k}^k f(x)dx$.

Solution.

1. (a) $f'(x) = 2xe^{(x^2)}$, $g'(x) = 2e^{2x}$
 (b) The second one.
2. (a) We know $f(t) = f(0)e^{kt} = 5e^{kt}$ and

$$2.5 = f(8.14) = 5e^{k8.14} \Rightarrow e^{k8.14} = \frac{1}{2} \Rightarrow k = -\frac{\ln 2}{8.14} \approx -0.085$$

Thus $f(t) = 5e^{-0.085t}$.

- (b) $f'(t) > 0$, $f^{-1}(x) = \frac{1}{0.085} \ln(x/5)$
 (c) The number of days to get decay to certain amount of Iodine.
3. (a)
- $$\frac{d}{dx}(e^{-x}e^{b+x}) = -e^{-x}e^{b+x} + e^{-x}e^{b+x} = 0$$
- (b) $e^{-0}e^{b+0} = e^b$
 (c) Take $b = 0$ in (b) we have $e^{-x}e^x = e^{-x}e^{0+x} = e^0 = 1 \Rightarrow e^{-x} = 1/e^x$.
 (d) Take $x = c$ in (b) we get $e^{-c}e^{b+c} = e^b$, use (c) we know $\frac{1}{e^c}e^{b+c} = e^b \Rightarrow e^{b+c} = e^b e^c$.
4. (a)

$$f'(x) = -\frac{x-\mu}{\sigma^3\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

So $f'(\mu) = 0$, and f attain maximum at $x = \mu$.

- (b)
 (c)

$$\int_{-3}^3 f(x)dx \approx \sum_{k=0}^{49} \frac{1}{50} f\left(-3 + \frac{6k}{50}\right) = \sum_{k=0}^{49} \frac{1}{50} e^{-\frac{(-3 + \frac{6k}{50})^2}{2}}$$

□

1.8 Worksheet for Section 7.3 and 7.5

- (a) Let $f(t) = a(b^t)$, with $f(-1) = 1/2$ and $f(2) = 108$. Find the values of a and b .
 (b) Let $x > 0$. Order the following functions from smallest to largest, and sketch their graphs:

$$3^{-x}, 3^x, 2^{-x}, 2^x, e^x$$

- (a) Let a and b be positive constants, with $a \neq 1$ and $b \neq 1$. Using Theorem 7.8, prove the general change of base formula

$$\log_b x = \log_b a \log_a x, \quad \text{for all } x > 0$$

- (b) We know that $\log_2 7 \approx 2.807355$, $\log_{15} 7 \approx 0.718565$ and $\log_7 15 \approx 1.391663$. Using (a) and whichever such approximations are relevant, approximate $\log_2 15$.
- For certain numerical values of c , the answer when we evaluate the integral $\int \frac{1}{x^2 + 6x + c} dx$ involves an arctangent. By completing the square in the denominator of the integrand, determine those values of c , and evaluate the corresponding integral for an arbitrary such value of c .
- Let g be a differentiable function defined on $[0, 1]$, with $|g(t)| < 3$ for $0 \leq t \leq 1$. Thus $16 - (g(t))^2$ is strictly positive on $[0, 1]$.

- (a) Substitute $u = g(t)$, and then evaluate the integral $\int \frac{g'(t)}{\sqrt{16 - (g(t))^2}} dt$.

- (b) Suppose $g(0) = 0$. Find a value of $g(1)$ so that $\int_0^1 \frac{g'(t)}{\sqrt{16 - (g(t))^2}} dt = \frac{\pi}{3}$.


- Jo drops a marble from her apartment window, 20 meters above the ground. At the same height but 12 meters away, Doe watches the marble fall. When the height of the marble above the ground is h , let θ be the angle between L and M , where L is the horizontal line joining Jo and Doe, and where M represents Doe's line of sight to the marble.

- (a) Draw a picture of the scene, including L , M , and θ . Put the horizontal axis at ground level, and indicate the position of the marble $h(t)$ meters above ground level at time t .
- (b) Find θ in terms of h , and then calculate $d\theta/dh$.
- (c) Assume that $h(t) = 20 - 4.9t^2$ until the marble hits the ground. Find $d\theta/dt$. Show geometrically why $d\theta/dt > 0$, and then tell why the formula for $d\theta/dt$ tells us that $d\theta/dt > 0$.

Note: It turns out that there is a value t_0 for which $d\theta/dt$ has a maximum value. For $\theta(t_0)$ the marble appears to be falling the fastest. Can you find the value of t_0 ?

Solution.

- (a) From $ab^{-1} = f(-1) = 1/2$ we know $a = \frac{b}{2}$, plug it into $ab^2 = f(2) = 108$ we get $b^3 = 216 \Rightarrow b = 6$, so $a = 3$.
 (b) Let's use red to indicate 3^x , blue to indicate 3^{-x} , green to indicate 2^x , yellow to indicate 2^{-x} and black to indicate e^x .



2. (a) The left hand side is $\log_b x = \frac{\ln x}{\ln b}$ which coincide with the right hand side

$$\log_b a \log_a x = \frac{\ln a \ln x}{\ln b \ln a} = \frac{\ln x}{\ln b}$$

- (b) We can simply use (a)

$$\log_2 15 = \log_2 7 \log_7 15 = 2.807355 \cdot 1.391663 \approx 3.90$$

3. First we complete the squares

$$x^2 + 6x + c = (x + 3)^2 + (c - 9)$$

Therefore

$$\int \frac{1}{x^2 + 6x + c} dx = \int \frac{1}{(x + 3)^2 + (c - 9)} dx = \frac{1}{\sqrt{c - 9}} \arctan \left(\frac{x + 3}{\sqrt{c - 9}} \right) + C$$

So $c > 9$.

4. (a)


$$\int \frac{g'(t)}{\sqrt{16 - (g(t))^2}} dt \stackrel{u=g(t)}{=} \int \frac{du}{\sqrt{16 - u^2}} = \arcsin \left(\frac{g(t)}{4} \right) + C$$

- (b) By Fundamental Theore of Calculus, we have

$$\begin{aligned} \int_0^1 \frac{g'(t)}{\sqrt{16 - (g(t))^2}} dt &= \arcsin \left(\frac{g(t)}{4} \right) \Big|_0^1 = \arcsin \left(\frac{g(1)}{4} \right) - \arcsin \left(\frac{g(0)}{4} \right) \\ &= \arcsin \left(\frac{g(1)}{4} \right) - \arcsin \left(\frac{0}{4} \right) \\ &= \arcsin \left(\frac{g(1)}{4} \right) = \frac{\pi}{3} \end{aligned}$$

$$\text{So } \frac{g(1)}{4} = \sin\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{2} \Rightarrow g(1) = 2\sqrt{3}.$$

5. The picture looks like



- (a) $h(t) = 20 - 4.9t^2$

- (b) By examining the picture we see that $\tan \theta = \frac{20 - h}{12}$, so $\theta = \arctan \left(\frac{20 - h}{12} \right)$, so

$$\frac{d\theta}{dh} = \frac{1}{\left(\frac{20-h}{12} \right)^2 + 1} \cdot \left(-\frac{1}{12} \right)$$

(c)

$$\frac{d\theta}{dt} = \frac{d\theta}{dh} \frac{dh}{dt} = \frac{1}{\left(\frac{20-h}{12}\right)^2 + 1} \cdot \left(-\frac{1}{12}\right) (-9.8t) = \frac{1}{\left(\frac{20-h}{12}\right)^2 + 1} \cdot \left(\frac{9.8t}{12}\right) > 0$$

There for $\frac{\theta}{dt} > 0$, and it achieves its maximum when it hits the ground, i.e. when $h(t_0) = 0 \Rightarrow t_0 = \sqrt{\frac{20}{4.9}}$.

□

1.9 Worksheet for Section 7.6

- Show that 1^∞ is an indeterminate form by finding 2 values of c for which $\lim_{x \rightarrow \infty} \left(1 + \frac{c}{x}\right)^x$ yields two different numbers.
 - Show that 0^∞ is a determinate form by looking at $\lim_{x \rightarrow \infty} f(x)^{g(x)}$, where $\lim_{x \rightarrow \infty} f(x) = 0$ and $\lim_{x \rightarrow \infty} g(x) = \infty$, and telling the behavior of the values of $f(x)^{g(x)}$ as x approaches ∞ .
- Exercise 51 in the textbook asks us to evaluate one of the limits that appeared in l'Hôpital's original text on calculus: $\lim_{x \rightarrow a} \frac{a^2 - ax}{a - \sqrt{ax}}$ with $a > 0$.
 - Use l'Hôpital's Rule to evaluate the limit.
 - Rewrite the numerator as the product of two expressions, one of which is the denominator. Then simplify the resulting expression, and evaluate the limit *without* using l'Hôpital's Rule.
- Consider $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}}$.
 - Apply l'Hôpital's Rule twice. What do you observe?
 - Factor out x^2 from $x^2 + 1$ in the denominator, then cancel, and next evaluate the limit without l'Hôpital's Rule.
 - Corroborate your answer to part (b) by plotting. and examining the portion of the graph of $y = x/\sqrt{x^2 + 1}$ for large values of x .
- Consider $\lim_{x \rightarrow 0} \frac{\sin x}{x + 1}$.
 - Use the l'Hôpital's Rule process to obtain value of the limit.
 - Use the limit rules and *not* l'Hôpital's Rule in order to obtain the value of the Limit.
 - In a sentence, tell why the result of (a) is wrong.
- Suppose a right circular cylinder with top and bottom is to be made in such a way that if the radius is r , then the height of the cylinder is 2^{-r} . Determine what happens to the volume V of the cylinder as r increases without bound.
- EXTRA: Solve Exercise 60 in Section 7.6, which is the following: Show that $\lim_{x \rightarrow 0^+} x^{(x^x)} = 0$.

Solution.

1. (a)

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e, \quad \lim_{x \rightarrow \infty} \left(1 + \frac{-1}{x}\right)^x = 1/e, \quad \lim_{x \rightarrow \infty} \left(1 + \frac{0}{x}\right)^x = 1 = e^0$$

In general, first we know

$$\lim_{x \rightarrow \infty} x \ln \left(1 + \frac{c}{x}\right) = \lim_{x \rightarrow \infty} \frac{\ln \left(1 + \frac{c}{x}\right)}{1/x} \stackrel{\text{l'H } \frac{0}{0}}{=} \lim_{x \rightarrow \infty} \frac{\frac{1}{1+\frac{c}{x}} \left(-\frac{1}{x^2}\right)}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{c}{x}} = c$$

so we have

$$\lim_{x \rightarrow \infty} \left(1 + \frac{c}{x}\right)^x = \lim_{x \rightarrow \infty} e^{\ln \left(1 + \frac{c}{x}\right)^x} = \lim_{x \rightarrow \infty} e^{x \ln \left(1 + \frac{c}{x}\right)} = e^{\lim_{x \rightarrow \infty} x \ln \left(1 + \frac{c}{x}\right)} = e^c$$

(b)

2. (a)

$$\lim_{x \rightarrow a} \frac{a^2 - ax}{a - \sqrt{ax}} \stackrel{\text{FH } \frac{0}{0}}{=} \lim_{x \rightarrow a} \frac{-a}{-\frac{\sqrt{a}}{2\sqrt{x}}} = \frac{-a}{-\frac{1}{2}} = 2a$$

(b)

$$\lim_{x \rightarrow a} \frac{a^2 - ax}{a - \sqrt{ax}} = \lim_{x \rightarrow a} \frac{(a - \sqrt{ax})(a + \sqrt{ax})}{a - \sqrt{ax}} = \lim_{x \rightarrow a} (a + \sqrt{ax}) = 2a$$

3. (a)


$$\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} = \lim_{x \rightarrow \infty} \frac{1}{\frac{x}{\sqrt{x^2 + 1}}} = \lim_{x \rightarrow \infty} \frac{\sqrt{x^2 + 1}}{x} = \lim_{x \rightarrow \infty} \frac{\frac{x}{\sqrt{x^2 + 1}}}{1} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}}$$

This has no progress.

(b)

$$\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} = \lim_{x \rightarrow \infty} \frac{x}{x\sqrt{1 + x^{-2}}} = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + x^{-2}}} = \frac{1}{1 + 0} = 1$$

(c)



4. (a)

$$\lim_{x \rightarrow 0} \frac{\sin x}{x + 1} = \lim_{x \rightarrow 0} \frac{\cos x}{1} = 1$$

(b)

$$\lim_{x \rightarrow 0} \frac{\sin x}{x + 1} = \frac{0}{0 + 1} = 0$$

(c) $\frac{0}{a}$ where $a \neq 0$ is not an indeterminate form.

5. First we know $V = \pi r^2 \cdot 2^{-r} = \frac{\pi r^2}{2^r}$, so

$$\lim_{r \rightarrow \infty} V = \lim_{r \rightarrow \infty} \frac{\pi r^2}{2^r} = \lim_{r \rightarrow \infty} \frac{2\pi r}{2^r \ln 2} = \lim_{r \rightarrow \infty} \frac{2\pi}{2^r (\ln 2)^2} = 0$$

6. First consider

$$\lim_{x \rightarrow 0^+} x \ln x = \lim_{x \rightarrow 0^+} \frac{\ln x}{1/x} = \lim_{x \rightarrow 0^+} \frac{1/x}{-1/x^2} = - \lim_{x \rightarrow 0^+} x = 0$$

So we know

$$\lim_{x \rightarrow 0^+} x^x = \lim_{x \rightarrow 0^+} e^{x \ln x} = e^0 = 1$$

And

$$\lim_{x \rightarrow 0^+} x^{(x^x)} = \lim_{x \rightarrow 0^+} e^{(x^x) \ln x} = e^{-\infty} = 0$$

□

1.10 Worksheet for Section 7.6 and 8.1

1. Determine which of the following limits exist as a number, which as ∞ , and which don't exist at all. For any limit that exists as a number, determine the number.

$$\begin{array}{lll} \text{(a)} & \text{(i)} \lim_{x \rightarrow 0^+} x^x & \text{(ii)} \lim_{x \rightarrow 0^+} (1/x)^x & \text{(iii)} \lim_{x \rightarrow \infty} x^{\sin x} \\ \text{(b)} & \text{(i)} \lim_{x \rightarrow 0^+} (1 - 1/x)^x & \text{(ii)} \lim_{x \rightarrow 0^+} (1 - 1/x)^{1/x} & \text{(iii)} \lim_{x \rightarrow \infty} (1 - 1/x)^x \end{array}$$

2. In the following, one integral in each pair can be evaluated by substitution and the other integral can be evaluated by integration by parts. In each case, write down which method applies, and evaluate the integral.

$$\begin{array}{ll} \text{(a)} & \text{(i)} \int x \cos(x^2) dx & \text{(ii)} \int x^2 \cos x dx \\ \text{(b)} & \text{(i)} \int \frac{x^2}{e^x} dx & \text{(ii)} \int x e^{(x^2)} dx \\ \text{(c)} & \text{(i)} \int \frac{1}{x^4} \ln x dx & \text{(ii)} \int \frac{1}{x} (\ln x)^4 dx \end{array}$$

3. Consider the integral $\int e^{\pi x} \sin(2x) dx$.

- (a) In half of your group, solve the integral by performing integration by parts (twice!) with $u = \sin(2x)$ and $du = e^{\pi x} dx$. In the other half of the group, solve the integral by performing integration by parts (twice!) with $u = e^{\pi x}$ and $dv = \sin(2x) dx$. Are the answers equivalent to each other? Explain in a sentence.
- (b) Which of integration by parts processes is easier, or are they both about the same? Explain your answer. (Note: Sometimes it is not so simple to determine which part of the integrand is better as u , and which is better as dv .)
4. In each of the following, try to use the technique of integration by parts with the given u . In each, explain why that choice of u is not effective. Then write down a different function u and also write down dv that will make integration by parts effective, and tell why you chose your u and dv . (Don't perform the integration, however.)

$$\text{(i)} \int e^{-3x} x^{10} dx; u = e^{-3x} \qquad \text{(ii)} \int x^4 \sin x dx; u = \sin x$$

Solution.

1. (a) (i)

$$\lim_{x \rightarrow 0^+} x^x = \lim_{x \rightarrow 0^+} e^{x \ln x} = e^0 = 1$$

- (ii)

$$\lim_{x \rightarrow 0^+} (1/x)^x = \lim_{x \rightarrow 0^+} 1/x^x = 1$$

- (iii) It doesn't exist since $\sin x$ oscillates between -1 and 1 , $x^{\sin x}$ oscillates between really small and really big, so it doesn't have a limit.

- (b) (i) $(1 - 1/x)^x = e^{x \ln(1-1/x)}$, but $\lim_{x \rightarrow 0^+} \ln(1 - 1/x)$ is not well defined.

- (ii) $(1 - 1/x)^{1/x} = e^{\ln(1-1/x)/x}$, but $\lim_{x \rightarrow 0^+} \ln(1 - 1/x)$ is not well defined.

- (iii) As in 1(a) of worksheet 7.6, we get e^{-1} .

2. (i)

$$\int x \cos(x^2) dx \stackrel{u=x^2}{=} \frac{1}{2} \int \cos u du = \frac{1}{2} \sin u + C = \frac{1}{2} \sin(x^2) + C$$

(ii)

$$\begin{aligned}
& \int x^2 \cos x dx \stackrel{u=x^2, dv=\cos x dx=d \sin x}{=} x^2 \sin x - \int 2x \sin x dx \\
& \stackrel{u=2x, dv=\sin x dx=d(-\cos x)}{=} x^2 \sin x + 2x \cos x - \int 2 \cos x dx \\
& = x^2 \sin x + 2x \cos x - 2 \sin x + C
\end{aligned}$$

(iii) (i)

$$\begin{aligned}
& \int \frac{x^2}{e^x} dx \stackrel{u=x^2, dv=e^{-x} dx=d(-e^{-x})}{=} -x^2 e^{-x} + \int 2x e^{-x} dx \\
& \stackrel{u=2x, dv=e^{-x} dx=d(-e^{-x})}{=} -x^2 e^{-x} - 2x e^{-x} + \int 2 e^{-x} dx \\
& = -x^2 e^{-x} - 2x e^{-x} - 2 e^{-x} + C
\end{aligned}$$

(ii)

$$\int x e^{(x^2)} dx \stackrel{u=x^2}{=} \frac{1}{2} \int e^u du = \frac{1}{2} e^u + C = \frac{1}{2} e^{(x^2)} + C$$

(iv) (i)

$$\begin{aligned}
& \int \frac{1}{x^4} \ln x dx \stackrel{u=\ln x, dv=x^{-4} dx=d(-1/3x^{-3})}{=} -\frac{1}{3x^3} \ln x + \int \frac{1}{3x^3} \frac{1}{x} dx \\
& = -\frac{1}{3x^3} \ln x - \frac{1}{9x^3} + C
\end{aligned}$$

(ii)

$$\int \frac{1}{x} (\ln x)^4 dx \stackrel{u=\ln x}{=} \int u^4 du = \frac{1}{5} u^5 + C = \frac{1}{5} (\ln x)^5 + C$$

3. (a) Assume $I = \int e^{\pi x} \sin(2x) dx$.

$$\begin{aligned}
I & \stackrel{u=\sin(2x), dv=e^{\pi x} dx=d(e^{\pi x}/\pi)}{=} \frac{e^{\pi x}}{\pi} \sin(2x) - \frac{2}{\pi} \int e^{\pi x} \cos(2x) dx \\
& \stackrel{u=\cos(2x), dv=e^{\pi x} dx=d(e^{\pi x}/\pi)}{=} \frac{e^{\pi x}}{\pi} \sin(2x) - \frac{2e^{\pi x} \cos(2x)}{\pi^2} - \frac{4}{\pi^2} I
\end{aligned} \tag{1.10.1}$$

$$\begin{aligned}
I & \stackrel{u=e^{\pi x}, dv=\sin(2x) dx=d(-\cos(2x)/2)}{=} -\frac{\cos(2x)}{2} e^{\pi x} + \frac{\pi}{2} \int e^{\pi x} \cos(2x) dx \\
& \stackrel{u=e^{\pi x}, dv=\cos(2x) dx=d(\sin(2x)/2)}{=} -\frac{\cos(2x)}{2} e^{\pi x} + \frac{\pi}{4} e^{\pi x} \sin(2x) - \frac{\pi^2}{4} I
\end{aligned} \tag{1.10.2}$$

Both equations give us

$$I = \frac{e^{\pi x} (\pi \sin(2x) - 2 \cos(2x))}{4 + \pi^2}$$

You will get the same answer since they the same integral after all!

(b) It is about the same difficulty.

4. (a) It is ineffective since it raised the power of x . If we choose $u = x^{10}$, then $du = 10x^9 dx$ which reduces down the power on x .
- (b) It is ineffective since it raised the power of x . If we choose $u = x^4$, then $du = 4x^3 dx$ which reduces down the power on x .

□

1.11 Sample Exam 2 MATH 141 - Calculus II

Instructions: Please put everything away and no calculators are permitted on this exam. Legibly print your full name on each answer sheet, and write and sign the honor pledge on the first page. Write your solutions neatly on the answer sheets, doing problem 1 on the first page, problem 2 on the second page, and so on. Continue your work onto the back of the page, if necessary. When you have finished, arrange the answer sheets in order and turn them in to your TA.

1. Let $f(x) = 3x + \sqrt{x+10}$.

(a) (10 points) Find the largest interval on which f has an inverse function f^{-1} . What are the domain and range of f^{-1} ?

(b) (10 points) Calculate $(f^{-1})'(0)$.

2. Find the derivative y' . You do not have to simplify your answer.

(a) (10 points)

$$y = 3 \log_{10}(2x) + 2^{x-1} + \sin^{-1}(x^3)$$

(b) (10 points)

$$y = (\sin^{-1}(x^2))^{\tan^{-1} x}$$

3. (a) (10 points) Let $f(x) = \sin\left(\cos^{-1}\left(\frac{5x}{7}\right)\right)$. Rewrite the function f so it does not contain a trigonometric or inverse trigonometric function.

(b) (10 points) Evaluate the integral

$$\int \frac{1}{x^2 + 3x + 3} dx$$

4. Find the limit.

(a) (10 points)

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{(1 - e^x)^2}$$

(b) (10 points)

$$\lim_{x \rightarrow \infty} \left(\frac{x}{x+1} \right)^{2x}$$

5. Evaluate the integral.

(a) (10 points)

$$\int_0^{\pi/4} x^2 \sin 2x dx$$

(b) (10 points)

$$\int_0^{\pi/2} e^{2x} \cos x dx$$

Solution.

1. (a) Since $f'(x) = 3 + \frac{1}{2\sqrt{x+10}} \geq 3$, f is always strictly increases, so f has an inverse on $[-10, \infty)$. The domain of f^{-1} is the range of f which is $[-30, \infty)$, and range of f^{-1} is the domain of f which is $[-10, \infty)$.

(b) First solve $0 = f(x) = 3x + \sqrt{x+10} \iff 3x = -\sqrt{x+10}$, so we have $9x^2 = x+10 \iff (9x-10)(x+1) = 0 \iff x = -1$ or $x = 10/9$ which we discard since it is not a solution (we can also see this because $x = -\sqrt{x+10}/3$ should be non-positive)

$$(f^{-1})'(0) = \frac{1}{f'(-1)} = \frac{1}{3 + \frac{1}{2 \cdot 3}} = \frac{6}{19}$$

2. (a) First we can change the base for log

$$y = \frac{3 \ln(2x)}{\ln 10} + e^{(x-1) \ln 2} + \sin^{-1}(x^3)$$

Therefore


$$y' = \frac{3}{(\ln 10)x} + 2^{x-1} \ln 2 + 3x^2 \sqrt{1+x^6}$$

- (b) First we rewrite it as

$$y = e^{\tan^{-1} x \ln(\sin^{-1}(x^2))}$$

$$y' = (\sin^{-1}(x^2))^{\tan^{-1} x} \left[\frac{\ln(\sin^{-1}(x^2))}{1+x^2} + \tan^{-1} x \frac{1}{\sin^{-1}(x^2)} \frac{1}{1+x^4} 2x \right]$$

3. (a) Suppose $\theta = \cos^{-1}\left(\frac{5x}{7}\right)$, then $\cos \theta = \frac{5x}{7}$, so $\sin \theta = \frac{\sqrt{49-25x^2}}{7}$



- (b) First complete the squares $x^2 + 3x + 3 = (x + \frac{3}{2})^2 + \frac{3}{4}$, then

$$\int \frac{dx}{x^2 + 3x + 3} = \int \frac{dx}{(x + \frac{3}{2})^2 + \frac{3}{4}} = \frac{2}{\sqrt{3}} \arctan\left(\frac{2x+3}{\sqrt{3}}\right) + C$$

4. (a)

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{(1 - e^x)^2} \stackrel{\text{L'H } \frac{0}{0}}{=} \lim_{x \rightarrow 0} \frac{\sin x}{-2e^x(1 - e^x)} = \lim_{x \rightarrow 0} \frac{1}{2e^x} \lim_{x \rightarrow 0} \frac{\sin x}{-(1 - e^x)} = \frac{1}{2} \lim_{x \rightarrow 0} \frac{\sin x}{-(1 - e^x)}$$

$$\stackrel{\text{L'H } \frac{0}{0}}{=} \frac{1}{2} \lim_{x \rightarrow 0} \frac{\cos x}{e^x} = \frac{1}{2}$$

- (b) We can covert this into an expression that we have dealt before

$$\lim_{x \rightarrow \infty} \left(\frac{x}{x+1}\right)^{2x} = \lim_{x \rightarrow \infty} \left(\frac{x+1}{x}\right)^{-2x} = \lim_{x \rightarrow \infty} \left(\left(x + \frac{1}{x}\right)^x\right)^{-2} = e^{-2}$$

5. (a)

$$\int_0^{\pi/4} x^2 \sin 2x dx \stackrel{u=x^2, dv=\sin(2x)dx=d(-\cos(2x)/2)}{=} -\frac{1}{2} x^2 \cos(2x) \Big|_0^{\pi/4} + \int_0^{\pi/4} x \cos(2x) dx$$

$$\stackrel{u=x, dv=\cos(2x)dx=d(\sin(2x)/2)}{=} \frac{1}{2} x \sin(2x) \Big|_0^{\pi/4} - \frac{1}{2} \int_0^{\pi/4} \sin(2x) dx$$

$$= \frac{\pi}{8} - \frac{1}{4} \cos(2x) \Big|_0^{\pi/4} = \frac{\pi}{8} - \frac{1}{4}$$

- (b) Suppose $I = \int_0^{\pi/2} e^{2x} \cos x dx$

$$I \stackrel{u=e^{2x}, dv=\cos x dx=d(\sin x)}{=} e^{2x} \sin x \Big|_0^{\pi/2} - 2 \int_0^{\pi/2} e^{2x} \sin x dx$$

$$\stackrel{u=e^{2x}, dv=\sin x dx=d(-\cos x)}{=} e^{\pi} + 2e^{2x} \cos x \Big|_0^{\pi/2} - 4I$$

$$\text{Solve the equal we get } I = \frac{e^{\pi} - 2}{5}$$

□

1.12 Worksheet for Section 8.2

- (Warmup): Let $f(x) = \sin x$ for $-\pi/2 \leq x \leq 0$, and R the region between the graph of f and the x -axis on $[-\pi/2, 0]$. Sketch R , and find the volume V of the solid obtained by revolving R about the x -axis.
- Consider the two integrals $\int \sin^8 x \cos^6 x dx$ and $\int \sin^7 x \cos^4 x dx$. Perform the integration for the integral that is easier to integrate.
 - Evaluate each of the following three integrals:
$$\int \tan^6 x \sec^8 x dx, \quad \int \tan^3 x \sec^4 x dx, \quad \int \tan^5 x \sec^3 x dx$$
- Consider the integral $\int \sin^3 x dx$.
 - Evaluate the integral by an appropriate reduction formula in Section 8.1, and write the constant added at the end as C_1 .
 - Evaluate the integral by the method of Section 8.2 (i.e., by factoring out $\sin x$ and by writing the rest of the integrand in terms of $\cos x$), and write the constant added at the end as C_2 .
 - Show that the results found in (a) and (b) are compatible, by finding a relationship between C_1 and C_2 . (*Hint*: Eliminate $\sin x$ from the answers in (a) and/or (b).)
- Let r be any positive number (rational or irrational), and consider $\int_0^{\pi/2} \sin^r x \cos^3 x dx$.
 - Without evaluating the integral, show that $0 < \int_0^{\pi/2} \sin^r x \cos^3 x dx < 2$.
 - Evaluate the given integral, and find the smallest positive value of s such that the value of the integral is always less than s , irrespective of the positive value of r .
- In a short paragraph, describe 3 different types of functions you have learned how to integrate in Section 8.2, and the method employed in integrating each type.

Solution.

1.

$$V = \int_{-\pi/2}^0 \pi \sin^2 x dx = \int_{-\pi/2}^0 \frac{\pi(1 - \cos(2x))}{2} dx = \pi \left(\frac{x}{2} - \frac{\sin(2x)}{4} \right) \Big|_{-\pi/2}^0 = \frac{\pi^2}{4}$$

2. (a)

$$\begin{aligned} \int \sin^7 x \cos^4 x dx &\stackrel{u=\cos x}{=} - \int (1 - u^2)^3 u^4 du = - \int (1 - u^2)^3 u^4 du \\ &= - \int u^4 - 3u^6 + 3u^8 - u^{10} du = - \frac{\cos^5 x}{5} + \frac{3 \cos^7 x}{7} - \frac{\cos^9 x}{3} + \frac{\cos^{11} x}{11} + C \end{aligned}$$

(b)

$$\begin{aligned} \int \tan^6 x \sec^8 x dx &\stackrel{u=\tan x}{=} \int u^6 (1 + u^2)^3 du = \int u^6 + 3u^8 + 3u^{10} + u^{12} du \\ &= \frac{\tan^7 x}{7} + \frac{\tan^9 x}{3} + \frac{3 \tan^{11} x}{11} + \frac{\tan^{13} x}{13} + C \end{aligned}$$

$$\int \tan^3 x \sec^4 x dx \stackrel{u=\sec x}{=} \int (u^2 - 1)u^3 du = \int u^5 - u^3 du = \frac{\sec^6 x}{6} - \frac{\sec^4 x}{4} + C$$

$$\begin{aligned}\int \tan^5 x \sec^3 x dx &\stackrel{u=\sec x}{=} \int (u^2 - 1)^2 u^2 du = \int u^6 - 2u^4 + u^2 du \\ &= \frac{\sec^7 x}{7} - \frac{2\sec^5 x}{5} + \frac{\sec^3 x}{3} + C\end{aligned}$$

3. (a) Recall the reduction formula

$$\int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx$$

Thus

$$\int \sin^3 x dx = -\frac{1}{3} \sin^2 x \cos x + \frac{2}{3} \int \sin x dx = -\frac{1}{3} \sin^2 x \cos x - \frac{2}{3} \cos x + C_1$$

- (b)

$$\int \sin^3 x dx \stackrel{u=\cos x}{=} -\int (1-u^2) du = -u + \frac{u^3}{3} + C_2 = \frac{\cos^3 x}{3} - \cos x + C_2$$

- (c) If we convert $\sin x$ into $\cos x$ in (a), it becomes

$$\int \sin^3 x dx = -\frac{1}{3} (1 - \cos^2 x) \cos x - \frac{2}{3} \cos x + C_1 = \frac{1}{3} \cos^3 x - \cos x + C_1$$

So we see that $C_1 = C_2$

4. (a) On $[0, \pi/2]$, $0 \leq \sin x \leq 1$, $0 \leq \cos x \leq 1$, and therefore $0 \leq \sin^r x \cos^3 x \leq 1$, by comparison property, we have

$$0 = \int_0^{\pi/2} 0 dx < \int_0^{\pi/2} \sin^r x \cos^3 x dx \leq \int_0^{\pi/2} 1 dx = \frac{\pi}{2} < 2$$

- (b)

$$\int_0^{\pi/2} \sin^r x \cos^3 x dx \stackrel{u=\sin x}{=} \int_0^1 u^r (1-u^2) du = \left. \frac{u^{r+1}}{r+1} - \frac{u^{r+3}}{r+3} \right|_0^1 = \frac{2}{(r+1)(r+3)}$$

It is easy to see that $I(r) = \int_0^{\pi/2} \sin^r x \cos^3 x dx$ as a function of r is decreasing, so it attains its minimum $2/3$ at $r = 0$.

- 5.

□

1.13 Worksheet for Section 8.3

1. (Warmup):

- (a) Let $g(t) = \frac{t^2}{\sqrt{4-t^2}}$, for $-2 < t < 2$. Find the largest interval containing $t = 1$ for which g^{-1} exists.

- (b) Find $\lim_{x \rightarrow \infty} (x^2 5^{-x})$, showing your work.

- (c) Evaluate $\int \sin^5 t \cos^6 t dt$.

2. In a sentence, describe the difference between integrating by ordinary substitution (as described in Section 5.6), and integrating by trigonometric substitution.
3. A trigonometric substitution is appropriate for only one of the following integrals, and ordinary substitution (as in Section 5.6) is appropriate for the other two. Evaluate each integral by using an appropriate method, and specify which method you use.

$$(a) \int x\sqrt{x^2-9}dx \qquad (b) \int \frac{1}{x\sqrt{x^2-9}}dx \qquad (c) \int \frac{x}{\sqrt{x^2-9}}dx$$

4. (a) Use integration by parts and trigonometric substitution in order to find $\int x \sin^{-1} x dx$.

(b) Find $\int x \tan^{-1} x dx$.

5. Find the area A of the region R between the graph of $f(x) = \frac{x^2}{\sqrt{4-x^2}}$ and the x -axis on the interval $[-1, 1]$.

6. (a) Use trigonometric substitution to show that if $a > 0$ and $b > 0$, then the region enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ has area $A = \pi ab$

(b) If $a = b = 1$ for part (a), and you want to find the area of the circular region (which is evidently π) by integration, do you need to use trigonometric substitution? Explain why, or why not.

Solution.

1. (a) First realize that $g(t) = 1 - \frac{4}{\sqrt{4-t^2}}$

$$g'(t) = \frac{t(8-t^2)}{(\sqrt{4-t^2})^3}$$

Which means that g decreases in $(-2, 0]$ and increases in $[0, 2)$, therefore, $[0, 2)$ is the largest interval containing $t = 1$ where g^{-1} exists.

(b)

$$\lim_{x \rightarrow \infty} (x^2 5^{-x}) = \lim_{x \rightarrow \infty} \frac{x^2}{5^x} \stackrel{\text{L'H } \infty}{=} \frac{2x}{(\ln 5)5^x} \stackrel{\text{L'H } \infty}{=} \frac{2}{(\ln 5)^2 5^x} = 0$$

(c)

$$\begin{aligned} \int \sin^5 t \cos^6 t dt &\stackrel{u=\cos t}{=} - \int (1-u^2)^2 u^6 du = \int -u^6 + 2u^8 - u^{10} du \\ &= -\frac{\cos^7 t}{7} + \frac{2 \cos^9 t}{9} - \frac{\cos^{11} t}{11} + C \end{aligned}$$

2. Ordinary u -substitution you substitute u for a function of x , for trigonometry substitution you substitute x for a trigonometric function.

3. (a)

$$\int x\sqrt{x^2-9}dx \stackrel{u=x^2-9}{=} \frac{1}{2} \int \sqrt{u} du = \frac{1}{3}(x^2-9)^{3/2} + C$$

(b)

$$\int \frac{1}{x\sqrt{x^2-9}}dx \stackrel{x=3\sec t}{=} \int \frac{1}{9 \sec t \tan t} 3 \sec t \tan t dt = \int \frac{1}{3} dt = \frac{t}{3} + C = \frac{\operatorname{arcsec}(x/3)}{3} + C$$

(c)

$$\int \frac{x}{\sqrt{x^2-9}}dx \stackrel{u=x^2-9}{=} \frac{1}{2} \int u^{-1/2} du = \sqrt{x^2-9} + C$$

4. (a)

$$\begin{aligned}
 \int x \sin^{-1} x dx &\stackrel{u=\sin^{-1} x, dv=xdx}{=} \frac{1}{2} x^2 \sin^{-1} x - \frac{1}{2} \int \frac{x^2}{\sqrt{1-x^2}} dx \stackrel{x=\sin t}{=} \\
 &\frac{1}{2} t \sin^2 t - \frac{1}{2} \int \frac{\sin^2 t}{\cos t} \cos t dt = \frac{1}{2} t \sin^2 t - \frac{t}{4} + \frac{\sin(2t)}{8} + C \\
 &= \frac{1}{2} t \sin^2 t - \frac{t}{4} + \frac{2 \sin t \cos t}{8} + C \\
 &= \frac{x^2 \sin^{-1} x}{2} - \frac{\sin^{-1} x}{4} + \frac{x \sqrt{1-x^2}}{4} + C
 \end{aligned}$$

(b)

$$\int x \tan^{-1} x dx \stackrel{u=\tan^{-1} x, dv=xdx}{=} \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{x^2+1} dx$$

We can do this via

$$\begin{aligned}
 &\stackrel{x=\tan t}{=} \frac{x^2 \arctan x}{2} - \frac{1}{2} \int \frac{\tan^2 t}{\sec^2 t} \sec^2 t dt = \frac{x^2 \arctan x}{2} - \frac{1}{2} \int \tan^2 t dt \\
 &= \frac{x^2 \arctan x}{2} - \frac{1}{2} \int \sec^2 t - 1 dt = \frac{x^2 \arctan x}{2} - \frac{\tan t - t}{2} + C \\
 &= \frac{x^2 \arctan x}{2} - \frac{x - \arctan x}{2} + C
 \end{aligned}$$

Or

$$= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int 1 - \frac{1}{x^2+1} dx = \frac{1}{2} x^2 \tan^{-1} x - \frac{x - \arctan x}{2} + C$$

5.

$$\begin{aligned}
 \int \frac{x^2}{\sqrt{4-x^2}} dx &\stackrel{x=2 \sin t}{=} \int \frac{4 \sin^2 t}{2 \cos t} 2 \cos t dt = \int 4 \sin^2 t dt \\
 &= 2 \int 1 - \cos(2t) dt = 2t - \sin(2t) + C = 2t - 2 \sin t \cos t + C \\
 &= 2 \arcsin(x/2) + x \sqrt{1 - \frac{x^2}{4}} + C
 \end{aligned}$$

So

$$A = \int_{-1}^1 \frac{x^2}{\sqrt{4-x^2}} dx = \left(2 \arcsin(x/2) - x \sqrt{1 - \frac{x^2}{4}} \right) \Big|_{-1}^1 = \frac{2\pi}{3} - \sqrt{3}$$

6. (a)

$$A = 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \stackrel{x=a \sin t}{=} 4 \int_0^{\pi/2} ab \cos^2 t dt = 4ab$$

(b) Yes.

□

1.14 Worksheet for Section 8.4 and 8.6

- Where applicable in (a) - (c), carry out the necessary division (as in Example 1 on p. 530) in preparation for integration. If division is not needed, give a reason why it is not needed.

$$(a) \frac{x^2}{x^2 - x + 1} \qquad (b) \frac{x - 1}{x^2 - x + 1} \qquad (c) \frac{x^4 + 4}{x^2 + 1}$$

2. In the following, write the rational function as a combination of factors like those appearing in (4) and/or (5) of Section 8.4, without finding the numerical values of the constants $A_1, \dots, B_1, \dots, C_1, \dots$

$$(a) \frac{6x^2 - 4x + 1}{(x - 2)^2} \qquad (b) \frac{x^3 + 2x^2 + 3x + 4}{x^4 - 16}$$

3. (a) Show that $\frac{x^2}{(x^2 - 4)^2} = \frac{1}{4} \left(\frac{1}{x - 2} + \frac{1}{x + 2} \right)^2$.
- (b) Use partial fractions to solve $\int \frac{x^2}{(x^2 - 4)^2} dx$.
- (c) Could the integral in (b) be solved by trigonometric substitution? Explain your answer.
- (d) Find $\int \frac{\sqrt{x+4}}{x^2} dx$. (Hint: Substitute $u = \sqrt{x+4}$, and then use the result in part (b)!)
4. (a) In a complete sentence, tell why there needs to be an even number of subintervals in Simpson's Rule.
- (b) Does there need to be an even number of subintervals in the Trapezoidal Rule? Explain your answer in a complete sentence.
5. Suppose f is continuous and positive on the interval $[x_{k-1}, x_{k+1}]$, and let $x_{k-1} < x_k < x_{k+1}$.
- (a) Draw the trapezoid above the interval $[x_{k-1}, x_k]$, with vertices $(x_{k-1}, 0)$, $(x_k, 0)$, $(x_k, f(x_k))$, $(x_{k-1}, f(x_{k-1}))$. Then find the area A of the trapezoid.
- (b) Using the result of (a), find the sum A_1 of the area of two trapezoids, the one trapezoid above the interval $[x_{k-1}, x_k]$ and the other trapezoid above the interval $[x_k, x_{k+1}]$. Draw a picture for the two trapezoids.
- (c) Using the formula derived in (b) as a guide, explain why the 2's appear inside the brackets of the Trapezoidal Rule ((2) in Section 8.6). Why is there no 2 accompanying $f(x_0)$ or $f(x_n)$?

Solution.

1. (a)

$$\frac{x^2}{x^2 - x + 1} = 1 + \frac{x - 1}{x^2 - x + 1}$$

- (b) Not needed, since the degree of numerator is less than that of the denominator.

- (c)

$$\frac{x^4 + 1}{x^2 + 4} = x^2 - 1 + \frac{5}{x^2 + 1}$$

2. (a)

$$\frac{6x^2 - 4x + 1}{(x - 2)^2} = \frac{A_1}{x - 2} + \frac{A_2}{(x - 2)^2} + \frac{A_3}{(x - 2)^3}$$

- (b)

$$\frac{x^3 + 2x^2 + 3x + 4}{x^4 - 16} = \frac{A_1}{x - 2} + \frac{B_1}{x + 2} + \frac{C_1x + D_1}{x^2 + 2}$$

3. (a)

$$\frac{1}{4} \int \left(\frac{1}{x - 2} + \frac{1}{x + 2} \right)^2 = \frac{1}{4} \int \left(\frac{2x}{(x - 2)(x + 2)} \right)^2 = \left(\frac{x}{x^2 - 4} \right)^2 = \frac{x^2}{(x^2 - 4)^2}$$

(b)

$$\begin{aligned}\int \frac{x^2}{(x^2-4)^2} dx &= \frac{1}{4} \int \left(\frac{1}{x-2} + \frac{1}{x+2} \right)^2 dx \\&= \frac{1}{4} \int \frac{1}{(x-2)^2} + \frac{1}{(x+2)^2} + \frac{2}{(x-2)(x+2)} dx \\&= \frac{1}{4} \int \frac{1}{(x-2)^2} + \frac{1}{(x+2)^2} + \frac{1/2}{x-2} + \frac{-1/2}{x+2} dx \\&= -\frac{1}{4(x-2)} - \frac{1}{4(x+2)} + \frac{1}{2} \ln|x-2| - \frac{1}{2} \ln|x+2| + C\end{aligned}$$

(c) We can substitute x for $2 \sec u$.


(d)

$$\int \frac{\sqrt{x+4}}{x^2} dx \stackrel{\substack{u=\sqrt{x+4} \\ x=u^2-4}}{=} \int \frac{u}{(u^2-4)^2} 2u du = 2 \int \frac{u^2}{(u^2-4)^2} du$$

4. (a) For Simpson's Rule, you need to add up areas under parabolas which is determined by three points and two intervals, so there needs to be an even number of subintervals.

(b) No need for Trapezoidal rules.

5.



(a)

$$A = \frac{x_k - x_{k-1}}{2} (f(x_{k-1}) + f(x_k))$$

(b)

$$A_1 = \frac{x_k - x_{k-1}}{2} (f(x_{k-1}) + f(x_k)) + \frac{x_{k+1} - x_k}{2} (f(x_k) + f(x_{k+1}))$$

(c) The middle sides get added twice whereas the end sides get only added once.

□


1.15 Worksheet for Section 8.6 and 8.7

1. Suppose f is continuous on $[a, b]$ and the graph of f is concave downward on (a, b) . Does this mean that the Trapezoidal Rule yields a number that is larger than, or a number that is smaller than, the integral $\int_a^b f(x) dx$? Draw a picture with your answer and explanation.
2. (a) Find the exact value A of $\int_0^1 x^3 dx$ by performing the integration.
- (b) Approximate the integral in (a) by Simpson's Rule with $n = 4$, obtaining the value B . Tell what the relationship between A and B is, and confirm it by using (8) in Section 8.6.

- (c) Tell why the Trapezoidal Rule for the integral in part (a) with any value of $n > 1$ is (should be) larger than the exact value of the integral. (*Hint:* Draw a graph of the function x^3 on the appropriate interval, and appropriate line segments corresponding to the Trapezoidal Rule.) Then corroborate the assertion by obtaining the Trapezoidal Rule value C with $n = 50$.
3. Consider $\int_{-1}^1 x^4 dx$. Find the exact value D of the integral, and Simpson's Rule's value E , where $n = 2$. By comparing the graphs of x^2 and x^4 , tell why D should indeed be less than E .
4. Find examples of functions f , g and h that are continuous on $[1, \infty)$, and such that
 (a) $\int_1^\infty f(x)dx = \infty$ (b) $\int_1^\infty g(x)dx = 2$ (c) $\int_1^\infty h(x)dx$ does not converge to equal ∞ or $-\infty$.
5. (a) Suppose f and g are continuous on $[a, \infty)$. Prove that if both $\int_a^\infty f(x)dx$ and $\int_a^\infty g(x)dx$ converge, then $\int_a^\infty (f(x) + g(x))dx$ converges.
 (b) Give an example of f and g are continuous on $[a, \infty)$, for which $\int_a^\infty (f(x) + g(x))dx$ converges, but neither $\int_a^\infty f(x)dx$ nor $\int_a^\infty g(x)dx$ converges
6. For (a) and (b), determine whether the area A of the region of the region between the graph of f and the x -axis on the interval $[0, \infty)$ is finite, or is infinite. If it is finite, determine its numerical value.
 (a) $\int_0^\infty \frac{x^3}{\sqrt{2+x^4}} dx$ (b) $\int_0^\infty \frac{x^3}{(2+x^4)^2} dx$

Solution.

1. The approximation is smaller than the actual integral since the trapezoids are below the graph of the function.



2. (a)

$$A = \int_0^1 x^3 dx = \frac{1}{4}$$

(b)

$$B = \frac{1/4}{3}(0^3 + 4 \cdot (1/4)^3 + 2 \cdot (1/2)^3 + 4 \cdot (3/4)^3) = \frac{1}{4}$$

$A = B$ because $(x^3)^{(4)} = 0$ and the error

$$E \leq \frac{\max |f^{(4)}(x)|}{180n^4} (b-a)^4 = 0$$


Which implies that $E = 0$.

(c) Similar to 1 but the function concave downwards.

3. Using Simpson's rule we have

$$E = \frac{1}{3}((-1)^4 + 4 \cdot 0^4 + 1^4) = \frac{2}{3}$$

We also know that $D = \int_{-1}^1 x^4 dx = \frac{2}{5}$. We see that x^2 is the parabolic approximation of x^4 passing $(-1, 1)$, $(0, 0)$, $(1, 1)$, and $D < E$.



4. (a) $f(x)$ could be x

(b) $g(x)$ could be $2/x^2$

(c) $h(x)$ could be $\sin x$

5. (a)

$$\int_a^\infty (f(x) + g(x)) dx = \int_a^\infty f(x) dx + \int_a^\infty g(x) dx$$

since the right hand side converges, so is the left hand side.

(b) $f(x) = x$, $g(x) = -x$.

6. (a)

$$\int \frac{x^3}{\sqrt{2+x^4}} dx \stackrel{u=2+x^4}{=} \frac{du}{4\sqrt{u}} = \frac{1}{2}\sqrt{u} + C = \frac{1}{2}\sqrt{2+x^4} + C$$

So

$$\int_0^\infty \frac{x^3}{\sqrt{2+x^4}} dx = \lim_{b \rightarrow \infty} \int_0^b \frac{x^3}{\sqrt{2+x^4}} dx = \lim_{b \rightarrow \infty} \left. \frac{1}{2}\sqrt{2+x^4} \right|_0^b = \infty$$

(b)

$$\int \frac{x^3}{(2+x^4)^2} dx \stackrel{u=2+x^4}{=} \frac{du}{4u^2} = -\frac{1}{4u} + C = -\frac{1}{4(2+x^4)} + C$$

So

$$\int_0^\infty \frac{x^3}{(2+x^4)^2} dx = \lim_{b \rightarrow \infty} \int_0^b \frac{x^3}{(2+x^4)^2} dx = \left(-\frac{1}{4(2+b^4)} \right) - \left(-\frac{1}{4(2+0)} \right) = \frac{1}{8}$$

□

2 Exams

2.1 Exam 1

(1) (10 points)

Let $f(x) = x^2$ and $g(x) = 2 - x^2$, and let Ω be the region in between these two curves.

- (a) Draw a picture of the region and a sketch of the solid generated by revolving Ω about the x -axis.
- (b) Write down an expression for the volume of this solid of revolution. DO NOT EVALUATE THE INTEGRAL.

Solution.

- (a)
- (b) First solve $x^2 = 2 - x^2 \Rightarrow x = \pm 1$, so

$$V = \int_{-1}^1 \pi[(2 - x^2)^2 - x^4]dx$$

□

(2) (10 points)

Consider the region Ω from Question 1, and assume that its area is $8/3$.

- (a) Write down the integrals that give the moments M_y and M_x about the y - and x -axis.
- (b) Write down expressions for the center of gravity (\bar{x}, \bar{y}) in the terms of these moments.
- (c) Find the values of \bar{x} and \bar{y} (with or without the integrals.)

Solution.

(a)

$$M_y = \int_{-1}^1 x(2 - 2x^2)dx, \quad , M_x = \frac{1}{2} \int_{-1}^1 [(2 - x^2)^2 - x^4]dx$$

(b)

$$\bar{x} = \frac{M_y}{A}, \quad \bar{y} = \frac{M_x}{A}$$

(c) We can use symmetry for both to see that

$$(\bar{x}, \bar{y}) = (0, 1)$$

□

(3) (10 points)

Consider a tank of water in the form of a half-cylinder on its side (see the picture), where the semi-circle has radius 2 feet, and the tank is 20 feet long (so a horizontal cross-section is a rectangle). A pump is placed at the top of a full tank of water, and the water is pumped out of the tank until the tank has depth 1 foot. How much work is done on the water in this process? Water weighs 62.5 lb/ft^3 . DO NOT EVALUATE THE INTEGRAL.

Hint: Place $x = 0$ at the bottom of the tank.

Solution.

$$W = \int_1^2 62.5 \cdot 20(16 - 4x^2)(2 - x)dx$$

□

(4) (10 points)

Find the length of the curve $y = \frac{2}{3}x^{3/2} + 5$ from $x = 0$ to $x = 3$. You do not need to simplify your answer.

Solution. First evaluate $f'(x) = x^{1/2}$, so

$$L = \int_0^1 \sqrt{1 + f'(x)^2} dx = \int_0^1 \sqrt{1 + x} dx \stackrel{u=1+x}{=} \int_1^2 \sqrt{u} du = \frac{2}{3} \left(u^{3/2} \right) \Big|_1^2 = \frac{14}{3}$$

□

(5) (10 points)

Consider a particle that is moving in a circular path given parametrically by

$$\begin{cases} x(t) = 3 + 2 \cos(2t) \\ y(t) = 2 - 2 \sin(2t) \end{cases} \quad \text{for } 0 \leq t \leq 4\pi$$

Find an expression for the speed $v(t)$ of the particle at time t . Determine the following information:

- Center of the circle.
- Radius of the circle.
- How many revolutions of the circle the particle makes.
- Which direction (clockwise or counterclockwise) the particle moves around the circle.

Solution. First we have $x'(t) = -4 \sin(2t)$, $y'(t) = -4 \cos(2t)$, and $v(t) = \sqrt{x'(t)^2 + y'(t)^2} = 4$ □

- The center of the circle is $(3, 2)$.
- The radius of the circle is 2.
- 4 revolutions.
- Clockwise.

2.2 Exam 2

(1) (points) Let $f(x) = x^5 + 3x^3 + 2x - 2$.

- Show that $f(x)$ has an inverse on the entire real line.
- Find $f^{-1}(-2)$ by inspection, and then find $(f^{-1})'(-2)$.

Note. you can do this without solving part (a).

Solution.

- $f'(x) = 5x^4 + 9x^2 + 2 \geq 2 > 0$, so f is strictly increasing, and it has an inverse on the entire real line.
- By inspection, $f(0) = -2$, so

$$(f^{-1})'(-2) = \frac{1}{f'(0)} = \frac{1}{2}$$

□

(2) (points) Find the derivative of

$$f(x) = (\arctan(x))^{2x} + 3^{7x+4} + \log_2(x^3(x+3)^9)$$

Do not simplify your answer.

Solution. First we simplify f

$$f(x) = e^{2x \ln(\arctan(x))} + 3^{7x+4} + 3 \log_2(x) + 9 \log_2(x+3)$$

Therefore

$$f'(x) = (\arctan x)^{2x} \left[2 \ln(\arctan x) + \frac{2x}{\arctan(x)(x^2+1)} \right] + (\ln 3) 3^{7x+4} \cdot 7 + \frac{3}{x \ln 2} + \frac{9}{(x+3) \ln 2}$$

□

(3) (points)

(a) Evaluate $\int \frac{dx}{\sqrt{8x-x^2}}$.

(b) Solve the equation $\sec(\arctan(x)) = 1$ for x .

Solution.

(a) First we complete the squares $8x - x^2 = 16 - (x-4)^2$ Thus

$$\int \frac{dx}{\sqrt{16 - (x-4)^2}} \stackrel{u=x-4}{=} \int \frac{du}{\sqrt{4^2 - u^2}} = \arcsin\left(\frac{u}{4}\right) + C = \arcsin\left(\frac{x-4}{4}\right) + C$$

(b) Suppose $\theta = \arctan x$, then $x = \tan \theta$, so $\sec(\arctan(x)) = \sec \theta =$

□

(4) (points)

(a) Which of the following are indeterminate forms? $0^\infty, \infty^0, \infty^\infty, 1^\infty, \infty^1, 1^0, 0^1, 1^1, 0^0$.

(b) Find the limit: $\lim_{x \rightarrow \infty} x^{(e^{-x})}$ (that is: x raised to the power of e^x)

Solution.

(a) The indeterminate forms are $\infty^0, 1^\infty, 0^0$.

(b)

$$\lim_{x \rightarrow \infty} x^{(e^{-x})} = e^{\lim_{x \rightarrow \infty} \frac{\ln x}{e^x}} \stackrel{\text{L'H } \frac{\infty}{\infty}}{=} e^{\lim_{x \rightarrow \infty} \frac{1}{x e^x}} = e^{\frac{1}{\infty}} = e^0 = 1$$

□

(5) (points)

(a) Evaluate $\int x^2 e^{2x} dx$.

(b) Evaluate $\int_0^1 x^2 e^{2x} dx$.

Solution.

(a)

$$\int x^2 e^{2x} dx$$

(b)

$$\int_0^1 x^2 e^{2x} dx$$

□