Precambrian Research 320 (2019) 484–485

Contents lists available at ScienceDirect

Precambrian Research

journal homepage: www.elsevier.com/locate/precamres

Corrigendum

Corrigendum to "Paleomagnetism, magnetic anisotropy and U-Pb baddeleyite geochronology of the early Neoproterozoic Blekinge-Dalarna dolerite dykes, Sweden" [Precambrian Res. 317 (2018) 14–32]

Zheng Gong[a,*], David A.D. Evans[a], Sten-Åke Elming[b], Ulf Söderlund[c,d], Johanna M. Salminen[e]

[a] Department of Geology and Geophysics, Yale University, 210 Whitney Avenue, New Haven, CT 06511, USA
[b] Department of Civil, Environmental and Natural Resources Engineering, Luleå University of Technology, SE-971 87 Luleå, Sweden
[c] Department of Geology, Lund University, SE-223 62 Lund, Sweden
[d] Swedish Museum of Natural History, Laboratory of Isotope Geology, SE-104 05 Stockholm, Sweden
[e] Department of Geosciences and Geography, University of Helsinki, Helsinki 00014, Finland

The authors regret that the incorrect version of Table 3 and Fig. 12 appeared in the paper. The corrected Table 3 and Fig. 12 are presented below. In Table 2, Group A mean direction should be Dec = 128.8°, Inc = 39.6°, α_{95} = 6.5°, and Group B mean direction should be Dec = 127.6°, Inc = 65.4°, α_{95} = 9.7°. These changes do not affect any main conclusion of the paper. The authors would like to apologize for any inconvenience caused.

Table 3
Paleomagnetic poles used to constrain the paleogeographic reconstruction of Baltica and Laurentia at ca. 1000–850 Ma.

#	Paleomagnetic Pole	Dec (°)	Inc (°)	α_{95} (°)	k	N/n	Plat (°N)	Plon (°E)	A_{95} (°)	S (°)	|λ| (°)	Age (Ma)	1	2	3	4	5	6	7	Q	Reference
Baltica																					
1	Hunnedalen dykes	294.0	−75.0	6.0	115	6/69	−41.0	222.0	10.5	13.4	61.8	848 ± 27[a]; 855 ± 59[b]	1	1	1	0	1	0	1	5	Walderhaug et al. (1999)
2	Egersund-Ogna anorthosite	325.9	−80.1	4.9	73	13/69	−42.1	200.4	9.0	17.3	70.8	900[c]	1	1	1	0	0	0	1	4	Brown and McEnroe (2004)
3	Egersund anorthosite	–	–	–	–	76/–	−43.5	213.7	3.6	–	–	900[c]	0	1	1	0	0	0	1	3	Stearn and Piper (1984); Walderhaug et al. (1999)
4	Rogaland Igneous Complex	269.0	−72.0	11.0	49	5/24	−45.9	238.4	18.2	19.4	57.0	869 ± 14[a]	1	0	1	1	0	0	1	4	Walderhaug et al. (2007)
5	Bjerkreim-Sokndal intrusion	303.4	−73.5	3.7	24	66/354	−35.9	217.9	6.0	26.1	59.4	916[c]	1	1	1	0	1	0	1	5	Brown and McEnroe (2015)
6	Mean 951–935 Ma Baltica pole	308.8	−39.6	6.5	42	12/116	−2.6	239.6	5.8	10.8	22.5	951–935[d]	1	1	1	1	1	1	1	7	This study
7	971 Ma BDD dykes VGP	307.6	−65.4	14.2	68	4/36	−27.0	230.4	14.9	13.0	47.5	971 ± 7[d]	1	0	1	0	1	1	1	5	This study
8	Laanila-Ristijärvi dykes	355.5	−40.0	17.5	51	3/7	−2.1	212.2	16.4	9.0	22.8	1042 ± 50[b]	0	0	1	1	1	0	0	3	Mertanen et al. (1996)
9	Bamble Intrusion mean	–	–	–	–	–	3.0	217.0	15.0	–	–	1100–1040	0	1	1	0	0	0	0	2	Meert and Torsvik (2003)
Laurentia																					
10	Adirondack microcline gneisses	289.2	−62.8	7.6	29	14/80	−18.4	151.1	10.5	22.0	44.2	960[c]	1	1	1	0	0	1	1	5	Brown and McEnroe (2012)

(*continued on next page*)

DOI of original article: https://doi.org/10.1016/j.precamres.2018.08.019
* Corresponding author.

https://doi.org/10.1016/j.precamres.2018.10.013
Received 15 October 2018; Accepted 30 October 2018
Available online 02 November 2018
0301-9268/ © 2018 Elsevier B.V. All rights reserved.

Table 3 (*continued*)

#	Paleomagnetic Pole	Dec (°)	Inc (°)	α_{95} (°)	k	N/n	Plat (°N)	Plon (°E)	A_{95} (°)	S (°)	|λ| (°)	Age (Ma)	1	2	3	4	5	6	7	Q	Reference
11	Adirondack metamorphic anorthosites and other rocks	283.9	−67.3	7.7	28	14/68	−25.1	149.0	11.6	24.3	50.1	970[c]	1	1	1	0	0	1	1	5	Brown and McEnroe (2012)
12	Adirondack fayalite granites	297.0	−75.8	3.9	199	8/40	−28.4	132.7	6.9	10.2	63.2	990[c]	1	1	1	0	0	1	1	5	Brown and McEnroe (2012)

Note: Dec = declination, Inc = inclination, α_{95} = radius of 95% confidence cone of the mean direction, k = precision parameter, N/n = number of site/sample, Plat/Plon = paleomagnetic pole latitude/longitude, A_{95} = radius of 95% confidence cone of the paleomagnetic pole, S = angular dispersion of VGPs, |λ| = absolute value of paleolatitude, Q = sum of quality criteria (Van der Voo, 1990).

[a] Ar-Ar biotite ages.
[b] Sm-Nd whole rock ages.
[c] Cooling ages.
[d] U-Pb baddeleyite ages.




Fig. 12.