

ML Homework 4

Liheng Cao

March 22, 2021

1

When w_0 increases, then the probability of predicting class 1 becomes higher, and vice versa.

$$h(x) = \frac{1}{1 + e^{-(w_0 + \dots)}}$$

This is because as w_0 gets bigger, the result of exponentiation gets smaller. As the denominator gets smaller, the quotient gets larger. $h(x)$ is used as the probability we will predict a certain class. The higher $h(x)$ is, the more likely we will predict class 1 (as opposed to class 0).

2

The logistic function is

$$\frac{1}{1 + e^{-\mathbf{w}^T \mathbf{x}}}$$

If we were to double \mathbf{w} , the result wouldn't actually change, as long as the threshold is 0.5.

The geometric interpretation is because $\mathbf{w}^T \mathbf{x}$ represents a (hyper)plane. Multiplying the equation of a plane doesn't affect any points that are on the plane (threshold = 0.5 \implies on hyperplane), but it makes the points that are off it seem further away, increasing the certainty with which we pick a class. For example, if $\mathbf{w}^T \mathbf{x} = 0$, then there is no change ($0 \cdot 2 = 0$). But if it's positive, it becomes a larger positive number, and vice versa.

3

(a)

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

(b) :




Figure 1: o: class 0, x: class 1

(c) $\text{TPR} = \frac{1}{3+1} = 1/4$

(d) $\text{FPR} = \frac{3}{3+1} = 3/4$

(e) $\text{accuracy} = \frac{3+3}{3+1+3+1} = 3/4$

(f) $\text{recall} = \frac{3}{3+1} = 3/4$

(g) $\text{precision} = \frac{3}{3+1} = 3/4$

(h)

$$-[0 \cdot \ln(0.389) + (1-0) \ln(1-0.389)] - \dots - [1 \cdot \ln(0.638) + (1-1) \ln(1-0.638)] = 4.25205106 \dots$$

(i)

$$-\left[0 \cdot \ln\left(\frac{1}{1 + \exp(-\mathbf{w}'^T \mathbf{X}^{(1)})}\right)\right] + \dots - \left[0 \cdot \ln\left(\frac{1}{1 + \exp(-\mathbf{w}'^T \mathbf{X}^{(8)})}\right)\right] = 3.0158793 \dots$$

Since \mathbf{w}' has a lower cross-entropy, it is a better fit for this data.

(j) After one iteration,

$$\begin{aligned}
 \mathbf{w} &= \mathbf{w} + \frac{\alpha}{N} \mathbf{x}^T (y - \sigma(\mathbf{xw})) \\
 &\quad \begin{bmatrix} 0.66 \\ -2.24 \\ -0.18 \end{bmatrix} \\
 &+ \frac{0.01}{N} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0.49 & 1.69 & 0.04 & 1 & 0.16 & 0.25 & 0.49 & 0.04 \\ 0.09 & 0.04 & 0.64 & 0.16 & 0.09 & 0 & 0 & 0.01 \end{bmatrix} \begin{pmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \sigma \left(\begin{bmatrix} 1 & 0.49 & 0.09 \\ 1 & 1.69 & 0.04 \\ 1 & 0.04 & 0.64 \\ 1 & 1 & 0.16 \\ 1 & 0.16 & 0.09 \\ 1 & 0.25 & 0 \\ 1 & 0.49 & 0 \\ 1 & 0.04 & 0.01 \end{bmatrix} \begin{bmatrix} 0.66 \\ -2.24 \\ -0.18 \end{bmatrix} \right) \end{pmatrix} \\
 &= \begin{bmatrix} 0.661 \dots \\ -2.239 \dots \\ -0.181 \dots \end{bmatrix}
 \end{aligned}$$

(k) These data points would contribute a medium amount to the new \mathbf{w} .

(l) These data points would contribute a low amount to the new \mathbf{w} .

(m) These data points would contribute a high amount to the new \mathbf{w} .

(n)

$$- \left[0 \cdot \ln \left(\frac{1}{1 + \exp(-\mathbf{w}_g^T \mathbf{X}^{(1)})} \right) \right] + \dots - \left[0 \cdot \ln \left(\frac{1}{1 + \exp(-\mathbf{w}_g^T \mathbf{X}^{(8)})} \right) \right] = 4.2519 \dots$$

This is a very minor decrease from 4.2520. This makes sense because gradient ascent is supposed to make the model fit better and have less error.

4

- lasso

$$-\lambda (|w_1| + \cdots + |w_d|) + \sum_{i=1}^N y^{(i)} \ln(h(x)) + (1 - y^{(i)}) \ln(1 - h(x))$$

- ridge

$$-\lambda (w_1^2 + \cdots + w_d^2) + \sum_{i=1}^N y^{(i)} \ln(h(x)) + (1 - y^{(i)}) \ln(1 - h(x))$$

- gradient

$$\mathbf{x}^T(y - \sigma(\mathbf{x}\mathbf{w})) + 2\lambda I' \mathbf{w}$$

- The training error increases, but the validation error decreases.

5

- (a) Predictors: average pitch; response: gender
- (b) Predictors: matrix of where the strokes were; response: letter or number written

6




Figure 2: o: no donation, x: donation

(a)

(b) I will choose the horizontal line $x_2 = 0.5 \implies 0.5 + 0 \cdot x_1 + w_2$

$$\mathbf{w}^T = [-0.5, 0, 1]$$

- (c) The least likely would be the one that is incorrect. In this case, it's same 3, or the one that earns 70k, follows 1 site, and did not donate. No calculator was needed for this!
- (d) They would not change the values of \hat{y} , but they would change the likelihoods. If the prediction was correct, then the likelihood would increase. If the predictions were incorrect, then the likelihood would decrease.