Supporting Information

Amplitude and midpoint statistics of the same chemotaxis cells demand incompatible receptor models

Huan Li

Shanghai Liangta Biotech Co., Ltd., Shanghai 201899, P. R. China

October 2026

This Supporting Information documents the complete computational output of the accompanying repository (`chemotaxis-receptor-layer`, release v1.0.1): every analysis script, the numerical results each produced, and the provenance of the third-party FRET archive. Numbers are transcribed verbatim from `results/`; scripts are plain Python 3 (numpy/scipy/matplotlib) and read the `.mat` files in `data_external/moore2024_fret/` directly.

Contents

1 Pipeline map: scripts and outputs **2**

2 P7 adjudication: $K_{1/2}$ medians vs the Weber line **3**

3 Adaptation-relay baseline audit **3**

4 Prospective mixed-ligand arms (code70) **3**

5 Third-party data provenance **3**

1 Pipeline map: scripts and outputs

Script	Computes	Archived output
code49_P7_blind_ adjudication_ FCD_Weber	P7 blind adjudication of the fold-change/Weber claim on the Moore 2024 archive	adjudication tables
code49b_P7_erratum_ readjudication	erratum re-adjudication (censoring-aware)	the code49b unit-table CSV (results/P7_ Moore2024_ FCD-Weber/code49b_ unit_table.csv)
code51chemo_receptor_ layer_ selfcheck	receptor-layer internal-consistency self-check	console log
code52_protocol_ artifact_test_ raw_FRET_ coordinate_analysis	protocol-artifact test in raw FRET coordinates	console/tables
code53_CheA_node_ mechanism_ kernel_ competition	CheA-node mechanism: kernel competition	console/tables
code54_K12_joint_fit_ rupture_ quantification	joint-fit rupture quantification on the $K_{1/2}$ ladder	console/tables
code55_multispecies_ local_ adaptation_joint_ test	multi-species / local-adaptation joint test	console/tables
code56_FRET_readout_ nonlinearity_ test	FRET readout-nonlinearity control	console/tables
code57_TCS_depletion_ robustness_ gain_ retuning	TCS-depletion robustness with gain retuning	console/tables
code58_P8_independent_ arm_ serine_Lasp	P8 independent arm (serine, L-aspartate)	console/tables
code59_residual_ waterfall_ window_ structure_rhoF	residual waterfall; window structure of ρ_F	console/tables
code60_heterogeneous_ population_ censoring_ test	heterogeneous-population censoring control	console/tables
code61_rowwise_Hill_ steepness_ test	row-wise Hill-steepness test	console/tables
code62_single_cell_same_ source_ statistics	same-source single-cell statistics	console/tables
code70_mixed_ligand_ validation_ arm_P1P2	prospective mixed-ligand validation arms P1/P2	chemotaxis_ prospective_arm.csv
chemotaxis_adaptation_ audit.py	adaptation-relay baseline audit on the full archive	chemotaxis_ adaptation_audit.json

2 P7 adjudication: $K_{1/2}$ medians vs the Weber line

Censoring-aware re-adjudication (code49b) of the FCD–Weber claim on the Moore et al. (2024) single-cell FRET archive. Table 2 is the archived unit table (verbatim rows).

Table 2: $K_{1/2}$ medians per background level vs Weber-line prediction.

Background (μM)	$K_{1/2}$ median (μM)	IQR lo–hi	censored / n	Weber pred. (μM)	dev. (dex)
0.0	2.033	1.583–3.105	179/268	2.282	−0.050
0.01	2.896	2.080–3.065	69/101	2.293	+0.101
0.1	2.168	1.769–2.844	35/57	2.399	−0.044

The per-cell unit table (`results/P7_Moore2024_FCD-Weber/code49b_unit_table.csv`) carries, per (background B, foreground F) condition: fold change r, threshold T, $\log_{10} r$, $\log_{10} T$, n cells, and the two arm statistics R_a, R_A; e.g. B=0.01, F=0.2 μM: $r = 21.0$, $T = 0.21$, $n = 101$, $R_a = 0.0655$, $R_A = 0.0209$.

3 Adaptation-relay baseline audit

Full-archive audit (`chemotaxis_adaptation_audit.json`): 2217 cells, 77,595 response events; median baseline activity after high- vs low-amplitude preceding stimuli 0.434 vs 0.448; median absolute response 0.608. The per-(preceding-dose) baseline bands are archived in the same JSON.

4 Prospective mixed-ligand arms (code70)

Frozen-then-measured prospective arm V1 (clause P-1b, amplitude table of the L-aspartate component). Verbatim rows of `chemotaxis_prospective_arm.csv`:

Table 3: Prospective arm V1: measured vs predicted amplitudes (tolerance: max |dev| = 0.08).

Foreground (μM)	measured	predicted	deviation	hit
5	0.047	0.119	−0.072	Y
10	0.085	0.152	−0.067	Y
20	0.139	0.187	−0.048	Y

5 Third-party data provenance

`data_external/moore2024_fret/` mirrors, verbatim, the 48-file single-cell FRET archive of Moore, Kamino, Kottou, Shimizu & Emonet, *Cell Syst.* 15, 628–638 (2024), Dryad doi:10.5061/dryad.nvx0k6dzz, under its CC0 1.0 public-domain dedication; please cite the original study. Receptor-layer parameter priors derive from Sourjik & Berg, *Proc. Natl Acad. Sci. USA* 99, 123–127 (2002); Shimizu, Tu & Berg, *Mol. Syst. Biol.* 6, 382 (2010); and Tu, Shimizu & Berg, *Proc. Natl Acad. Sci. USA* 105, 14855–14860 (2008); fold-change readout evidence from Lazova et al., *Proc. Natl Acad. Sci. USA* 108, 13870–13875 (2011). Full notes in `data_external/PROVENANCE.md`.