



# Logistic Regression Inference

---

Julie Deeke

Statistics with Python Course Developer



# Cartwheel Data


Random sample of 25 adults attempted cartwheels

**Primary Variable of interest:** Cartwheel completion



Based on age, can we predict whether a cartwheel is completed?

# Logistic Regression Line



## Generalized Linear Model Regression Results

**Dep. Variable:** CompleteGroup **No. Observations:** 25  
**Model:** GLM **Df Residuals:** 23  
**Model Family:** Binomial **Df Model:** 1  
**Link Function:** logit **Scale:** 1.0  
**Method:** IRLS **Log-Likelihood:** -12.534  
**Date:** Tue, 27 Nov 2018 **Deviance:** 25.068  
**Time:** 16:11:20 **Pearson chi2:** 22.4  
**No. Iterations:** 6

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$$b_1 \pm t^* \text{se}(b_1)$$

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$$b_1 \pm t^* \text{se}(b_1)$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$$b_1 \pm t^* se(b_1)$$


	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$b_1 \pm t^* se(b_1)$

  $df = n-2$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545



# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

$(-0.1442, 0.5634)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Confidence Interval

**Best Estimate  $\pm$  Margin of Error**

**Sample slope  $\pm$  “a few”  $\cdot$  estimated standard error**

$df = n - 2$

$b_1 \pm t^* se(b_1)$

$0.2096 \pm 2.069 (0.171)$

$(-0.1442, 0.5634)$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

0 is included

# IVQ

Does it seem reasonable that there is a significant slope?  
*No*

We'll continue working through the hypothesis testing framework before coming back to the answer of the IVQ.

# Hypothesis Testing

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic:  $\frac{b_1 - 0}{\text{s.e.}(b_1)}$

← null hypothesis

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

Test statistic: 
$$\frac{b_1 - 0}{\text{s.e.}(b_1)}$$

	coef	std err	z	P> z	[0.025	0.975]
<b>Intercept</b>	-4.4213	4.429	-0.998	0.318	-13.101	4.259
<b>Age</b>	0.2096	0.171	1.225	0.221	-0.126	0.545



# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171}$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Hypothesis Testing

$$H_0: \beta_1 = 0$$

$$H_a: \beta_1 \neq 0$$

*need significance level*

With our p-value of about 0.221, we would fail to reject the null hypothesis and cannot conclude that we have a significantly linear relationship between age and the log odds of the probability of successfully completing a cartwheel.

$$\text{Test statistic: } \frac{b_1 - 0}{\text{s.e.}(b_1)} = \frac{0.2096}{0.171} = 1.225$$

	coef	std err	z	P> z	[0.025	0.975]
Intercept	-4.4213	4.429	-0.998	0.318	-13.101	4.259
Age	0.2096	0.171	1.225	0.221	-0.126	0.545

# Summary

Logistic regression for a predicted variable with two options  
~correct/incorrect or success/failure

# Summary

Logistic regression for a predicted variable with two options

~correct/incorrect or success/failure

Confidence Intervals and Hypothesis Tests follow similar format as linear regression

~changing interpretation to the logistic context

# Summary

Logistic regression for a predicted variable with two options

~correct/incorrect or success/failure

Confidence Intervals and Hypothesis Tests follow similar format as Linear Regression

~changing interpretation to the logistic context

Coming up, we'll look at an example from NHANES using blood pressure and smoking