

# 机器学习与数据挖掘

*Machine Learning & Data Mining*


权小军 教授

中山大学数据科学与计算机学院


[quanxj3@mail.sysu.edu.cn](mailto:quanxj3@mail.sysu.edu.cn)

# **Preface**

# Preface




# Preface



# Preface

## The Evils of “Hard Assignments”?



- Clusters may overlap
- Some clusters may be “wider” than others
- Distances can be deceiving!

# **Lecture 16:**

# **Expectation Maximization**

# A Generative View of Clustering

- We need a sensible measure of what it means to cluster the data well
  - This makes it possible to judge different methods
  - It may help us decide on the number of clusters
- An obvious approach is to imagine that the data was produced by a generative model
  - Then we adjust the model parameters to maximize the probability that it would produce exactly the data we observed

# Generative Models

- We model the joint distribution as,

$$p(\mathbf{x}, z) = p(\mathbf{x}|z)p(z)$$

- But in unsupervised clustering we do not have the class labels  $z$ .
- What can we do instead?

$$p(\mathbf{x}) = \sum_z p(\mathbf{x}, z) = \sum_z p(\mathbf{x}|z)p(z)$$

- This is a **mixture model**.




# The General GMM assumption

- $P(Z)$ : There are  $k$  components
- $P(X|Z)$ : Each component generates data from a **multivariate Gaussian** with mean  $\mu_i$  and covariance matrix  $\Sigma_i$

# The General GMM assumption

Each data point is sampled from a **generative process**:

1. Choose component  $i$  with probability  $P(z=i)$
2. Generate datapoint  $\sim N(m_i, \Sigma_i)$



# Gaussian Mixture Model (GMM)

Most common mixture model: [Gaussian mixture model](#) (GMM)

- A GMM represents a **distribution** as

$$p(\mathbf{x}) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)$$

with  $\pi_k$  the [mixing coefficients](#), where:

$$\sum_{k=1}^K \pi_k = 1 \quad \text{and} \quad \pi_k \geq 0 \quad \forall k$$

- GMM is a density estimator
- GMMs are **universal approximators of densities** (if you have enough Gaussians). Even diagonal GMMs are universal approximators.
- In general mixture models are very powerful, but harder to optimize

# Gaussian Mixture Model (GMM)

Most common mixture model: Gaussian mixture model (GMM)

- A GMM represents a **distribution** as

$$p(\mathbf{x}) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)$$

with  $\pi_k$  the **mixing coefficients**, where:

$$\sum_{k=1}^K \pi_k = 1 \quad \text{and} \quad \pi_k \geq 0 \quad \forall k$$

- GMM is a density estimator

In probability and statistics, **density estimation** is the construction of an estimate, based on observed data, of an unobservable underlying probability density function.

- In general mixture models are very powerful, but harder to optimize

# Gaussian Mixture Model (GMM)

Most common mixture model: Gaussian mixture model (GMM)

- A GMM represents a **distribution** as

$$p(\mathbf{x}) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)$$


with  $\pi_k$  the **mixing coefficients**, where:

$$\sum_{k=1}^K \pi_k = 1 \quad \text{and} \quad \pi_k \geq 0 \quad \forall k$$


- GMM is a density estimator
- GMMs are **universal approximators of densities** (if you have enough Gaussians).
- In general mixture models are very powerful, but harder to optimize

# Visualizing a Mixture of Gaussians – 1D Gaussians

- If you fit a Gaussian to data:




- Now, we are trying to fit a GMM (with  $K = 2$  in this example):




# Visualizing a Mixture of Gaussians – 2D Gaussians

## Old Faithful Data Set






Single Gaussian



Mixture of two Gaussians

# Visualizing a Mixture of Gaussians – 2D Gaussians





# Fitting GMMs: Maximum Likelihood

- Maximum likelihood maximizes

$$\ln p(\mathbf{X}|\pi, \mu, \Sigma) = \sum_{n=1}^N \ln \left( \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}^{(n)}|\mu_k, \Sigma_k) \right)$$

$$\text{w.r.t } \Theta = \{\pi_k, \mu_k, \Sigma_k\}$$

- How would you optimize this?
- Can we have a closed form update?
- Don't forget to satisfy the constraints on  $\pi_k$  and  $\Sigma_k$

# Latent Variable

- Our original representation had a hidden (latent) variable  $z$  which would represent which Gaussian generated our observation  $\mathbf{x}$ , with some probability
- Let  $z \sim \text{Categorical}(\boldsymbol{\pi})$  (where  $\pi_k \geq 0$ ,  $\sum_k \pi_k = 1$ )

■ Then:

$$\begin{aligned} p(\mathbf{x}) &= \sum_{k=1}^K p(\mathbf{x}, z = k) \\ &= \sum_{k=1}^K \underbrace{p(z = k)}_{\pi_k} \underbrace{p(\mathbf{x} | z = k)}_{\mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)} \end{aligned}$$

- This breaks a complicated distribution into simple components.

# Latent Variable Models

- Some model variables may be unobserved, either at training or at test time, or both
- If occasionally unobserved they are missing, e.g., undefined inputs, missing class labels, erroneous targets
- Variables which are always unobserved are called **latent variables**, or sometimes **hidden variables**
- We may want to intentionally introduce latent variables to model complex dependencies between variables – this can actually simplify the model
- Form of divide-and-conquer: use simple parts to build complex models
- In a **mixture model**, the identity of the component that generated a given datapoint is a latent variable

# Back to GMM

- A Gaussian mixture distribution:  $p(\mathbf{x}) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)$
- We had:  $\mathbf{z} \sim \text{Categorical}(\boldsymbol{\pi})$  (where  $\pi_k \geq 0$ ,  $\sum_k \pi_k = 1$ )

- Joint distribution:  $p(\mathbf{x}, \mathbf{z}) = p(\mathbf{z})p(\mathbf{x}|\mathbf{z})$

- Log-likelihood:

$$\begin{aligned}\ell(\boldsymbol{\pi}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) &= \ln p(\mathbf{X} | \boldsymbol{\pi}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \sum_{n=1}^N \ln p(\mathbf{x}^{(n)} | \boldsymbol{\pi}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) \\ &= \sum_{n=1}^N \ln \sum_{z^{(n)}=1}^K p(\mathbf{x}^{(n)} | z^{(n)}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) p(z^{(n)} | \boldsymbol{\pi})\end{aligned}$$

- Note: We have a hidden variable  $z^{(n)}$  for every observation
- General problem: sum inside the log
- How can we optimize this?

# Back to GMM

- A Gaussian mixture distribution:  $p(\mathbf{x}) = \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x} | \mu_k, \Sigma_k)$
- We had:  $\mathbf{z} \sim \text{Categorical}(\boldsymbol{\pi})$  (where  $\pi_k \geq 0$ ,  $\sum_k \pi_k = 1$ )

- Joint distribution:  $p(\mathbf{x}, \mathbf{z}) = p(\mathbf{z})p(\mathbf{x}|\mathbf{z})$

- Log-likelihood:

$$\begin{aligned}\ell(\boldsymbol{\pi}, \mu, \Sigma) &= \ln p(\mathbf{X} | \boldsymbol{\pi}, \mu, \Sigma) = \sum_{n=1}^N \ln p(\mathbf{x}^{(n)} | \boldsymbol{\pi}, \mu, \Sigma) \\ &= \sum_{n=1}^N \ln \sum_{k=1}^K p(\mathbf{x}^{(n)} | z^{(n)}; \mu, \Sigma) p(z^{(n)} | \boldsymbol{\pi})\end{aligned}$$

## Relation to neural networks!

- Note: We have a hidden variable  $z^{(n)}$  for every observation
- General problem: sum inside the log
- How can we optimize this?

# Maximum Likelihood

- If we knew  $z^{(n)}$  for every  $x^{(n)}$ , the maximum likelihood problem is easy:

$$\ell(\boldsymbol{\pi}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \sum_{n=1}^N \ln p(x^{(n)}, z^{(n)} | \boldsymbol{\pi}, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \sum_{n=1}^N \ln p(\mathbf{x}^{(n)} | z^{(n)}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) + \ln p(z^{(n)} | \boldsymbol{\pi})$$

- We would get this:

$$\begin{aligned}\mu_k &= \frac{\sum_{n=1}^N 1_{[z^{(n)}=k]} \mathbf{x}^{(n)}}{\sum_{n=1}^N 1_{[z^{(n)}=k]}} \\ \Sigma_k &= \frac{\sum_{n=1}^N 1_{[z^{(n)}=k]} (\mathbf{x}^{(n)} - \mu_k)(\mathbf{x}^{(n)} - \mu_k)^T}{\sum_{n=1}^N 1_{[z^{(n)}=k]}} \\ \pi_k &= \frac{1}{N} \sum_{n=1}^N 1_{[z^{(n)}=k]}\end{aligned}$$

# How Can We Fit a Mixture of Gaussians?

- Optimization uses the **Expectation Maximization algorithm**, which alternates between two steps:
  1. **E-step**: Compute the posterior probability over  $z$  given our current model
    - i.e. how much do we think each Gaussian generates each datapoint.
  2. **M-step**: Assuming that the data really was generated this way, change the parameters of each Gaussian to maximize the probability that it would generate the data it is currently responsible for.

# Relation to k-Means

## ■ The K-Means Algorithm:

1. **Assignment step**: Assign each data point to the closest cluster
2. **Refitting step**: Move each cluster center to the center of gravity of the data assigned to it



# Relation to k-Means

## ■ The K-Means Algorithm:

1. Assignment step: Assign each data point to the closest cluster
2. Refitting step: Move each cluster center to the center of gravity of the data assigned to it

## ■ The EM Algorithm:

1. **E-step**: Compute the posterior probability over  $z$  given our current model
2. **M-step**: Maximize the probability that it would generate the data it is currently responsible for.

# Expectation Maximization for GMM Overview

- Elegant and powerful method for finding maximum likelihood solutions for models with latent variables

## 1. E-step:

- In order to adjust the parameters, we must first solve the inference problem: Which Gaussian generated each datapoint?
- We cannot be sure, so it's a distribution over all possibilities.

$$\gamma_k^{(n)} = p(z^{(n)} = k | \mathbf{x}^{(n)}; \pi, \mu, \Sigma)$$

## 2. M-step:

- Each Gaussian gets a certain amount of posterior probability for each datapoint.
- We fit each Gaussian to the weighted datapoints
- We can derive closed form updates for all parameters

# Where does EM come from? I

- Remember that optimizing the likelihood is hard because of the sum inside of the log. Using  $\Theta$  to denote all of our parameters:

$$\ell(\mathbf{X}, \Theta) = \sum_i \log(P(\mathbf{x}^{(i)}; \Theta)) = \sum_i \log \left( \sum_j P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta) \right)$$

- We can use a common trick in machine learning, introduce a new distribution,  $q$ :

$$\ell(\mathbf{X}, \Theta) = \sum_i \log \left( \sum_j q_j \frac{P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta)}{q_j} \right)$$

- Now we can swap them! Jensen's inequality - for **concave** function (like log)

$$f(\mathbb{E}[x]) = f \left( \sum_i p_i x_i \right) \geq \sum_i p_i f(x_i) = \mathbb{E}[f(x)]$$

# Where does EM come from? II

- Applying Jensen's,

$$\sum_i \log \left( \sum_j q_j \frac{P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta)}{q_j} \right) \geq \sum_i \sum_j q_j \log \left( \frac{P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta)}{q_j} \right)$$

- Maximizing this lower bound will force our likelihood to
- increase. But how do we pick a  $q_j$  that gives a good bound?

# EM derivation

- We got the sum outside but we have an inequality.

$$\ell(\mathbf{X}, \Theta) \geq \sum_i \sum_j q_j \log \left( \frac{P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta)}{q_j} \right)$$

- Lets fix the current parameters to  $\Theta^{old}$  and try to find a good  $q_i$
- What happens if we pick  $q_j = p(z^{(i)} = j | \mathbf{x}^{(i)}, \Theta^{old})$ ?

$$\frac{P(\mathbf{x}^{(i)}, z^{(i)}; \Theta)}{p(z^{(i)}=j | \mathbf{x}^{(i)}, \Theta^{old})} = P(\mathbf{x}^{(i)}; \Theta^{old}) \text{ and the inequality becomes an equality!}$$

- We can now define and optimize

$$\begin{aligned} Q(\Theta) &= \sum_i \sum_j p(z^{(i)} = j | \mathbf{x}^{(i)}, \Theta^{old}) \log \left( P(\mathbf{x}^{(i)}, z^{(i)} = j; \Theta) \right) \\ &= \mathbb{E}_{P(z^{(i)} | \mathbf{x}^{(i)}, \Theta^{old})} [\log \left( P(\mathbf{x}^{(i)}, z^{(i)}; \Theta) \right)] \end{aligned}$$

- We ignored the part that doesn't depend on  $\Theta$


# EM derivation

- So, what just happened?
- Conceptually: We don't know  $z^{(i)}$  so we average them given the current model.
- Practically: We define a function  $Q(\Theta) = \mathbb{E}_{P(z^{(i)}|\mathbf{x}^{(i)}, \Theta^{old})}[\log(P(\mathbf{x}^{(i)}, z^{(i)}; \Theta))]$  that lower bounds the desired function and is equal at our current guess.
- If we now optimize  $\Theta$  we will get a better lower bound!

$$\log(P(\mathbf{X}|\Theta^{old})) = Q(\Theta^{old}) \leq Q(\Theta^{new}) \leq \log(P(\mathbf{X}|\Theta^{new}))$$

- We can iterate between **expectation** step and **maximization** step and the lower bound will always improve (or we are done)

# Visualization of the EM Algorithm



- The EM algorithm involves alternately computing a lower bound on the log likelihood for the current parameter values and then maximizing this bound to obtain the new parameter values.

# General EM Algorithm

1. Initialize  $\Theta^{old}$
2. E-step: Evaluate  $p(\mathbf{Z}|\mathbf{X}, \Theta^{old})$  and compute

$$Q(\Theta, \Theta^{old}) = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \Theta^{old}) \ln p(\mathbf{X}, \mathbf{Z}|\Theta)$$

3. M-step: Maximize

$$\Theta^{new} = \arg \max_{\Theta} Q(\Theta, \Theta^{old})$$

4. Evaluate log likelihood and check for convergence (or the parameters). If not converged,  $\Theta^{old} = \Theta^{new}$ , Go to step 2



# GMM E-Step: Responsibilities

Lets see how it works on GMM:

- Conditional probability (using Bayes rule) of  $\mathbf{z}$  given  $\mathbf{x}$

$$\begin{aligned}\gamma_k = p(z = k|\mathbf{x}) &= \frac{p(z = k)p(\mathbf{x}|z = k)}{p(\mathbf{x})} \\ &= \frac{p(z = k)p(\mathbf{x}|z = k)}{\sum_{j=1}^K p(z = j)p(\mathbf{x}|z = j)} \\ &= \frac{\pi_k \mathcal{N}(\mathbf{x}|\mu_k, \Sigma_k)}{\sum_{j=1}^K \pi_j \mathcal{N}(\mathbf{x}|\mu_j, \Sigma_j)}\end{aligned}$$

- $\gamma_k$  can be viewed as the **responsibility** of cluster  $k$  towards  $\mathbf{x}$

# GMM E-Step

- Once we computed  $\gamma_k^{(i)} = p(z^{(i)} = k | \mathbf{x}^{(i)})$  we can compute the expected likelihood

$$\begin{aligned} & \mathbb{E}_{P(z^{(i)} | \mathbf{x}^{(i)})} \left[ \sum_i \log(P(\mathbf{x}^{(i)}, z^{(i)} | \Theta)) \right] \\ &= \sum_i \sum_k \gamma_k^{(i)} \left( \log(P(z^i = k | \Theta)) + \log(P(\mathbf{x}^{(i)} | z^{(i)} = k, \Theta)) \right) \\ &= \sum_i \sum_k \gamma_k^{(i)} \left( \log(\pi_k) + \log(\mathcal{N}(\mathbf{x}^{(i)}; \mu_k, \Sigma_k)) \right) \\ &= \sum_k \sum_i \gamma_k^{(i)} \log(\pi_k) + \sum_k \sum_i \gamma_k^{(i)} \log(\mathcal{N}(\mathbf{x}^{(i)}; \mu_k, \Sigma_k)) \end{aligned}$$

- We need to fit  $k$  Gaussians, just need to weight examples by  $\gamma_k$

# GMM M-Step

- Need to optimize

$$\sum_k \sum_i \gamma_k^{(i)} \log(\pi_k) + \sum_k \sum_i \gamma_k^{(i)} \log(\mathcal{N}(\mathbf{x}^{(i)}; \mu_k, \Sigma_k))$$

- Solving for  $\mu_k$  and  $\Sigma_k$  is like fitting  $k$  separate Gaussians but with weights  $\gamma_k^{(i)}$
- Solution is similar to what we have already seen:

$$\begin{aligned}\mu_k &= \frac{1}{N_k} \sum_{n=1}^N \gamma_k^{(n)} \mathbf{x}^{(n)} \\ \Sigma_k &= \frac{1}{N_k} \sum_{n=1}^N \gamma_k^{(n)} (\mathbf{x}^{(n)} - \mu_k)(\mathbf{x}^{(n)} - \mu_k)^T \\ \pi_k &= \frac{N_k}{N} \quad \text{with} \quad N_k = \sum_{n=1}^N \gamma_k^{(n)}\end{aligned}$$

# EM Algorithm for GMM

- **Initialize** the means  $\mu_k$ , covariances  $\Sigma_k$  and mixing coefficients  $\pi_k$
- Iterate until convergence:

**E-step:** Evaluate the responsibilities given current parameters







$$\gamma_k^{(n)} = p(z^{(n)} | \mathbf{x}) = \frac{\pi_k \mathcal{N}(\mathbf{x}^{(n)} | \mu_k, \Sigma_k)}{\sum_{j=1}^K \pi_j \mathcal{N}(\mathbf{x}^{(n)} | \mu_j, \Sigma_j)}$$

**M-step:** Re-estimate the parameters given current responsibilities

$$\begin{aligned}\mu_k &= \frac{1}{N_k} \sum_{n=1}^N \gamma_k^{(n)} \mathbf{x}^{(n)} \\ \Sigma_k &= \frac{1}{N_k} \sum_{n=1}^N \gamma_k^{(n)} (\mathbf{x}^{(n)} - \mu_k)(\mathbf{x}^{(n)} - \mu_k)^T \\ \pi_k &= \frac{N_k}{N} \quad \text{with} \quad N_k = \sum_{n=1}^N \gamma_k^{(n)}\end{aligned}$$

Evaluate log likelihood and check for convergence

$$\ln p(\mathbf{X} | \pi, \mu, \Sigma) = \sum_{n=1}^N \ln \left( \sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}^{(n)} | \mu_k, \Sigma_k) \right)$$



# Mixture of Gaussians vs. K-means

- EM for mixtures of Gaussians is just like a soft version of K-means, with **fixed priors and covariance**
- Instead of hard assignments in the E-step, we do **soft assignments** based on the softmax of the squared Mahalanobis distance from each point to each cluster.
- Each center moved by **weighted means** of the data, with weights given by soft assignments
- In K-means, weights are 0 or 1

# EM alternative approach \*

- Our goal is to maximize

$$p(\mathbf{X}|\Theta) = \sum_{\mathbf{z}} p(\mathbf{X}, \mathbf{z}|\Theta)$$

- Typically optimizing  $p(\mathbf{X}|\Theta)$  is difficult, but  $p(\mathbf{X}, \mathbf{Z}|\Theta)$  is easy
- Let  $q(\mathbf{Z})$  be a distribution over the latent variables. For any distribution  $q(\mathbf{Z})$  we have

$$\ln p(\mathbf{X}|\Theta) = \mathcal{L}(q, \Theta) + KL(q||p(\mathbf{Z}|\mathbf{X}, \Theta))$$


where

$$\begin{aligned}\mathcal{L}(q, \Theta) &= \sum_{\mathbf{z}} q(\mathbf{Z}) \ln \left\{ \frac{p(\mathbf{X}, \mathbf{Z}|\Theta)}{q(\mathbf{Z})} \right\} \\ KL(q||p) &= - \sum_{\mathbf{z}} q(\mathbf{Z}) \ln \left\{ \frac{p(\mathbf{Z}|\mathbf{X}, \Theta)}{q(\mathbf{Z})} \right\}\end{aligned}$$

# EM alternative approach \*


- The KL-divergence is always positive and have value 0 only if  $q(Z) = p(\mathbf{Z}|\mathbf{X}, \Theta)$
- Thus  $L(q, \Theta)$  is a lower bound on the likelihood

$$L(q, \Theta) \leq \ln p(\mathbf{X}|\Theta)$$






# Visualization of E-step



- The  $q$  distribution equal to the posterior distribution for the current parameter values  $\theta^{\text{old}}$ , causing the lower bound to move up to the same value as the log likelihood function, with the KL divergence vanishing.

# Visualization of M-step



- The distribution  $q(\mathbf{Z})$  is held fixed and the lower bound  $L(q, \Theta)$  is maximized with respect to the parameter vector  $\Theta$  to give a revised value  $\Theta^{new}$ . Because the KL divergence is nonnegative, this causes the log likelihood  $\ln p(\mathbf{X}|\Theta)$  to increase by at least as much as the lower bound does.

# E-step and M-step \*

$$\ln p(\mathbf{X}|\Theta) = \mathcal{L}(q, \Theta) + KL(q||p(\mathbf{Z}|\mathbf{X}, \Theta))$$

- In the E-step we maximize  $q(\mathbf{Z})$  w.r.t the lower bound  $L(q;_{old})$
- This is achieved when  $q(\mathbf{Z}) = p(\mathbf{Z}|\mathbf{X}, \Theta^{old})$
- The lower bound  $L$  is then

$$\begin{aligned}\mathcal{L}(q, \Theta) &= \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \Theta^{old}) \ln p(\mathbf{X}, \mathbf{Z}|\Theta) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \Theta^{old}) \ln p(\mathbf{Z}|\mathbf{X}, \Theta^{old}) \\ &= Q(\Theta, \Theta^{old}) + \text{const}\end{aligned}$$

- with the content the entropy of the  $q$  distribution, which is independent of  $\Theta$
- In the M-step the quantity to be maximized is the expectation of the complete data log-likelihood
- Note that  $\Theta$  is only inside the logarithm and optimizing the complete data likelihood is easier

# GMM Recap

- A probabilistic view of clustering - Each cluster corresponds to a different Gaussian.
- Model using **latent variables**.
- General approach, can replace Gaussian with other distributions (continuous or discrete)
- More generally, mixture model are very powerful models, **universal approximator**
- Optimization is done using the **EM** algorithm.

# EM Recap

- A general algorithm for optimizing many latent variable models.
- Iteratively computes a lower bound then optimizes it.
- Converges but maybe to a local minima.
- Can use multiple restarts.
- Can initialize from k-means
- Limitation - need to be able to compute  $P(z | \mathbf{x}; \Theta)$ , not possible for more complicated models.

# Acknowledges

---

Slides from:

- CSC 411 Fall 2018 Machine Learning and Data Mining, Toronto University
- David Sontag, New York University

# Thank you!

权小军 中山大学数据科学与计算机学院