

| | | Frequency of Inadequacy | | | | | | | | | | | Intensity of Inadequacy | | | | | | | | | | | | |
|--------|---------------------|---------------------------------|-------|---------|--------------------|---------|-------|---------------------------------|---------|-------|--------------------|-------|-------------------------|--------|---------------------------------|--------|---------|--------------------|--------|---------|---------------------------------|---------|--------|--------------------|--|
| | | Reported Individual Consumption | | | | | | Allocated Household Consumption | | | | | | | Reported Individual Consumption | | | | | | Allocated Household Consumption | | | | |
| | | wrt Age Group Req | | | wrt Individual Req | | | wrt Age Group Req | | | wrt Individual Req | | | | wrt Age Group Req | | | wrt Individual Req | | | wrt Age Group Req | | | wrt Individual Req | |
| Sex | Age Range | BMI | India | FAO/WHO | IOM | FAO/WHO | IOM | India | FAO/WHO | IOM | FAO/WHO | IOM | BMI | India | FAO/WHO | IOM | FAO/WHO | IOM | India | FAO/WHO | IOM | FAO/WHO | IOM | | |
| Female | infant (2-3) | NA | 89.1% | 80.3% | 88.7% | 78.7% | 81.4% | 80.7% | 86.4% | 88.1% | 74.5% | 54.0% | NA | -54.3% | -46.7% | -51.2% | -43.3% | -46.8% | -31.0% | -30.6% | -35.1% | -26.1% | -22.9% | | |
| Male | infant (2-3) | NA | 92.7% | 84.3% | 90.8% | 82.5% | 86.9% | 73.9% | 81.4% | 87.5% | 71.6% | 53.2% | NA | -53.4% | -48.1% | -51.1% | -45.6% | -46.7% | -31.4% | -32.0% | -34.5% | -27.9% | -23.9% | | |
| Female | child (3-10) | NA | 87.6% | 75.3% | 80.7% | 58.9% | 69.1% | 71.5% | 85.5% | 87.6% | 71.3% | 55.0% | NA | -38.5% | -34.6% | -36.1% | -28.4% | -31.3% | -27.7% | -31.9% | -33.9% | -26.9% | -22.7% | | |
| Male | child (3-10) | NA | 85.9% | 79.4% | 83.0% | 64.1% | 70.4% | 71.0% | 85.1% | 87.5% | 73.2% | 56.5% | NA | -37.5% | -35.5% | -37.2% | -29.9% | -31.7% | -27.9% | -32.7% | -34.5% | -27.0% | -22.4% | | |
| Female | adolescent (10-18) | NA | 88.8% | 80.4% | 73.8% | 67.7% | 52.6% | 85.8% | 83.7% | 84.0% | 69.4% | 46.7% | NA | -37.5% | -32.5% | -29.7% | -27.9% | -25.0% | -35.4% | -32.2% | -32.8% | -27.0% | -22.9% | | |
| Male | adolescent (10-18) | NA | 90.7% | 86.1% | 88.3% | 71.6% | 58.1% | 84.6% | 86.3% | 87.4% | 71.2% | 49.6% | NA | -39.6% | -37.1% | -37.7% | -29.3% | -26.2% | -33.8% | -32.5% | -33.8% | -26.8% | -21.2% | | |
| Female | young adult (18-30) | 67.9% | 82.7% | 80.7% | 83.6% | 71.0% | 72.0% | 69.0% | 87.3% | 86.6% | 70.9% | 52.7% | -14.7% | -37.5% | -35.6% | -36.0% | -32.0% | -31.7% | -31.1% | -35.7% | -34.8% | -27.4% | -22.7% | | |
| Male | young adult (18-30) | 76.3% | 82.8% | 83.7% | 87.6% | 75.4% | 30.5% | 85.7% | 85.3% | 87.2% | 73.1% | 46.9% | -13.7% | -37.6% | -37.3% | -39.4% | -31.6% | -24.6% | -39.9% | -32.5% | -34.9% | -27.8% | -20.2% | | |
| Female | adult (30-60) | 59.2% | 84.5% | 81.0% | 81.7% | 75.7% | 68.5% | 86.9% | 85.9% | 86.5% | 70.9% | 46.9% | -15.6% | -37.3% | -34.3% | -34.0% | -32.2% | -30.3% | -41.6% | -33.1% | -33.7% | -26.8% | -21.7% | | |
| Male | adult (30-60) | 69.1% | 87.8% | 85.0% | 88.1% | 78.6% | 20.3% | 95.1% | 85.3% | 87.2% | 72.7% | 47.5% | -13.9% | -39.4% | -37.2% | -38.0% | -33.3% | -23.6% | -46.1% | -31.8% | -33.6% | -27.0% | -20.9% | | |
| Female | senior (60+) | 74.8% | 86.3% | 73.0% | 77.0% | 59.5% | 50.0% | 97.9% | 83.3% | 85.1% | 68.7% | 45.0% | -20.5% | -36.9% | -32.7% | -32.6% | -29.0% | -26.8% | -51.6% | -30.8% | -32.9% | -25.2% | -20.6% | | |
| Male | senior (60+) | 76.7% | 92.9% | 82.9% | 89.8% | 64.9% | 13.9% | 97.3% | 83.4% | 86.0% | 65.3% | 34.8% | -18.3% | -42.3% | -32.9% | -37.9% | -28.2% | -22.0% | -48.1% | -30.4% | -33.3% | -24.0% | -19.1% | | |

Note:

Weights generated using IFPRI sample weights.

Consumption share is equal to individual Adult Equivalents/Total Household Adult Equivalents.

India is

WHO/FAO

IOM