The Asymmetric Metric-Tensor and the Thermodynamic Equation of State

Binyamin Tsadik Bair-Mosheh

September 19, 2026

Abstract

This document resumes Einstein's effort to unify gravitation and electromagnetism by generalizing the classical symmetric metric into a complete Asymmetric Metric-Tensor ($\hat{U}_{\mu\nu}$). By relaxing the assumption of an irrotational vacuum baseline, the tensor forward-models both macroscopic spacetime curvature (General Relativity) and discrete quantum kinematics (Electromagnetism and Mass) as orthogonal, coupled strain components of a continuous fluid manifold.

The scope of this manuscript constructs the primary tensor, computes its principal eigenvalues, and applies symmetry-breaking Lie derivatives to resolve the trace contractions of the matrix. These operations derive the four fundamental forces as boundary conditions of a single continuous pressure gradient. Beyond recovering the admittance-modified Navier-Stokes equation, the Kinematic Lagrangian and the Schwarzschild metric, the tensor extracts the cosmological constant (Λ), the physical Planck mass yield limit, the Compton cavitation boundary, anomalous galactic kinematics without dark matter, and cosmological redshift via thermodynamic metric capacity.

The capability of a single tensor to reproduce these scale invariant phenomena validates the asymmetric continuous baseline. Advanced phenomenological applications emerge as predictive corollaries. By formalizing the operational methodology of these extensions, the tensor establishes itself as a unified forward-modeling engine for continuous physics, validating Einstein's historical intuition that physical unification requires an asymmetric geometric extension.

1. Kinematic Parametrization (Clebsch Variables)

A continuous metric governed by a scalar potential (dh) is irrotational ($d(dh) = 0$). In order to capture the dynamics of topological defects and rotational shear, we first parameterize the macroscopic velocity field as a 1-form ($\mathbf{v}$) by using Clebsch variables [11].

Spatial vorticity is described by using a continuous velocity 1-form via orthogonal conjugate components:

$$\mathbf{v}(\mathbf{x}, t) = dh + \mathbf{v}_\perp + \lambda_\Omega d\beta_\Omega \tag{1}$$

Where dh represents the irrotational and longitudinal phase (compression and translation), whereas $\mathbf{v}_\perp$ represents propagating transverse shear, and $\lambda_\Omega d\beta_\Omega$ represents the internal rotational phase (topological twisting). The exterior derivative (d) is applied with respect to the flat Euclidean background (δ_{ij} in $\mathbb{R}^3$).

Taking the exterior derivative of the rotational velocity field extracts the internal topological vorticity 2-form ($\mathbf{\Omega}$):

$$\begin{aligned} \mathbf{\Omega} &= d(\lambda_\Omega d\beta_\Omega) \\ \mathbf{\Omega} &= d\lambda_\Omega \wedge d\beta_\Omega \end{aligned} \tag{2}$$

This parametrization represents the kinematic behavior of the metric: the irrotational phase represents longitudinal pressure, while the Clebsch wedge product represents spatial vorticity and the shear stress required for invariant rest mass [51].

2. The Thermodynamic Budget (Bernoulli Conservation)

Integrating the steady-flow Euler equation for an incompressible fluid ($\int \frac{dP}{\rho} + \mathbf{v} \cdot d\mathbf{v} = 0$) along a streamline to the phase limit (c) defines the finite, background yield-stress. This maximum limit establishes the Critical Pressure ($P_c = \frac{1}{2}\rho_\tau c^2$). By the conservation of energy (Bernoulli's Principle) [31], the sum of static pressure (P_{static}) and all applied kinematic loads equates to the unperturbed limit.

Parsing the two scale domains of structural load:

- **Macroscopic Baseline Load (ΔP_{macro}):** The depletion of ambient capacity induced via cosmic expansion and gravitational bodies. Integrating the macroscopic Hubble parameter ($v = H_0 r$) to the causal boundary determines the velocity field ($v_0(r) = c(1 - e^{-r/R_h})$). This background pressure yields $\Delta P_{macro} = \frac{1}{2}\rho_\tau v_0^2$, establishing an equation of state dictating maximum metric compliance for an observer at the origin ($r = 0, v_0 = 0$) and minimum compliance at the causal horizon ($v_0 \to c$) [2] (derived in Section 15).
- **Microscopic Load ($P_{dyn} + P_{shear}$):** The kinetic translation and structural vorticity of a boundary.

The Nested Ledger:
The ambient capacity of a frame, pre-strained by the background, establishes the Ambient Pressure ($P_{ambient}$):

$$P_{ambient} = P_c - \Delta P_{macro} = P_c - \frac{1}{2}\rho_\tau v_0^2 \tag{3}$$

This macroscopic pressure deficit translates into the Gravitational Compliance (α_g):

$$\alpha_g^2 = \frac{P_{ambient}}{P_c} = 1 - \frac{v_0^2}{c^2} \implies P_{ambient} = P_c \alpha_g^2 \tag{4}$$

According to the Bernoulli Principle, kinematics consume the available ambient capacity. Dynamic pressure (P_{dyn}) is quantified by taking the root-sum-square of orthogonal velocities, relative to the background density (ρ_τ):

$$P_{dyn} = \frac{1}{2}\rho_\tau \left(v_\parallel^2 + v_\perp^2 + v_{\circlearrowleft}^2 \right) \tag{5}$$

The distortional shear stress (P_{shear}) is governed by the magnitude of the Clebsch vorticity ($\mathbf{\Omega}$).

Subtracting this load establishes the remaining available static pressure (P_{static}), which acts as a structural ledger for the remaining capacity:

$$P_{static} = P_{ambient} - (P_{dyn} + P_{shear}) \tag{6}$$

The Ambient Compliance (α_a) quantifies the remaining capacity available:

$$\alpha_a^2 = \frac{P_{static}}{P_{ambient}} \tag{7}$$

The Total Scalar Admittance (α_s^2) of the coordinate is determined by taking the product of the nested compliances:

$$\alpha_s^2 \equiv \frac{P_{static}}{P_c} = \left(\frac{P_{ambient}}{P_c} \right) \left(\frac{P_{static}}{P_{ambient}} \right) = \alpha_g^2 \alpha_a^2 \tag{8}$$

3. Assembly of the Asymmetric Metric-Tensor ($\hat{U}_{\mu\nu}$)

The corresponding thermodynamic loads hold representation in metric geometry; the parameters map to a dimensionless matrix. An asymmetric second-rank tensor decomposes into a symmetric component (volumetric strain) and an antisymmetric component (deviatoric shear strain). This geometric summation is structurally isomorphic to the Clifford geometric product ($uv = u \cdot v + u \wedge v$), formulating spacetime as a continuous multivector and defining the metric as a micropolar Cosserat continuum [13, 23, 16]:

$$\hat{U}_{\mu\nu} = \hat{U}_{(\mu\nu)} + \hat{U}_{[\mu\nu]} \equiv S_{\mu\nu} + A_{\mu\nu} \tag{9}$$

To satisfy covariance, the metric is constructed via the continuous 4-velocity (u_μ) and the spatial projection tensor ($h_{\mu\nu}$). The symmetric sector ($S_{\mu\nu}$) and the antisymmetric sector ($A_{\mu\nu}$) split as:

$$\hat{U}_{\mu\nu} = \underbrace{-\alpha_s^2 u_\mu u_\nu + \frac{1}{\alpha_s^2} h_{\mu\nu}}_{S_{\mu\nu}} + \underbrace{\alpha \left(\frac{u_\mu v_{\perp,\nu} - u_\nu v_{\perp,\mu}}{c} \right) + \alpha(t_p)\omega_{\mu\nu}}_{A_{\mu\nu}} \tag{10}$$

Where the covariant tensor variables are:

- u_μ: **Continuum 4-Velocity.** The normalized, timelike vector field ($u_\mu u^\mu = -1$) representing the local rest frame of the coordinate metric. In this comoving frame, the spatial components evaluate to zero ($u^i = 0$) and the temporal component evaluates to $u^0 = 1$, isolating bulk geometric strain to the symmetric diagonals.
- $h_{\mu\nu}$: **Spatial Projection Tensor.** The symmetric tensor ($h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$) that projects the metric onto the orthogonal spatial hypersurface, where $g_{\mu\nu}$ is the unperturbed Minkowski background signature $(-, +, +, +)$.
- α_s: **Total Scalar Admittance.** The dimensionless phase capacity bounded by thermodynamic saturation, representing the scalar contraction of the symmetric sector.
- α: **Viscoelastic Modulus Ratio.** The dimensionless transverse coupling constant. Tracks the fractional capacity of the metric to sustain transverse shear relative to bulk compression.
- $v_{\perp,\mu}$: **Transverse Slip Vector.** The orthogonal spatial velocity representing solenoidal slip, coupling to the time-space components via the antisymmetric wedge product.
- $\omega_{\mu\nu}$: **Vorticity Tensor.** The antisymmetric component representing the curl of the Clebsch rotational phase.
- t_p: **Structural Relaxation Time.** The temporal phase limit ($t_p = l_p/c$) that normalizes the vorticity into dimensionless strain.

3.1 The Symmetric Diagonals (Volumetric Strain)

The symmetric components govern the bulk geometric packing, driven by the ambient ratio.

- **Temporal Component ($\hat{U}_{00}$):** Phase update capacity. It resolves to the negative dimensionless static pressure ($\hat{U}_{00} = -\alpha_g^2 \alpha_a^2$). The time-time component is the scalar product of gravitational and kinematic time dilation.

- **Spatial Components ($\hat{U}_{ii}$):** Effective volumetric density and spatial packing of the coordinate grid. Maps to the inverse dimensionless static pressure: $\hat{U}_{ii} = \frac{1}{\alpha_g^2 \alpha_a^2}$.

3.2 The Antisymmetric Off-Diagonals (Transverse Shear)

The antisymmetric components capture the dynamic sliding and twisting of the metric. Translating these physical observables into a dimensionless strain matrix is achieved via normalization:

- **Viscoelastic Throttling (α):** The metric capacity to sustain transverse shear acts as a geometric fraction of its bulk compressive capacity. The off-diagonals are attenuated by the Viscoelastic Modulus Ratio (α).
- **Transverse Velocity ($\hat{U}_{0i}$):** The temporal rate of transverse displacement ($v_{\perp,i}$). Normalized by the acoustic phase limit (c), this defines the continuous strain of the Electric Field: $\hat{U}_{0i} = -\alpha\left(\frac{v_{\perp,i}}{c}\right)$.
- **Rotational Vorticity ($\hat{U}_{ij}$):** The exterior derivative of the metric's rotational phase ($d\lambda_\Omega \wedge d\beta_\Omega$). Because vorticity is a physical frequency (Ω_k), it is scaled by the structural relaxation time of the metric ($t_p \equiv l_p/c$) to yield a dimensionless strain. This determines the dimensionless Deborah Number of the metric, generating the Magnetic Field: $\hat{U}_{ij} = \alpha(t_p\Omega_k)$.

3.3 The Final Tensor Matrix

Superimposing the symmetric and antisymmetric components yields the 4×4 Asymmetric Metric-Tensor for a Cartesian coordinate referenced against the unperturbed vacuum (c):

$$\hat{U}_{\mu\nu} = \begin{bmatrix} -\alpha_s^2 & -\alpha\left(\frac{v_{\perp,x}}{c}\right) & -\alpha\left(\frac{v_{\perp,y}}{c}\right) & -\alpha\left(\frac{v_{\perp,z}}{c}\right) \\ \alpha\left(\frac{v_{\perp,x}}{c}\right) & \frac{1}{\alpha_s^2} & -\alpha(t_p\Omega_z) & \alpha(t_p\Omega_y) \\ \alpha\left(\frac{v_{\perp,y}}{c}\right) & \alpha(t_p\Omega_z) & \frac{1}{\alpha_s^2} & -\alpha(t_p\Omega_x) \\ \alpha\left(\frac{v_{\perp,z}}{c}\right) & -\alpha(t_p\Omega_y) & \alpha(t_p\Omega_x) & \frac{1}{\alpha_s^2} \end{bmatrix} \tag{11}$$

Where the Cartesian matrix variables are specified as:

- $v_{\perp,i}/c$: **Dimensionless Transverse Slip.** The temporal rate of spatial sliding along the i-axis, normalized by the acoustic phase limit. Generates the electric field vector (E_i).
- $t_p\Omega_i$: **Dimensionless Vorticity.** The spatial curl of the Clebsch phase around the i-axis, normalized by the structural relaxation limit. Generates the magnetic field vector (B_i).

This architecture enables cross-coupling. As gravity, velocities, or internal vorticities increase, the scalar capacity (α_s^2) approaches zero. The spatial diagonals expand, generating a dense boundary that limits the metric's capacity to sustain transverse waves in the off-diagonals (enforcing the coordinate limits of the speed of light).

4. The Continuous Field Equation

Because $\hat{U}_{\mu\nu}$ represents dimensionless strain, it balances against the physical stress that caused the deformation. Applying generalized Hooke's Law, the dimensionless tensor is multiplied by the thermodynamic stiffness of the baseline vacuum (P_c).

Macroscopic field propagation is governed by the global background. The resulting field equates to the continuous asymmetric Cauchy stress tensor ($\hat{T}_{\mu\nu}$) of the overarching metric. Topological defects emerge as steady-state circulation structures embedded within this continuous medium.

The Unified Field Equation is defined as the continuum's stress deviation from the flat Minkowski baseline ($\eta_{\mu\nu}$):

$$\hat{T}_{\mu\nu} = P_c\Big(\hat{U}_{\mu\nu} - \eta_{\mu\nu}\Big) \tag{12}$$

To maintain covariance near topological boundaries where dynamic pressure approaches the baseline capacity ($P_{dyn} \to P_c$), this linear relation transitions to a non-linear hyperelastic

extension. To evaluate this limit, we define the second invariant constraint (I_2) not as the trace, but as the full scalar contraction of the strain tensor ($I_2 = \hat{U}_{\alpha\beta}\hat{U}^{\alpha\beta}$).

At the unperturbed Minkowski baseline ($\eta_{\mu\nu}$), this invariant evaluates to 4:

$$I_2^{(vac)} = \eta_{\mu\nu}\eta^{\mu\nu} = (-1)(-1) + (1)(1) + (1)(1) + (1)(1) = 4 \tag{13}$$

This invariant bounds the stress tensor to prevent infinite deformation at the boundary:

$$\hat{T}_{\mu\nu} = P_c \left(\hat{U}_{\mu\nu} - \eta_{\mu\nu} \right) + 2C_1 \left(I_2 - 4 \right) \hat{U}_{\mu\nu} \tag{14}$$

Because weak-field coordinates operate far below the saturation threshold ($I_2 \approx 4$), the hyperelastic coefficient (C_1, possessing units of thermodynamic pressure) term attenuates, recovering the linear baseline for standard field interactions.

4.1 Energy Density and Structural Stress

Isolating the temporal (00) slot computes the continuous equivalent of energy density. Given $\eta_{00} = -1$:

$$\begin{aligned} \hat{T}_{00} &= P_c \left(-\frac{P_{static}}{P_c} - (-1) \right) \\ \hat{T}_{00} &= P_c - P_{static} \\ \hat{T}_{00} &= \Delta P_{macro} + P_{dyn} + P_{shear} \end{aligned} \tag{15}$$

Energy density ($\hat{T}_{00}$) reduces to the sum of the macroscopic gravitational deficit and the internal kinematic pressure of the mass defect.

Evaluating the individual spatial components (no sum) determines the continuous structural stress. Given $\eta_{ii} = 1$:

$$\hat{T}_{ii} = P_c \left(\frac{P_c - P_{static}}{P_{static}} \right) = P_c \left(\frac{\Delta P_{macro} + P_{dyn} + P_{shear}}{P_{static}} \right) \tag{16}$$

The physical coordinate stress maps to the total applied pressure load scaled by the inverse of the available static capacity. As structural loads saturate the metric ($P_{static} \to 0$), the physical spatial stress diverges ($\hat{T}_{ii} \to \infty$), establishing the cavitation limit without a singularity.

4.2 The Cosmological Constant & Metric Density (ρ_τ)

Because the Unified Field Equation (12) evaluates the deviatoric stress applied by kinematic defects, an unperturbed coordinate space ($\hat{U}_{\mu\nu} = \eta_{\mu\nu}$) evaluates to zero ($\hat{T}_{\mu\nu} = 0$). To derive the baseline metric density (ρ_τ), the background stress tensor is defined as ($\Sigma_{\mu\nu}^{(vac)} = -P_c\eta_{\mu\nu}$), which establishes the isotropic tension of the manifold. We evaluate the spatial surface integral of this vacuum stress across the minimum resolvable coordinate boundary (a sphere of Planck radius, $A_p = 4\pi l_p^2$).

By divergence equilibrium, the total outward spatial force exerted by the static vacuum stress (P_c) on this boundary computes to $F = P_c(4\pi l_p^2)$. Because rest energy is evaluated over one full phase cycle (derived formally in Section 14 as $E = mc^2$), this boundary equates to the tension limit of the manifold ($F_p = c^4/G$).

Equating these bounds sets the microscopic yield pressure:

$$P_c(4\pi l_p^2) = \frac{c^4}{G} \implies P_c = \frac{c^4}{4\pi G l_p^2} \tag{17}$$

Substituting the geometric definition of the Planck area ($l_p^2 = \hbar G/c^3$) defines the metric stiffness:

$$P_c = \frac{c^7}{4\pi\hbar G^2} \tag{18}$$

Equating this to dynamic pressure ($P_c = \frac{1}{2}\rho_\tau c^2$) extracts the baseline metric density:

$$\begin{aligned} \frac{1}{2}\rho_\tau c^2 &= \frac{c^7}{4\pi\hbar G^2} \\ \rho_\tau &= \frac{2P_c}{c^2} = \frac{1}{2\pi}\left(\frac{c^5}{\hbar G^2}\right) \end{aligned} \tag{19}$$

The density of the metric is fixed as $1/2\pi$ of the Planck density.

The Macroscopic Limit (The Epoch-Dependent Cosmological Parameter Λ): The continuous manifold is bounded by the causal horizon ($R_h = c{\cdot}t$). Because the total tension of the spacetime manifold (c^4/G) acts as an invariant geometric constant, the force projected across any continuous 3D isotropic spherical boundary remains conserved ($F = P \cdot A = c^4/G$).

At the microscopic limit, this isolates the Planck stiffness ($P_c \cdot 4\pi l_p^2 = c^4/G$). As the universe expands, this invariant tension distributes over the causal area ($4\pi R_h^2$). The observed macroscopic vacuum pressure (P_{macro}) is therefore the geometric dilution of the baseline metric stiffness:

$$P_{macro}(4\pi R_h^2) = P_c(4\pi l_p^2) = \frac{c^4}{G} \tag{20}$$

Isolating the macroscopic pressure yields:

$$P_{macro} = \frac{c^4/G}{4\pi R_h^2} = \frac{c^4}{4\pi G R_h^2} \tag{21}$$

Substituting this scaled macroscopic pressure into the definition of the Cosmological Scalar ($\Lambda = 8\pi G P_{macro}/c^4$) yields:

$$\begin{aligned} \Lambda &= \frac{8\pi G}{c^4}\left(\frac{c^4}{4\pi G R_h^2}\right) \\ \Lambda &= \frac{2}{R_h^2} \end{aligned} \tag{22}$$

Substituting the present-day Hubble radius ($R_h = c/H_0 \approx 1.37 \times 10^{26}$ m) calculates the local snapshot of the metric:

$$\begin{aligned} \Lambda &= \frac{2}{(1.37 \times 10^{26}\ \mathrm{m})^2} \\ \Lambda &= 1.06 \times 10^{-52}\ \mathrm{m}^{-2} \end{aligned} \tag{23}$$

This reproduces the Planck 2018 empirical measurement within 4%: ($\Lambda_{obs} \approx 1.1{\times}10^{-52}$ m^{-2}) [42].

The Poisson Limit:

Extracting the individual spatial components (no sum) of the continuous Cauchy stress tensor from the baseline field equation yields:

$$\hat{T}_{ii} = P_c\Big(\hat{U}_{ii} - \eta_{ii}\Big) \tag{24}$$

In the weak-field limit, the symmetric spatial diagonal acts as a linear perturbation of the volumetric packing. Because gravitational potential (Φ) represents a negative geometric energy

state ($P_{dyn} = -\rho_\tau \Phi$), the spatial strain expands inversely to the dropping scalar admittance ($\hat{U}_{ii} = 1/\alpha_s^2 \approx 1 - \frac{2\Phi}{c^2}$). Substituting this and the flat Minkowski baseline ($\eta_{ii} = 1$) isolates the continuous structural stress:

$$\hat{T}_{ii} = P_c \left(1 - \frac{2\Phi}{c^2} - 1\right)$$
$$\hat{T}_{ii} = P_c \left(-\frac{2\Phi}{c^2}\right) \tag{25}$$

Substituting the stiffness ($P_c = \frac{1}{2}\rho_\tau c^2$) eliminates the phase limits. This defines the restorative mechanical stress holding the mass defect in place. Because the potential is negative ($\Phi < 0$), the tensor generates a positive tensile stress with a negative spatial gradient:

$$\hat{T}_{ii} = \left(\frac{1}{2}\rho_\tau c^2\right)\left(-\frac{2\Phi}{c^2}\right)$$
$$\hat{T}_{ii} = -\rho_\tau \Phi \tag{26}$$

4.3 The Macroscopic Reduction (The Einstein Field Equation)

The classical formalism of General Relativity resolves as a macroscopic limit of the Unified Field Equation.

1. The Macroscopic Boundary Constraint:
At the cosmological scale, the restoring thermodynamic stiffness of the continuum is governed by the scaled macroscopic vacuum pressure (P_{macro}). Applying this boundary constraint to the Unified Field Equation limits the macroscopic Cauchy stress ($\hat{T}_{\mu\nu}^{macro}$):

$$\hat{T}_{\mu\nu}^{macro} = P_{macro}\left(\hat{U}_{\mu\nu} - \eta_{\mu\nu}\right) \tag{27}$$

2. The Defect Density and Thermodynamic Substitution:
As derived in Section 4.2, the restoring macroscopic pressure is parameterized by the expanding causal horizon ($P_{macro} = \frac{c^4}{4\pi G R_h^2}$). Macroscopic mass-energy introduces topological defects, mapping the defect density to the classical Stress-Energy Tensor ($T_{\mu\nu}$). Standard GR assumes a symmetric stress-energy tensor because macroscopic objects possess negligible net spin density.

3. The Volumetric Determinant (Jacobian Equivalence):
Because the metric trace ($I_1 \equiv \hat{U}_\mu^\mu$) governs volumetric strain, the scalar admittance (α_s^2) is equivalent to the normalized volume (the Jacobian determinant) of the local spacetime metric:

$$\alpha_s^2 \equiv \frac{\sqrt{-g}}{\sqrt{-g_0}} \equiv J_{normalized} \tag{28}$$

Where $\sqrt{-g_0}$ represents the unperturbed vacuum baseline. This identity formalizes the 1-to-1 mapping between the permittivity of the vacuum (α_s^2) and the volumetric compression of the manifold ($J_{normalized}$).

4. The Differential Incompatibility Mapping (The Kröner Formulation):
To map the zero-order asymmetric strain ($\hat{U}_{\mu\nu}$) to the second-order curvature tensor ($G_{\mu\nu}$) without manually stripping the antisymmetric components ($A_{\mu\nu}$), we apply the mathematics of continuum defect theory.

Macroscopic mass acts as a Volterra dislocation [52] within the continuous metric. The geometric consequence of the continuum warping to accommodate these spatial defects is measured

by the Saint-Venant incompatibility tensor (Inc) [45, 30]. We apply this continuous differential operator to the covariant strain tensor:

$$G_{\mu\nu} \equiv \text{Inc}(\hat{U}_{\mu\nu}) \approx \partial_\alpha \partial_\beta (\hat{U}_{\mu\nu} - \eta_{\mu\nu}) \epsilon^{\alpha\mu\rho} \epsilon^{\beta\nu\sigma} \eta_{\rho\sigma} \tag{29}$$

5. Torsion Nullification at the Macroscopic Limit (Coarse-Graining):
Because the strain tensor contains both symmetric volumetric strain ($S_{\mu\nu}$) and antisymmetric transverse shear ($A_{\mu\nu}$), the incompatibility operator inherently evaluates both. To prove that the antisymmetric sector decouples from the macroscopic large-scale curvature, we apply the mathematical formalism of spatial coarse-graining over a macroscopic volume V where $V \gg l_p^3$.

The macroscopic observable strain is defined as the volume average of the microscopic fields:

$$\langle \hat{U}_{\mu\nu} \rangle_V = \frac{1}{V} \int_V (S_{\mu\nu} + A_{\mu\nu})\, d^3x \tag{30}$$

In unpolarized mass distributions, transverse shear (quantum vorticity/spin and electromagnetic dipoles) are isotropically distributed. Their phase orientations are randomized, meaning their static vector sum across the macroscopic volume V statistically cancels:

$$\frac{1}{V} \int_V A_{\mu\nu}^{static} d^3x = 0 \implies \langle A_{\mu\nu}^{static} \rangle_V = 0 \tag{31}$$

Conversely, propagating electromagnetic waves exist as zero-mean oscillatory perturbations ($\Box A_{\mu\nu}^{dynamic} = 0$). While their kinematic energy propagates through the metric, their linear spatial average across macroscopic volumes evaluates to zero.

Because the Saint-Venant incompatibility tensor (Inc) is a linear differential operator with respect to strain, the incompatibility of the macroscopic average is equal to the average of the microscopic incompatibility:

$$\text{Inc}(\langle A_{\mu\nu} \rangle_V) = \langle \text{Inc}(A_{\mu\nu}) \rangle_V = 0 \tag{32}$$

Furthermore, this cancellation aligns with the established constraints of Einstein-Cartan-Sciama-Kibble (ECSK) theory [25, 46]. In ECSK spacetime, the antisymmetric torsion tensor is coupled to the spin density of matter. Unlike symmetric curvature, torsion is non-propagating; it vanishes outside the defect.

The smoothing of quantum properties ($\text{Inc}(A_{\mu\nu}) \to 0$) does not apply to either a locally polarized dynamic or to macroscopic asymmetry ($\langle A_{\mu\nu} \rangle_V \neq 0$) such as cosmological torsion events.

However, General Relativity operates under the physical reality that most of the bulk mass distributions are randomly distributed. Under this assumption, the surviving equation relies solely on the symmetric incompatibility ($G_{\mu\nu} \approx \text{Inc}(S_{\mu\nu})$) to generate classical large-scale gravitational curvature, preserving the continuous vacuum's capacity to propagate zero-mean transverse waves.

6. The Classical Limit and Scale Invariance:
As formalized by Kröner and extended to gravitational geometry by Katanaev and Volovich [24], the structural incompatibility of metric strain is directly proportional to the defect density tensor. When mapping it to spacetime, this defect density tensor functions identically to the classical Stress-Energy Tensor ($T_{\mu\nu}$) [26]. Scaling this defect density by the thermodynamic stiffness limits of the vacuum ($8\pi G/c^4$) equates the geometric curvature to the physical stress-energy:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \tag{33}$$

The Einstein Field Equation resolves as the macroscopic, unpolarized compatibility condition of the full unified field. The antisymmetric components ($A_{\mu\nu}$) are never zeroed out or

removed from the baseline tensor ($\hat{U}_{\mu\nu}$). While their net curvature footprint averages to zero for randomly oriented bulk mass, they remain active across all scales. They govern quantum mechanics and electrodynamics, and when coherent, they scale proportionally to dictate macroscopic rotational kinematics at the galactic level, demonstrating the universal scale-invariance of the overarching field equations.

5. The Equation of Motion (Divergence Equilibrium)

Energy-momentum conservation emerges via the spacetime divergence of the stress tensor ($\nabla_\mu \hat{T}^{\mu\nu} = 0$). The covariant derivative incorporates the non-linear affine connection ($\Gamma^\lambda_{\mu\nu}$) constructed in Section 12. However, the antisymmetric torsion ($T^\lambda_{\mu\nu} = \Gamma^\lambda_{\mu\nu} - \Gamma^\lambda_{\nu\mu}$) decays at the boundaries ($r \gg r_c$), producing a symmetric, torsion-free connection. Under this weak-field limit, the spatial geometry flattens against the Minkowski background ($\eta_{\mu\nu}$), and the covariant divergence converges to the standard partial derivative ($\partial_\mu \hat{T}^{\mu\nu} \approx 0$).

To extract the equation of motion, the macroscopic momentum density ($\hat{T}^{0i}$) is derived from the covariant parameters of the metric tensor ($\hat{U}^{\mu\nu}$). Evaluating the temporal-spatial component of Equation (10) with the normalized 4-velocity ($u^\mu = \gamma(1, v^i/c)$) and the spatial projection tensor ($h^{0i} = \eta^{0i} + u^0 u^i = \gamma^2 v^i/c$) isolates the symmetric field strain:

$$S^{0i} = -\alpha_s^2 u^0 u^i + \frac{1}{\alpha_s^2} h^{0i} = -\gamma^{-2}\left(\gamma^2 \frac{v^i}{c}\right) + \gamma^2\left(\gamma^2 \frac{v^i}{c}\right) = \frac{v^i}{c}(\gamma^4 - 1) \tag{34}$$

Applying the weak-field expansion ($\gamma^4 - 1 \approx 2v^2/c^2$) evaluates the strain. Multiplying by the baseline stiffness ($P_c = \frac{1}{2}\rho_\tau c^2$) translates this strain into the Cauchy stress momentum density ($\hat{T}^{0i} = P_c S^{0i}$):

$$\hat{T}^{0i} = \left(\frac{1}{2}\rho_\tau c^2\right)\frac{v^i}{c}\left(\frac{2v^2}{c^2}\right) = 2\left(\frac{1}{2}\rho_\tau \frac{v^2}{c^2}\right) c v^i \tag{35}$$

Defining density as the mass equivalent of the localized pressure ($\rho_{dyn} = P_{dyn}/c^2 = \frac{1}{2}\rho_\tau v^2/c^2$), the tensor outputs the Eulerian momentum density flux ($\hat{T}^{0i} = 2c\rho_{dyn} v^i$).

The symmetric spatial components (S^{ij}) are calculated using the weak-field Taylor expansion ($\gamma^2 \approx 1 + v^2/c^2$), producing the spatial Cauchy stress ($\hat{T}^{ij}$):

$$S^{ij} = -\alpha_s^2 u^i u^j + \frac{1}{\alpha_s^2} h^{ij} = -\gamma^{-2}\left(\gamma^2 \frac{v^i v^j}{c^2}\right) + \gamma^2\left(\delta^{ij} + \gamma^2 \frac{v^i v^j}{c^2}\right) \approx \left(1 + \frac{v^2}{c^2}\right)\delta^{ij} + 2\frac{v^2}{c^2}\frac{v^i v^j}{c^2} \tag{36}$$

Multiplying by the baseline stiffness translates this strain into the spatial Cauchy stress ($\hat{T}^{ij} = P_c(S^{ij} - \delta^{ij}) - \tau^{ij}$), dropping the higher-order velocity term ($\frac{v^4}{c^4} \approx 0$) in the weak field limit:

$$\hat{T}^{ij} \approx P_c\left(\frac{v^2}{c^2}\right)\delta^{ij} + P_c\left(2\frac{v^2}{c^2}\frac{v^i v^j}{c^2}\right) - \tau^{ij} = \left(\frac{1}{2}\rho_\tau v^2\right)\delta^{ij} + \left(\rho_\tau \frac{v^2}{c^2}\right) v^i v^j - \tau^{ij} \tag{37}$$

Because $P_{total} = \frac{1}{2}\rho_\tau v^2$, the tensor produces spatial flux:

$$\hat{T}^{ij} = P_{total}\delta^{ij} + 2\rho_{dyn} v^i v^j - \tau^{ij} \tag{38}$$

5.1 The Equation of Motion (Navier-Stokes)

To construct the equation of motion independent of scalar energy density variations, the 4D spacetime divergence is expanded across its temporal and spatial components ($\nabla_0 \hat{T}^{0i} + \nabla_j \hat{T}^{ji} = 0$). By applying the affine connection contraction ($\Gamma^\lambda_{\mu\lambda} = \partial_\mu \ln\sqrt{-g}$), the covariant momentum transport resolves across curved coordinates. In the previously established macroscopic weak-field limit, this expansion simplifies to the partial derivative domain.

Substituting the derived macroscopic stress components ($\hat{T}^{0i} = 2c\rho_{dyn}v^i$ and $\hat{T}^{ij} = P_{total}\delta^{ij} + 2\rho_{dyn}v^iv^j - \tau^{ij}$):

$$\frac{1}{c}\partial_t(2c\rho_{dyn}v^i) + \partial_j(P_{total}\delta^{ij} + 2\rho_{dyn}v^iv^j - \tau^{ij}) = 0 \tag{39}$$

Dividing the entire equilibrium equation by 2 isolates the dynamic density (ρ_{dyn}) relative to the structural pressure:

$$\partial_t(\rho_{dyn}v^i) + \partial_j\left(\frac{1}{2}P_{total}\delta^{ij} + \rho_{dyn}v^iv^j - \frac{1}{2}\tau^{ij}\right) = 0 \tag{40}$$

Applying the product rule to the convective momentum terms expands the temporal and spatial derivatives:

$$v^i\partial_t\rho_{dyn} + \rho_{dyn}\partial_t v^i + \frac{1}{2}\partial^i P_{total} + v^i\partial_j(\rho_{dyn}v^j) + \rho_{dyn}v^j\partial_j v^i - \frac{1}{2}\partial_j\tau^{ij} = 0 \tag{41}$$

Grouping the kinetic terms extracts the acceleration from the volumetric mass flux:

$$\rho_{dyn}(\partial_t v^i + v^j\partial_j v^i) + v^i\left[\partial_t\rho_{dyn} + \partial_j(\rho_{dyn}v^j)\right] = -\frac{1}{2}\partial^i P_{total} + \frac{1}{2}\partial_j\tau^{ij} \tag{42}$$

Assuming steady-state Eulerian mass conservation for the dynamic density envelope ($\partial_t\rho_{dyn} + \nabla\cdot(\rho_{dyn}\mathbf{v}) = 0$), the remaining kinetic term is the definition of the Eulerian Material Derivative ($\frac{D\mathbf{v}}{Dt} \equiv \partial_t\mathbf{v} + (\mathbf{v}\cdot\nabla)\mathbf{v}$), determining the coordinate acceleration of the coordinate frame:

$$\rho_{dyn}\frac{D\mathbf{v}}{Dt} = -\frac{1}{2}\nabla P_{total} + \frac{1}{2}\nabla\cdot\boldsymbol{\tau} \tag{43}$$

1. The Stress Tensor Partition:
The Cauchy stress on the right side of Equation (43) is driven by the components of the Tensor.

- **Symmetric Volumetric Stress ($-\frac{1}{2}\nabla P_{total}$):** The spatial diagonals ($\hat{U}_{ii}$) govern volumetric packing. A boundary condition upon the local frame identifies the total coordinate stress (P_{total}) as the superposition of the macroscopic force of the environment ($-\frac{1}{2}\nabla P_{ambient}$) and the compressive surface force of the defect ($-\frac{1}{2}\nabla P_{dyn}$).
- **Antisymmetric Deviatoric Shear ($\frac{1}{2}\nabla\cdot\boldsymbol{\tau}$):** The off-diagonals control transverse shear and the divergence of Clebsch vorticity ($\nabla\cdot\boldsymbol{\Omega}$). This transverse stress is attenuated by the Viscoelastic Modulus Ratio (α).

Substituting the partitioned stress components into the momentum equation yields:

$$\rho_{dyn}\frac{D\mathbf{v}}{Dt} = -\frac{1}{2}\nabla P_{ambient} - \frac{1}{2}\nabla P_{dyn} + \frac{1}{2}\alpha\big(\nabla\cdot\mathbb{T}_\perp - \mu\nabla\cdot(d\lambda_\Omega\wedge d\beta_\Omega)\big) \tag{44}$$

2. Kinematic Normalization:
To resolve the dimensional units from force density (N/m^3) into kinematic acceleration (m/s^2) the equation is divided by the baseline density (ρ_τ).

This defines the mass-independent kinematic pressure gradients and stresses (absorbing the geometric halving factor):

$$\mathcal{P}_{ambient} = \frac{P_{ambient}}{2\rho_\tau} \tag{45}$$

$$\mathcal{P}_{dyn} = \frac{P_{dyn}}{2\rho_\tau} \tag{46}$$

$$\mathbb{T}_{\perp,kin} = \frac{\mathbb{T}_\perp}{2\rho_\tau} \tag{47}$$

$$\nu = \frac{\mu}{2\rho_\tau} \tag{48}$$

Applying these invariant dimensional substitutions constructs the kinematic equation of motion:

$$\left(\frac{\rho_{dyn}}{\rho_\tau}\right)\frac{D\mathbf{v}}{Dt} = -\nabla\mathcal{P}_{ambient} - \nabla\mathcal{P}_{dyn} + \alpha\big(\nabla\cdot\mathbb{T}_{\perp,kin} - \nu\nabla\cdot(d\lambda_\Omega\wedge d\beta_\Omega)\big) \tag{49}$$

The divergence equilibrium conserves total angular momentum by transferring macroscopic deviatoric shear into the internal spin of the coordinate phase. The tensor establishes the Weak Equivalence Principle mechanically: macroscopic gravitational acceleration ($-\nabla\mathcal{P}_{ambient}$) operates independently of internal mass, while mechanical propulsion is bounded by the dynamic density limits of the coordinate frame.

5.2 The Archimedean Limit of the Geodesic Equation

A void submerged within a pressure gradient experiences a restorative buoyant force directed toward the lowest static pressure. Newtonian gravity emerges as macroscopic metric buoyancy directly from the geodesic equation of motion.

The equation of motion for a test coordinate in a continuous metric is governed by the geodesic equation ($\frac{d^2x^i}{d\tau^2} = -\Gamma^i_{\mu\nu}\frac{dx^\mu}{d\tau}\frac{dx^\nu}{d\tau}$). This derivation isolates the Newtonian limit by evaluating a static metric ($\partial_0\hat{U}_{\mu\nu} = 0$) and a non-relativistic test coordinate ($dx^i/d\tau \ll c$). While relativistic or time-varying geometries require the full non-linear evaluation of the metric connection, this boundary condition reduces the equation to the temporal Christoffel symbol ($a^i \approx -c^2\Gamma^i_{00}$).

For a static metric, the connection evaluates as $\Gamma^i_{00} \approx -\frac{1}{2}\eta^{ij}\partial_j\hat{U}_{00}$. Substituting this derivative determines the coordinate acceleration:

$$\mathbf{a} \approx \frac{1}{2}c^2\nabla\hat{U}_{00} \tag{50}$$

Equating the perturbation to the kinematic metric definition ($\hat{U}_{00} = -P_{static}/P_c$) limits the acceleration as a function of the static pressure gradient:

$$\begin{aligned}
\mathbf{a} &= \frac{1}{2}c^2\nabla\left(-\frac{P_{static}}{P_c}\right)\\
\mathbf{a} &= -\frac{c^2}{2\left(\frac{1}{2}\rho_\tau c^2\right)}\nabla P_{static}\\
\mathbf{a} &= -\frac{1}{\rho_\tau}\nabla P_{static}
\end{aligned} \tag{51}$$

The physical force ($\mathbf{F}_g$) acting upon a particle is derived by multiplying this coordinate acceleration against the particle's entrained hydrodynamic added mass (m_τ):

$$\mathbf{F}_g = m_\tau\mathbf{a} = -\left(\frac{m_\tau}{\rho_\tau}\right)\nabla P_{static} \tag{52}$$

The total phase capacity consumed within the continuous metric functions as an Effective Thermodynamic Displacement ($V_{eff} = m_\tau/\rho_\tau$). This effective volume represents the total hydrodynamic added mass rather than the physical volume of the cavitation. Substituting this effective displacement constructs the equation of state:

$$\mathbf{F}_g = -V_{eff}\nabla P_{static} \tag{53}$$

To derive the static pressure gradient without external kinematic postulates, the overarching metric must be evaluated as a continuous 3D spatial geometry. In a static continuous medium,

the macroscopic scalar strain generated by a spherically symmetric cavitation boundary resolves via the radial Laplace equation ($\nabla^2 P_{dyn} = 0$).

Expanding the Laplacian operator in spherical coordinates establishes the structural conservation of the spatial pressure gradient across the expanding coordinate area (r^2):

$$\frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2\frac{\partial P_{dyn}}{\partial r}\right) = 0 \tag{54}$$

Integrating the equation once extracts the strain gradient. The constant of integration ($-k$) represents the structural scalar magnitude of the defect:

$$\begin{aligned} r^2\frac{\partial P_{dyn}}{\partial r} &= -k \\ \nabla P_{dyn} &= -\frac{k}{r^2}\hat{\mathbf{r}} \end{aligned} \tag{55}$$

Integrating this gradient radially outward dictates the continuous scalar dynamic pressure field:

$$\begin{aligned} P_{dyn}(r) &= \int -\frac{k}{r^2}\,dr \\ P_{dyn}(r) &= \frac{k}{r} \end{aligned} \tag{56}$$

By Bernoulli conservation ($P_{static} = P_c - P_{dyn}$), the spatial gradient of the static pressure equates to the inverse dynamic gradient:

$$\nabla P_{static} = -\nabla P_{dyn} = -\left(-\frac{k}{r^2}\hat{\mathbf{r}}\right) = \frac{k}{r^2}\hat{\mathbf{r}} \tag{57}$$

Inserting this restorative pressure gradient into the metric buoyancy equation (53) calculates the physical force exerted upon an advecting boundary:

$$\begin{aligned} \mathbf{F}_g &= -V_{eff}\nabla P_{static} \\ \mathbf{F}_g &= -\left(\frac{m_\tau}{\rho_\tau}\right)\left(\frac{k}{r^2}\hat{\mathbf{r}}\right) \end{aligned} \tag{58}$$

To bridge this continuum mechanic to classical kinematics, empirical mass (M) maps to the geometry. The continuous metric does not treat mass as an independent input parameter; rather, it evaluates here as the structural magnitude of the defect (k) normalized by the metric density (ρ_τ) and the macroscopic displacement limit (G).

Applying the geometric limits of macroscopic measurement ($G = l_p c^2/m_p$ and $\rho_\tau = m_p/2\pi l_p^3$) strips the empirical constants. The hydrodynamic volume displacement defining this boundary evaluates in Section 8, but the scalar isomorphism resolves the boundary ratio as a Planck-scale hemisphere:

$$M \equiv \frac{k}{\rho_\tau G} \implies k = \frac{M}{2\pi t_p^2} \tag{59}$$

Substituting this normalized definition into the dynamic pressure field recovers the kinematic profile:

$$P_{dyn} = \frac{M}{2\pi t_p^2 r} = \rho_\tau\frac{GM}{r} \tag{60}$$

Substituting the identical definition into the buoyant force equation isolates the Newtonian limit:

$$\mathbf{F}_g = -\frac{GMm_\tau}{r^2}\hat{\mathbf{r}} \tag{61}$$

The tensor geodesic equation functions as the physical definition of Archimedes' Principle. Universal gravitation manifests as the symmetric spatial gradients of the continuous metric, adopting macroscopic mass as a consequence of dynamic pressure.

6. Unified Action & Singularity Resolution

The total unified action emerges as a consequence of the tensor contraction.

6.1 Unified Action

To evaluate the global energy equilibrium of the tensor, the action is extracted via the trace contraction of the Metric-Tensor against itself ($\hat{U}_{\mu\nu}\hat{U}^{\mu\nu}$). Because the tensor decomposes into symmetric ($S_{\mu\nu}$) and antisymmetric ($A_{\mu\nu}$) components, the algebraic expansion produces:

$$\hat{U}_{\mu\nu}\hat{U}^{\mu\nu} = S_{\mu\nu}S^{\mu\nu} + A_{\mu\nu}A^{\mu\nu} + 2S_{\mu\nu}A^{\mu\nu} \tag{62}$$

By the rules of tensor calculus, the inner product of a symmetric matrix and an antisymmetric matrix reduces to zero ($S_{\mu\nu}A^{\mu\nu} = 0$).

Because macroscopic expansion ($v_0(r)$) acts as radial volumetric strain, the strain partitions into independent scalars:

$$\hat{U}_{\mu\nu}\hat{U}^{\mu\nu} = S_{\mu\nu}S^{\mu\nu} + A_{\mu\nu}A^{\mu\nu} \tag{63}$$

These two macroscopic actions emerge from a single constraint:

- **The Gravitational Action ($S_{\mu\nu}S^{\mu\nu}$):** The contraction of the symmetric diagonals captures the volumetric compression and spatial packing of the coordinate grid. Because the unified stress tensor is divergence-free ($\nabla_\mu \hat{T}^{\mu\nu} = 0$), Lovelock's Theorem [34] mandates that its symmetric geometric representation must map to the Einstein Tensor ($\hat{T}^{\mu\nu} \propto G^{\mu\nu}$). Applying the background boundary scaling derived in Section 4, this proportionality defines the macroscopic curvature limit:

$$P_{macro}(S_{\mu\nu} - \eta_{\mu\nu}) = \frac{c^4}{8\pi G}G_{\mu\nu} \implies S_{\mu\nu} = \frac{c^4}{8\pi G P_{macro}}G_{\mu\nu} + \eta_{\mu\nu} \tag{64}$$

  Because the cosmological constant equates to this ratio ($\Lambda = 8\pi G P_{macro}/c^4$), substituting this limit ($S_{\mu\nu} = \frac{1}{\Lambda}G_{\mu\nu} + \eta_{\mu\nu}$) resolves the strain against classical curvature invariants:

$$\begin{aligned} S_{\mu\nu}S^{\mu\nu} &= \left(\frac{1}{\Lambda}G_{\mu\nu} + \eta_{\mu\nu}\right)\left(\frac{1}{\Lambda}G^{\mu\nu} + \eta^{\mu\nu}\right) \\ S_{\mu\nu}S^{\mu\nu} &= \frac{1}{\Lambda^2}G_{\mu\nu}G^{\mu\nu} + \frac{2}{\Lambda}G_{\mu\nu}\eta^{\mu\nu} + \eta_{\mu\nu}\eta^{\mu\nu} \end{aligned} \tag{65}$$

  Because the Einstein Tensor is defined as $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}R\eta_{\mu\nu}$, expanding the squared term ($G_{\mu\nu}G^{\mu\nu}$) cancels the scalar traces:

$$G_{\mu\nu}G^{\mu\nu} = \left(R_{\mu\nu} - \frac{1}{2}R\eta_{\mu\nu}\right)\left(R^{\mu\nu} - \frac{1}{2}R\eta^{\mu\nu}\right) = R_{\mu\nu}R^{\mu\nu} - R^2 + \frac{1}{4}R^2(4) = R_{\mu\nu}R^{\mu\nu} \tag{66}$$

  Substituting this identity, the symmetric contraction maps to a quadratic gravitational Lagrangian:

$$S_{\mu\nu}S^{\mu\nu} = \frac{1}{\Lambda^2}R_{\mu\nu}R^{\mu\nu} - \frac{2}{\Lambda}R + 4 \tag{67}$$

  Therefore, volumetric strain contraction generates the Ricci scalar (R) alongside the higher-order quadratic curvature ($R_{\mu\nu}R^{\mu\nu}$) required for topological boundary regularization.

- **The Electromagnetic Action ($A_{\mu\nu}A^{\mu\nu}$):** The contraction of the antisymmetric off-diagonals quantifies propagating transverse shear and Clebsch vorticity. Because these terms carry the modulus ratio (α), the resulting scalar matches the Maxwell invariant ($E^2 - B^2$) attenuated by the transverse coupling limit.

Gravity and electromagnetism manifest as orthogonal mechanical responses (longitudinal compression versus transverse shear) originating from the same thermodynamic budget.

6.2 Strain Incompatibility & Riemann Curvature

Macroscopic curvature of the metric is derived via the continuous deformation of the Tensor ($\hat{U}_{\mu\nu}$) against the geometric constraints of a Euclidean background.

1. Cauchy Strain Extraction:

The symmetric spatial components of the tensor ($S_{ij} = 1/\alpha_s^2$) constrain the effective volumetric packing. The symmetric Cauchy strain tensor (ε_{ij}) maps to the deviation of these spatial diagonals from the flat unperturbed background (δ_{ij}):

$$\varepsilon_{ij} = \frac{1}{2}\left(S_{ij} - \delta_{ij}\right) \tag{68}$$

2. The Saint-Venant Compatibility Operator:

An arbitrary strain field (ε_{ij}) is constrained within a continuous medium via the Saint-Venant Compatibility Condition ($\mathrm{Inc}(\boldsymbol{\varepsilon})_{ijkl} = 0$).

The Incompatibility Tensor measures the failure of the continuum to satisfy this closure. It is constructed as the linear combination of the second-order spatial derivatives of the strain field:

$$\mathrm{Inc}(\boldsymbol{\varepsilon})_{ijkl} = \partial_j\partial_l\varepsilon_{ik} + \partial_i\partial_k\varepsilon_{jl} - \partial_j\partial_k\varepsilon_{il} - \partial_i\partial_l\varepsilon_{jk} \tag{69}$$

3. Topological Dislocation & The Burgers Vector:

In an unperturbed vacuum, the Clebsch vorticity is zero ($\boldsymbol{\Omega} = 0$), the spatial displacement (u_i) is single-valued, and the incompatibility tensor reduces to zero ($\mathrm{Inc}(\boldsymbol{\varepsilon}) = 0$). The metric operates as flat Euclidean space.

The generation of discrete mass corresponds to the formation of a rotational Clebsch vortex. Because the metric circulates around the topological core, the displacement field (u_i) becomes multi-valued. Integrating the spatial displacement around a closed contour (C) of the vortex core produces a phase gap, defined as the Burgers vector (b_i) [39]:

$$\oint_C du_i = b_i \tag{70}$$

This functions physically as a Volterra dislocation. Because the continuous streamlines cannot close around the rotational core without a phase shift, the strain field is structurally incompatible ($\mathrm{Inc}(\boldsymbol{\varepsilon}) \neq 0$).

4. Macroscopic Riemann Curvature:

To conserve continuum stability across this dislocation, the surrounding spatial metric yields from the flat Euclidean plane to accommodate the missing phase.

In the weak-field limit of infinitesimal strain, the Saint-Venant incompatibility tensor ($\mathrm{Inc}(\boldsymbol{\varepsilon})_{ijkl}$) generates the combination of second-order spatial derivatives comprising the spatial components of the linearized Riemann Curvature Tensor (R_{ijkl}) [27]:

$$R^{(lin)}_{ijkl} = \mathrm{Inc}(\boldsymbol{\varepsilon})_{ijkl} \tag{71}$$

As structural loads approach the cavitation boundary ($r \to r_c$), the strain remains finite. Evaluating this non-linear regime maps the continuous deformation to the Kröner incompatibility tensor. The fully non-linear Riemann curvature, including the self-interacting connection terms ($\Gamma\Gamma$), recovers the finite strain incompatibility required to resolve the topological rotational defect.

6.3 Volumetric Cavitation & The Metric Lagrangian

1. The Symmetric Boundary (Volumetric Cavitation):

The spatial diagonals ($S_{ii} = 1/\alpha_s^2$) represent the compliance of the coordinate grid. Setting

the dynamic pressure ($P_{dyn}(r) = \rho_\tau GM/r$) equal to the critical vacuum capacity ($P_c = \frac{1}{2}\rho_\tau c^2$) collapses the Total Scalar Admittance ($\alpha_s \to 0$):

$$P_{dyn}(r_c) = P_c \implies \cancel{\rho_\tau}\frac{GM}{r_c} = \frac{1}{2}\cancel{\rho_\tau}c^2 \tag{72}$$

Calculating the cavitation radius outputs the Schwarzschild boundary ($r_c = 2GM/c^2$).

2. The Symmetric Lagrangian Components:
The total action partitions into symmetric (gravitational) and antisymmetric (gauge) sectors. Targeting the symmetric tensor ($S_{\mu\nu}$) evaluates the macroscopic strain.

The kinematic gradient energy formulates the square of the 4-gradient, mapping to the bosonic scalar propagation term:

$$\mathcal{L}_{grad} = -(\partial_\alpha S_{\mu\nu})(\partial^\alpha S^{\mu\nu}) \equiv -|D\phi|^2 \tag{73}$$

The static potential resolves the trace contraction ($S_{\mu\nu}S^{\mu\nu} \equiv m^2\phi^2$). Volumetric contraction reduces the equation to the quadratic curvature invariant ($R_{\mu\nu}R^{\mu\nu}$). Under the harmonic gauge condition ($\partial_\mu S^{\mu\nu} = \frac{1}{2}\partial^\nu S^\mu_\mu$), the Ricci curvature scales proportionally to the d'Alembertian ($\Box S_{\mu\nu}$):

$$\mathcal{L}_{static} \propto -R_{\mu\nu}R^{\mu\nu} \propto -(\Box S_{\mu\nu})(\Box S^{\mu\nu}) \tag{74}$$

To satisfy dimensional analysis, the kinetic gradient ($1/L^2$) and the higher-order curvature potential ($1/L^4$) are balanced by multiplying the resulting potential with the square of the cavitation radius (r_c^2). This constructs the raw Lagrangian prior to integration:

$$\mathcal{L}_{sym} \propto -(\partial_\alpha S_{\mu\nu})(\partial^\alpha S^{\mu\nu}) - r_c^2(\Box S_{\mu\nu})(\Box S^{\mu\nu}) \tag{75}$$

3. Integration by Parts (The d'Alembertian Operator):
Because the tensor evaluates dynamic strain over a flat Minkowski background ($\sqrt{-\eta} = 1$), the continuous action ($\mathcal{S} = \int \mathcal{L}_{sym}\, d^4x$) transforms the gradient term via integration by parts using standard partial derivatives. Bounding the surface terms to zero at infinity generates the linear 4D d'Alembertian:

$$\int -(\partial_\alpha S_{\mu\nu})(\partial^\alpha S^{\mu\nu})\, d^4x = \int S_{\mu\nu}\Box S^{\mu\nu}\, d^4x \tag{76}$$

Substituting this identity constructs the requisite geometric form required for higher-order gravity:

$$\mathcal{L}_{sym} \propto S_{\mu\nu}\Box S^{\mu\nu} - r_c^2(\Box S_{\mu\nu})(\Box S^{\mu\nu}) \tag{77}$$

6.4 Euler-Lagrange Variation & Singularity Resolution

To evaluate the macroscopic field equations, the principle of least action is applied to the total Lagrangian, summing the derived geometric strain ($\mathcal{L}_{geom}$) and the internal dynamic mass ($\mathcal{L}_{matter}$). Applying the higher-order Euler-Lagrange variation to this total action ensures structural equilibrium across the continuum:

$$\frac{\delta\mathcal{L}_{total}}{\delta S_{\mu\nu}} = \frac{\partial\mathcal{L}}{\partial S_{\mu\nu}} - \partial_\alpha\frac{\partial\mathcal{L}}{\partial(\partial_\alpha S_{\mu\nu})} + \Box\frac{\partial\mathcal{L}}{\partial(\Box S_{\mu\nu})} = 0 \tag{78}$$

Because the physical mass density (ρ_{dyn}) is explicitly extracted from the trace contraction of the tensor itself (Section 8.3), the matter action does not require external parameters. Classical mechanics filters directly through the symmetric spatial diagonals (S_{ii}), where the variation of the matter term defines the source coupling ($\delta\mathcal{L}_{matter}/\delta S_{ii} = -\kappa\rho_{dyn}$). Formulating the total classical Lagrangian yields:

$$\mathcal{L}_{classical} = \frac{1}{2}S_{ii}\Box S_{ii} - \frac{r_c^2}{2}(\Box S_{ii})^2 - \kappa\rho_{dyn}S_{ii} \tag{79}$$

At the static weak-field limit ($P_{dyn} = 0$), the scalar potential (Φ) drives the volumetric contraction linearly:

$$S_{ii} \approx 1 - \frac{2\Phi}{c^2} \tag{80}$$

Evaluating the variation at this static limit converts the d'Alembertian to the 3D Laplacian ($\Box \to \nabla^2$). Separating the strain into coupled differential operators:

$$\nabla^2 S_{ii} - r_c^2 \nabla^4 S_{ii} \propto \rho_{dyn} \tag{81}$$

Substituting Equation (80) directly into the variation isolates the scalar potential. The resulting negative sign on the spatial derivative is absorbed by the source coupling constant ($\kappa \equiv -8\pi G/c^2$), producing the 4th-order Modified Poisson Equation [43, 48]:

$$\nabla^2 \Phi - r_c^2 \nabla^4 \Phi = 4\pi G \rho_{dyn} \tag{82}$$

The Laplacian (∇^2) dominates the geometric curvature. At macroscopic coordinate boundaries where the spatial radius exceeds the boundary radius ($r \gg r_c$), the r_c^2 coefficient forces the $1/L^4$ biharmonic operator (∇^4) to attenuate to zero, restoring the Newtonian field limit ($\nabla^2\Phi \approx 4\pi G\rho_{dyn}$). Conversely, the negative biharmonic operator governs microscopic boundaries ($r \to r_c$). This strain-gradient elasticity restricts infinite compression, resolving the singularity. Because the cavitation limit operates as a cutoff, substituting the mass limit extracts the Schwarzschild radius ($r_c = 2GM/c^2$), truncating the depth of the metric and bridging continuum mechanics to macroscopic gravitational collapse.

6.5 The Unified Wave Operator & Phase Propagation

Applying the 4D d'Alembert wave operator ($\Box = \partial_\alpha \partial^\alpha$) to the full asymmetric tensor evaluates the continuous propagation of geometric strain across the Minkowski baseline. Because the derivative distributes linearly, the operator separates the tensor into orthogonal wave equations:

$$\Box \hat{U}_{\mu\nu} = \Box S_{\mu\nu} + \Box A_{\mu\nu} \tag{83}$$

1. The Symmetric Wave (Longitudinal Propagation):
Applying the operator to the spatial and temporal diagonals specifies the propagation of volumetric pressure perturbations. Taking the d'Alembertian of the macroscopic field equation (Equation 27) constructs the linearized gravitational wave equation:

$$\Box S_{\mu\nu} = \frac{8\pi G}{\Lambda c^4} \Box \hat{T}^{macro}_{(\mu\nu)} \tag{84}$$

This operator drives the longitudinal geometric strain (metric compression and expansion) through the continuum.

2. The Antisymmetric Wave (Transverse Propagation):
Applying the operator to the off-diagonals calculates the propagation of rotational shear. Because the antisymmetric matrix tracks the kinematic Clebsch fields, the geometric derivative forms the full transverse wave:

$$\Box A_{\mu\nu} = \alpha \Box \left(\frac{u_\mu v_{\perp,\nu} - u_\nu v_{\perp,\mu}}{c} \right) + \alpha t_p \Box \omega_{\mu\nu} \tag{85}$$

This operator guides the physical propagation of the electromagnetic gauge field.

3. The Source-Free Vacuum Limit:
In a source-free vacuum, the macroscopic propagation of the stress tensor equates to zero ($\Box \hat{T}_{\mu\nu} = 0$). By the unified field equation, this sets the geometric wave operator to zero:

$$\Box \hat{U}_{\mu\nu} = 0 \implies \left(-\frac{1}{c^2}\frac{\partial^2}{\partial t^2} + \nabla^2 \right) \hat{U}_{\mu\nu} = 0 \tag{86}$$

Because both the symmetric diagonals and antisymmetric off-diagonals are bounded by the identical geometric wave operator, longitudinal volumetric strain (gravity) and transverse rotational strain (electromagnetism) are forced to propagate at the identical acoustic phase limit of the continuum (c).

7. Conformal Symmetry & The Lie Derivative Constraints

The Conformal Group $SO(4,2)$ bounds the causal structure of the metric. Applying the 15 vector fields (ξ^μ) to the Asymmetric Strain Tensor via the Lie derivative ($\mathcal{L}_\xi$) identifies generic symmetry-breaking conditions.

Standard Conformal Killing Equations operate on symmetric metrics; therefore, the Lie derivative distributes across the asymmetric tensor. The tensor partitions into symmetric volumetric diagonals ($S_{\mu\nu}$) and antisymmetric shear/vorticity off-diagonals ($A_{\mu\nu}$):

$$\hat{U}_{\mu\nu} = S_{\mu\nu} + A_{\mu\nu} \tag{87}$$

The Lie derivative produces the kinematic transport along a flow path. Substituting the decomposed tensor isolates the symmetric and antisymmetric transformation rules:

$$\mathcal{L}_\xi \hat{U}_{\mu\nu} = \xi^\alpha \partial_\alpha (S_{\mu\nu} + A_{\mu\nu}) + (S_{\alpha\nu} + A_{\alpha\nu})\partial_\mu \xi^\alpha + (S_{\mu\alpha} + A_{\mu\alpha})\partial_\nu \xi^\alpha \tag{88}$$

7.1 The Poincare Subgroup (10 Generators)

The Poincare subgroup tests for volume-preserving transport. For Poincare symmetry to hold, the Lie derivative of the full tensor must equal zero ($\mathcal{L}_\xi \hat{U}_{\mu\nu} = 0$).

1. Translations (P_μ, 4 Generators):
For a constant translation parameter ϵ^μ, the vector field assigns $\xi^\mu = \epsilon^\mu$. The coordinate derivative vanishes ($\partial_\nu \xi^\alpha = 0$). Substituting this into Equation (88) extracts the position-dependent gradient of the tensor:

$$\mathcal{L}_\xi \hat{U}_{\mu\nu} = \epsilon^\alpha (\partial_\alpha S_{\mu\nu} + \partial_\alpha A_{\mu\nu}) \tag{89}$$

For translation symmetry, the gradient of both conjugate fields must vanish ($\partial_\alpha \hat{U}_{\mu\nu} = 0$), enforcing equilibrium. Substituting the macroscopic Hubble velocity field into the temporal diagonal ($\hat{U}_{00} \propto \Delta P_{macro}$) generates a temporal gradient ($\partial_0 \hat{U}_{00} \neq 0$). Furthermore, an ambient pressure gradient ($-\nabla \mathcal{P}_{ambient}$) leaves a spatial gradient ($\partial_i \hat{U}_{\mu\nu} \neq 0$). This demonstrates that macroscopic expansion and ambient gravity break translation symmetry.

2. Lorentz Transformations ($M_{\mu\nu}$, 6 Generators):
The vector field is constructed by an antisymmetric rotation matrix ($\omega_{\mu\nu} = -\omega_{\nu\mu}$) such that $\xi^\mu = \omega^\mu{}_\alpha x^\alpha$. The spatial derivative computes the constant transformation matrix ($\partial_\nu \xi^\mu = \omega^\mu{}_\nu$).

Substituting this into the distributed Lie derivative expands the rotational shear acting upon the full tensor:

$$\mathcal{L}_\xi \hat{U}_{\mu\nu} = (\omega^\alpha{}_\beta x^\beta)\partial_\alpha \hat{U}_{\mu\nu} + (S_{\alpha\nu} + A_{\alpha\nu})\omega^\alpha{}_\mu + (S_{\mu\alpha} + A_{\mu\alpha})\omega^\alpha{}_\nu \tag{90}$$

To separate the geometric couplings, we contract the symmetric tensor components ($S_{\mu\nu}$). Using the covariant formulation $S_{\mu\nu} = -\alpha_s^2 u_\mu u_\nu + \frac{1}{\alpha_s^2} h_{\mu\nu}$, the unperturbed isotropic spatial metric cancels the antisymmetric transformation ($\omega_{\mu\nu} + \omega_{\nu\mu} = 0$). The symmetric contraction outputs the 4-velocity projection:

$$S_{\alpha\nu}\omega^\alpha{}_\mu + S_{\mu\alpha}\omega^\alpha{}_\nu = \left(\frac{1}{\alpha_s^2} - \alpha_s^2\right)(u_\nu u_\alpha \omega^\alpha{}_\mu + u_\mu u_\alpha \omega^\alpha{}_\nu) \tag{91}$$

This residual vanishes for spatial rotations ($SO(3)$) because the temporal 4-velocity is orthogonal to spatial vorticity, preserving the volumetric trace. However, the 3 Lorentz boosts (M_{0i}) mix the temporal and spatial capacities, generating a residual for the time-space components:

$$S_{\alpha i}\omega^{\alpha}{}_{0} + S_{0\alpha}\omega^{\alpha}{}_{i} = -\omega_{0i}\left(\frac{1}{\alpha_s^2} - \alpha_s^2\right) \tag{92}$$

Because the Lie derivative preserves index symmetry, the symmetric and antisymmetric constraints must be satisfied independently ($\mathcal{L}_\xi S_{\mu\nu} \propto S_{\mu\nu}$ and $\mathcal{L}_\xi A_{\mu\nu} \propto A_{\mu\nu}$).

Computing the symmetric sector for a Lorentz boost (M_{0i}), the time-space components ($0i$) target the residual:

$$(\mathcal{L}_\xi S)_{0i} = \xi^\alpha \partial_\alpha S_{0i} - \omega_{0i}\left(\frac{1}{\alpha_s^2} - \alpha_s^2\right) \tag{93}$$

For Lorentz symmetry to hold across the symmetric sector, this residual must vanish. Because this symmetric time-space residual cannot algebraically cancel against the antisymmetric transverse sector ($A_{\mu\nu}$), Lorentz boost symmetry is broken unless the continuum is static ($\alpha_s = 1$).

Assessing the antisymmetric sector independently exposes the rotational couplings:

$$(\mathcal{L}_\xi A)_{\mu\nu} = \xi^\alpha \partial_\alpha A_{\mu\nu} + (A_{\mu\alpha}\omega^{\alpha}{}_{\nu} - \omega_{\mu\alpha}A^{\alpha}{}_{\nu}) \tag{94}$$

This demonstrates that while spatial rotations (M_{ij}) correspond to the antisymmetric internal Clebsch vorticity ($\omega_{\mu\nu}$), kinematic translation through the continuum ($v \to c$, $\alpha_s \to 0$) introduces a symmetric residual that structurally breaks the Lorentz invariance of the metric.

7.2 Dilation & Special Conformal Transformations (5 Generators)

The Dilation (D) and Special Conformal (K_μ) vector fields scale the 4D coordinate volume. Because these paths compress or expand the metric, the constraint utilizes the Conformal Killing Equation. The Lie derivative resolves to a scalar multiple of the tensor:

$$\mathcal{L}_\xi \hat{U}_{\mu\nu} = f(x)\hat{U}_{\mu\nu} \tag{95}$$

1. The Dilation Generator (D):
The vector field is parameterized by a constant scalar σ, constructed as $\xi^\mu = \sigma x^\mu$. The coordinate derivative yields the Kronecker delta ($\partial_\nu \xi^\mu = \sigma\delta^\mu_\nu$).

Substituting these definitions into the Lie derivative unfolds the operator:

$$\begin{aligned}\mathcal{L}_\xi \hat{U}_{\mu\nu} &= (\sigma x^\alpha)\partial_\alpha \hat{U}_{\mu\nu} + \hat{U}_{\alpha\nu}(\sigma\delta^\alpha_\mu) + \hat{U}_{\mu\alpha}(\sigma\delta^\alpha_\nu) \\ \mathcal{L}_\xi \hat{U}_{\mu\nu} &= \sigma x^\alpha \partial_\alpha \hat{U}_{\mu\nu} + 2\sigma\hat{U}_{\mu\nu}\end{aligned} \tag{96}$$

To satisfy scale invariance ($f(x) = 2\sigma$), the position-dependent gradient must vanish ($\sigma x^\alpha \partial_\alpha \hat{U}_{\mu\nu} = 0$). This constraint requires a homogeneous metric field. Discrete topological boundaries manifest as Volterra dislocations. Because the rotational Clebsch phase generates a Burgers vector, the metric violates the Saint-Venant compatibility condition ($\mathrm{Inc}(\boldsymbol{\varepsilon}) \neq 0$). This strain incompatibility ensures the metric varies with coordinate position ($\partial_\alpha \hat{U}_{\mu\nu} \neq 0$), structurally breaking scale invariance.

2. Special Conformal Transformations (K_μ, 4 Generators):
The vector field operates via a constant parameter (b^μ) with dimensions of inverse length (L^{-1}):

$$\xi^\mu = 2(x_\beta b^\beta)x^\mu - (x_\beta x^\beta)b^\mu \tag{97}$$

Expanding the spatial derivative retrieves the local scaling and rotational shift:

$$\partial_\alpha \xi^\mu = 2b_\alpha x^\mu + 2(x_\beta b^\beta)\delta^\mu_\alpha - 2x_\alpha b^\mu \tag{98}$$

Substituting this transformation into the Lie derivative distributes the non-inertial strain terms:

$$\mathcal{L}_\xi \hat{U}_{\mu\nu} = \xi^\alpha \partial_\alpha \hat{U}_{\mu\nu} + 4(x_\beta b^\beta)\hat{U}_{\mu\nu} + 2x^\alpha(b_\mu \hat{U}_{\alpha\nu} + b_\nu \hat{U}_{\mu\alpha}) - 2b^\alpha(x_\mu \hat{U}_{\alpha\nu} + x_\nu \hat{U}_{\mu\alpha}) \tag{99}$$

Contracting the trailing antisymmetric covariant combinations against the metric reveals that the spatial components (i, j) mutually cancel due to spatial isotropy. However, mixing the time-space components produces a residual strain:

$$2(x^0 b^1 - x^1 b^0)\left(\frac{1}{\alpha_s^2} - \alpha_s^2\right) \tag{100}$$

This residual vanishes only in a static vacuum ($\alpha_s = 1$). Under dynamic pressure ($\alpha_s < 1$), the Special Conformal transformation induces non-inertial shear across the metric boundaries.

To calculate the remaining volumetric trace $4(x_\beta b^\beta)\hat{U}_{\mu\nu}$, the global parameter (b^μ) must align dimensionally to a physical field without violating the product rule of the constant Lie generator. We designate b^μ as the global symmetry generator bounding a locally uniform kinematic acceleration field (a^μ, where $\partial_\alpha a^\mu = 0$) across the phase limit ($b^\mu \equiv a^\mu/c^2$). A non-uniform local field ($\partial_\alpha a^\mu \neq 0$) generates tidal shearing, which breaks the conformal symmetry and sets the structural stability limit of the translating defect.

Substituting the spatial material derivative for an accelerating volume ($a^i = -\frac{1}{\rho_\tau}\nabla^i P_{dyn}$) derives the thermodynamic scaling constraint:

$$\begin{aligned} 4(x_\beta b^\beta) &= 4\left(x_i \frac{a^i}{c^2}\right) \\ 4(x_\beta b^\beta) &= -\frac{4}{\rho_\tau c^2}\left(x_i \nabla^i P_{dyn}\right) \\ 4(x_\beta b^\beta) &= -\frac{2}{P_c}\left(x_i \nabla^i P_{dyn}\right) \end{aligned} \tag{101}$$

This geometric scaling factor operates proportionally to the directional derivative of dynamic pressure relative to the critical vacuum capacity ($P_c = \frac{1}{2}\rho_\tau c^2$). A conformal transformation manifests as a topological boundary propagating through a scalar pressure gradient, where the convergence of the coordinate lines governs the refraction of the trajectory.

3. Derivation of the Gordon Optical Metric:
This directly alters the apparent phase velocity of transverse shear waves (light) propagating through the medium. The continuous metric acts as a Gradient-Index (GRIN) optical medium.

The macroscopic index of refraction (n) of the vacuum is formulated as the ratio of the unperturbed baseline capacity (P_c) to the available static capacity (P_{static}):

$$n(x) = \frac{P_c}{P_c - P_{dyn}} = \frac{1}{\alpha_s^2} \tag{102}$$

As dynamic pressure increases ($P_{dyn} \to P_c$), the scalar admittance collapses ($\alpha_s^2 \to 0$) and the refractive index diverges ($n \to \infty$), reproducing the black hole horizon as an optical phase limit.

To formalize the trajectory of a shear wave through this pressure gradient, the conformal scaling constraint (Equation 101) maps to the Gordon optical metric [21]. If the defect translates with a normalized four-velocity ($\hat{u}_\mu \hat{u}^\mu = -1$), the effective optical spacetime ($\hat{U}^{opt}_{\mu\nu}$) experienced by the propagating wave is constructed by superimposing the refractive penalty onto the baseline metric:

$$\hat{U}^{opt}_{\mu\nu} = \hat{U}_{\mu\nu} + \left(1 - \frac{1}{n^2}\right)\hat{u}_\mu \hat{u}_\nu \tag{103}$$

Substituting the thermodynamic formulation of the refractive index from Equation (102) determines the acoustic metric:

$$\hat{U}^{opt}_{\mu\nu} = \hat{U}_{\mu\nu} + \left(1 - \alpha_s^4\right) \hat{u}_\mu \hat{u}_\nu \tag{104}$$

Under the conformal transformation (K_μ), acceleration (b^μ) generates a dynamic pressure gradient, affecting scalar admittance (α_s). The continuous coordinate transformation causes null geodesics ($ds^2 = 0$) to bend, reproducing gravitational lensing as acoustic refraction of a wave traversing a density gradient.

7.3 Kinematic Implications of Symmetry Breaking

The Lie derivative assessments confirm that Lorentz and conformal symmetries hold for a static continuum ($\alpha_s = 1$). Under applied kinematic strain, the asymmetric tensor generates residuals proportional to the dynamic pressure gradient (∇P_{dyn}) and the temporal capacity variance ($1/\alpha_s^2 - \alpha_s^2$). Substituting these conformal boundary constraints into the Gordon metric (Equation 103) demonstrates that impedance dictates the phase velocity of the vacuum. Consequently, null geodesic deflection and acoustic horizons emerge from the thermodynamic limits.

7.4 Conserved Noether Currents

By Noether's theorem, the invariances of the Poincare Lie derivatives determine the conserved kinematic currents.

For a continuous unified Lagrangian action ($\mathcal{L}_{metric}$) invariant under a generalized coordinate transformation ($\delta x^\mu = \xi^\mu$), the conserved Noether current density (J^μ) expands as:

$$J^\mu = \frac{\partial \mathcal{L}_{metric}}{\partial(\partial_\mu \hat{U}_{\alpha\beta})} \mathcal{L}_\xi \hat{U}_{\alpha\beta} - \xi^\mu \mathcal{L}_{metric} \tag{105}$$

Substituting the zero-residual Lie symmetries from Section 7.1 isolates the orthogonal conserved currents of the asymmetric metric:

1. Translational Symmetry (Energy-Momentum Conservation):
Under uniform spacetime translation ($\xi^\mu = \epsilon^\mu$), the Lie derivative yields zero gradient ($\partial_\alpha \hat{U}_{\mu\nu} = 0$). The Noether current equates to the macroscopic continuous field equation ($\hat{T}^\mu{}_\nu$). The continuity equation constrains the 4-divergence to zero:

$$\partial_\mu \hat{T}^{\mu\nu} = 0 \tag{106}$$

This operation conserves continuous energy and linear momentum.

2. Lorentz Symmetry (Total Angular Momentum & Spin):
Under spatial rotation ($\xi^\mu = \omega^\mu{}_\alpha x^\alpha$), the macroscopic energy-momentum tensor defines orbital angular momentum ($L^{\mu\alpha\beta} = x^\alpha \hat{T}^{\mu\beta} - x^\beta \hat{T}^{\mu\alpha}$). Because the tensor incorporates an antisymmetric transverse sector ($A_{\mu\nu}$), the Noether current generates an internal Clebsch spin tensor ($\Sigma^{\mu\alpha\beta}$):

$$J^\mu_{(Lorentz)} = M^{\mu\alpha\beta} = L^{\mu\alpha\beta} + \Sigma^{\mu\alpha\beta} \tag{107}$$

Because the antisymmetric rotational couplings resolve the residual ($A_{\mu\alpha}\omega^\alpha{}_\nu - \omega_{\mu\alpha}A^\alpha{}_\nu = 0$), the divergence of the total angular momentum converges to zero ($\partial_\mu M^{\mu\alpha\beta} = 0$). Internal Clebsch vorticity recovers conserved quantum spin.

7.5 The Dressing Field Method (Gauge-Invariant Composites)

The continuous extraction of physical degrees of freedom from the metric avoids spontaneous symmetry breaking and gauge-fixing artifacts via the Dressing Field Method (DFM) [19]. In

general relativistic gauge field theories, the topological obstruction to global gauge-fixing (the Gribov-Singer ambiguity [22, 47]) is bypassed by constructing composite, gauge-invariant variables directly from the fields.

The Asymmetric Metric-Tensor ($\hat{U}_{\mu\nu}$) operates as a dressed field space, utilizing the Clebsch parameters to extract the invariant content of the continuum without requiring an absolute background manifold.

1. $U(1)$ Gauge Dressing (Electromagnetism):
The Clebsch parametrization (Equation 1) executes an Abelian DFM operation. In standard DFM formalism, a bare gauge connection (A) and a dressing field (u) combine to form a gauge-invariant composite ($\hat{A} = A + du$).

Within the asymmetric tensor, the transverse slip 1-form ($v_\perp$) operates as the bare gauge connection (A). The irrotational phase scalar (h) operates as the compensatory dressing field (u). Under a local $U(1)$ gauge transformation parameterized by χ, the bare fields transform as:

$$v_\perp^g = v_\perp - d\chi, \quad h^g = h + \chi \tag{108}$$

The composite kinematic field ($\hat{\mathbf{v}}$) is constructed via the DFM addition:

$$\hat{\mathbf{v}}^g = v_\perp^g + dh^g = (v_\perp - d\chi) + d(h + \chi) = v_\perp + dh = \hat{\mathbf{v}} \tag{109}$$

The composite resolves as gauge-invariant ($\delta_\chi \hat{\mathbf{v}} = 0$). Consequently, the antisymmetric off-diagonals ($\hat{U}_{0i} \propto \hat{\mathbf{v}}_i$) representing electromagnetic transverse shear are dressed physical observables rather than gauge-dependent potentials.

2. Diffeomorphism Invariance (Diff(M)) and Relational Spacetime:
To satisfy relational physics, the continuous metric is not evaluated against a fixed Euclidean stage. DFM for diffeomorphisms requires a dressing field that transforms under Diff(M) to counteract the Lie derivative ($\mathcal{L}_\xi$) of the bare coordinate manifold.

The Clebsch rotational phases ($\lambda_\Omega, \beta_\Omega$) act as these relational dressing fields. Under a diffeomorphism generated by a vector field ξ, the fields co-define the invariant geometric vorticity ($\hat{\Omega}$):

$$\hat{\Omega}_{\mu\nu} = \partial_{[\mu}\lambda_\Omega \partial_{\nu]}\beta_\Omega \tag{110}$$

By establishing internal spatial coordinates through this exterior product intersection ($d\lambda_\Omega \wedge d\beta_\Omega$), the metric defines its own reference frame.

This relational structure evaluates physical states as boundary transitions. Internal composition resolves not as a discrete volume, but as nested geometric yield limits scaling down to the Planck boundary. A phase-locked transition layer (defined via Clebsch vorticity) acts as a Quantum Reference Frame (QRF) [20, 44]. Because the tensor provides an invariant mapping across these nested scales, evaluating covariance across distinct boundaries (ranging from microscopic cavitation limits to macroscopic shear layers) utilizes transformations of the second kind (active gauge transformations that switch the physical reference boundary). This maps kinematic strain and physical action between relational interfaces, establishing force as a geometric boundary condition rather than an intrinsic property.

3. Invariant Path Integral Quantization:
Because the Asymmetric Metric-Tensor ($\hat{U}_{\mu\nu}$) is assembled from these dressed, composite variables, it constitutes an invariant field space. The geometric components evaluated in the Lagrangian (Section 6.3) are devoid of redundant gauge orbits. By evaluating continuous physical states utilizing these dressed variables, an invariant path integral quantization prevents Gribov obstructions, permitting deterministic convergence at the yield limits.

8. The Mass Mechanism (Topological Tensor Yield)

While the conformal scaling constraints are restricted by the Lie derivatives (Section 7), rest mass emerges as a phase transition where kinematic loads decouple metric continuity.

8.1 The Tensor Yield Criterion (The VEV Origin)

The symmetric spatial diagonals of the tensor define the volumetric density of the coordinate grid. Because the spatial diagonals are derived via the inverse scalar admittance ($S_{ii} = 1/\alpha_s^2$), substituting the thermodynamic ratio ($\alpha_s^2 = P_{static}/P_c$) equates the strain to the available static capacity:

$$S_{ii} = \frac{P_c}{P_{static}} \tag{111}$$

Substituting the Bernoulli equilibrium ($P_{static} = P_c - P_{dyn}$) and the kinematic pressure of metric circulation ($P_{dyn} = \frac{1}{2}\rho_\tau v^2$) translates the coordinate expansion into a velocity-dependent boundary:

$$S_{ii} = \frac{P_c}{P_c - \frac{1}{2}\rho_\tau v^2} \tag{112}$$

Because the total static capacity of the vacuum equates to the phase limit of the continuum ($P_c = \frac{1}{2}\rho_\tau c^2$), the diagonals evaluate as a function of the velocity ratio:

$$\begin{aligned} S_{ii} &= \frac{\frac{1}{2}\rho_\tau c^2}{\frac{1}{2}\rho_\tau c^2 - \frac{1}{2}\rho_\tau v^2} \\ S_{ii} &= \frac{1}{1 - \frac{v^2}{c^2}} \end{aligned} \tag{113}$$

Taking the limit as velocity approaches the speed of light ($v \to c$) forces the spatial component to diverge:

$$\lim_{v \to c} S_{ii} = \lim_{v \to c} \frac{1}{1 - \frac{v^2}{c^2}} = \infty \tag{114}$$

Because the overarching continuum cannot sustain an infinite spatial coordinate geometry, the internal structure yields. The metric cavitates, thereby conserving the thermodynamic boundary (P_c).

This critical threshold acts as the Vacuum Expectation Value (v_{vev}) [51]. It is the specific velocity where dynamic pressure equals the total ambient capacity, forcing a geometric phase transition from a continuous manifold into a localized topological defect:

$$P_{dyn} = P_{ambient} \implies v_{vev}^2 = \frac{2P_{ambient}}{\rho_\tau} \tag{115}$$

Redefining the kinematic state relative to the baseline defines the longitudinal acoustic breathing mode ($\delta v(x)$):

$$v(x) = v_{vev} + \delta v(x) \tag{116}$$

8.2 The Topological Phase Lock (Mass Generation)

Before structural yield, the metric sustains continuous independent wave states (e.g., irrotational compression and solenoidal transverse shear).

When the kinematic strain crosses the VEV threshold ($v > v_{vev}$), the spatial diagonals ($1/\alpha_s^2$) diverge to infinity, and the metric undergoes cavitation.

Continuous phase conservation locks the unresolved kinetic energy into the antisymmetric off-diagonals ($A_{\mu\nu}$) of the metric tensor. The metric transitions from a compliant medium into a rotational boundary (a Clebsch vortex):

$$\alpha_s \to 0 \implies \text{Metric Cavitation} \implies \text{Discrete Rest Mass} \tag{117}$$

This topological yield converts the continuous metric into invariant mass.

8.3 The Kinematic Lagrangian & Orthogonal Mass Extraction

The scalar energy state is determined by the trace contraction of the tensor ($\hat{U}_{\mu\nu}\hat{U}^{\mu\nu} = S_{\mu\nu}S^{\mu\nu} + A_{\mu\nu}A^{\mu\nu}$). Expanding this contraction against the covariant definitions extracts the kinematic wave velocities.

By applying the covariant 4-velocity orthogonality constraints ($u_\mu u^\mu = -1$, $u_\mu v_\perp^\mu = 0$, and $u_\mu h^{\mu\nu} = 0$) it reduces to:

$$S_{\mu\nu}S^{\mu\nu} = \left(-\alpha_s^2 u_\mu u_\nu + \frac{1}{\alpha_s^2}h_{\mu\nu}\right)\left(-\alpha_s^2 u^\mu u^\nu + \frac{1}{\alpha_s^2}h^{\mu\nu}\right) = \alpha_s^4 + \frac{3}{\alpha_s^4} \tag{118}$$

Applying a weak-field Taylor expansion ($\alpha_s^2 = 1-\beta^2 \implies \alpha_s^4 \approx 1-2\beta^2$ and $\alpha_s^{-4} \approx 1+2\beta^2$), the trace simplifies to:

$$S_{\mu\nu}S^{\mu\nu} \approx (1-2\beta^2) + 3(1+2\beta^2) = 4 + 4\frac{v^2}{c^2} \tag{119}$$

This expansion operates as an approximation of the classical weak-field limit, not a physical law. As velocity approaches the phase limit ($v \to c$), the non-linear trace restricts the boundary. This approximation produces the total static capacity.

The antisymmetric trace expansion filters the squares of the transverse velocity ($v_\perp$) and internal vorticity ($\omega_{\mu\nu}$). Because vorticity is orthogonal to the temporal rest frame ($u_\mu\omega^{\mu\nu} = 0$), the cross-terms vanish. Expanding the covariant wedge product validates the remaining cancellation:

$$\begin{aligned}
A_{\mu\nu}A^{\mu\nu} &= \frac{\alpha^2}{c^2}(u_\mu v_{\perp,\nu} - u_\nu v_{\perp,\mu})(u^\mu v^{\perp,\nu} - u^\nu v^{\perp,\mu}) + \alpha^2 t_p^2 \omega_{\mu\nu}\omega^{\mu\nu} \\
A_{\mu\nu}A^{\mu\nu} &= \frac{\alpha^2}{c^2}\Big((u_\mu u^\mu)(v_{\perp,\nu}v^{\perp,\nu}) - (u_\mu v^{\perp,\mu})(v_{\perp,\nu}u^\nu) - (u_\nu v^{\perp,\nu})(v_{\perp,\mu}u^\mu) + (u_\nu u^\nu)(v_{\perp,\mu}v^{\perp,\mu})\Big) + \alpha^2 t_p^2 \omega_{\mu\nu}\omega^{\mu\nu} \\
A_{\mu\nu}A^{\mu\nu} &= \frac{\alpha^2}{c^2}\Big((-1)(v_\perp^2) - (0) - (0) + (-1)(v_\perp^2)\Big) + \alpha^2 t_p^2 \omega_{\mu\nu}\omega^{\mu\nu} \\
A_{\mu\nu}A^{\mu\nu} &= -2\alpha^2\frac{v_\perp^2}{c^2} + \alpha^2 t_p^2 \omega_{\mu\nu}\omega^{\mu\nu}
\end{aligned} \tag{120}$$

Because these traces measure dimensionless strain across four spacetime dimensions, subtracting the unperturbed Minkowski background trace ($\hat{U}_{\mu\nu}\hat{U}^{\mu\nu} - 4$) isolates the kinematic variance. Normalizing this trace by the spacetime dimensionality (1/4), mirroring standard classical field formulations ($\mathcal{L} \propto -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$), and multiplying by the background pressure (P_c) extracts the continuous kinematic Lagrangian.

Substituting the symmetric trace variance ($4\frac{v^2}{c^2}$) and expanding the thermodynamic coefficient ($P_c = \frac{1}{2}\rho_\tau c^2$):

$$\begin{aligned}
\mathcal{L}_{kin} &= \frac{1}{4}P_c\Big(\hat{U}_{\mu\nu}\hat{U}^{\mu\nu} - 4\Big) \\
\mathcal{L}_{kin} &= \frac{1}{4}\left(\frac{1}{2}\rho_\tau c^2\right)\left(4\frac{v^2}{c^2} + A_{\mu\nu}A^{\mu\nu}\right) \\
\mathcal{L}_{kin} &= \left(\frac{1}{2}\rho_\tau v^2\right) + \frac{1}{4}P_c\big(A_{\mu\nu}A^{\mu\nu}\big)
\end{aligned} \tag{121}$$

Because $\frac{1}{2}\rho_\tau v^2$ defines dynamic pressure (P_{dyn}), and the scaled antisymmetric trace produces transverse structural shear (P_{shear}):

$$\mathcal{L}_{kin} = P_{dyn} + P_{shear} \tag{122}$$

Mass density (ρ_{dyn}) is extracted from this tensor operation. Dividing the dynamic pressure component of the symmetric trace contraction by the squared phase boundary (c^2) outputs the physical density ($\rho_{dyn} = P_{dyn}/c^2$).

1. The Quantum Cavitation Boundary (Compton Limit):
While the symmetric trace bounds the macroscopic gravitational limit, the antisymmetric trace specifies the microscopic boundary. The term ($\alpha^2 t_p^2 \omega_{\mu\nu}\omega^{\mu\nu}$) represents internal shear, confining the rest mass energy of a topological defect ($E = mc^2$).

Because the cavitated boundary operates as an acoustic standing wave, its spatial circulation is phase-locked. The scalar magnitude of the spatial vorticity maps to the temporal phase frequency ($E = \hbar\omega$), resolving the frequency ($\omega = mc^2/\hbar$). The radius required to confine this rotational phase ($v = c$) evaluates as the phase velocity divided by the frequency:

$$r_c = \frac{c}{\omega} = \frac{c}{(mc^2/\hbar)} = \frac{\hbar}{mc} \tag{123}$$

This derives the reduced Compton wavelength as the cavitation radius of the antisymmetric tensor.

2. The Topological Yield Transition (Planck Mass):
Because the generic metric tensor ($\hat{U}_{\mu\nu}$) limits singularities at $r = r_c$, equating the symmetric gravitational boundary ($r_s = 2GM/c^2$) to the antisymmetric shear boundary ($r_c = \hbar/mc$) identifies the geometric scale where volumetric compression (P_{dyn}) and rotational shear (P_{shear}) exert equal thermodynamic strain:

$$\frac{\hbar}{mc} = \frac{2GM}{c^2} \implies M = \frac{M_P}{\sqrt{2}} \tag{124}$$

This isolates the Planck mass scale ($M = M_P/\sqrt{2}$) as the transition threshold between continuous gravitation and discrete quantum localization. The $1/\sqrt{2}$ coefficient derives from the metric compliance limit governing inertia. As established in the derivation of kinematic inertia, the structural compliance (α_s) determines the metric's capacity to bound mass density:

$$\alpha_s = \left(1 - \frac{P_{dyn} + P_{shear}}{P_c}\right)^{1/2} \tag{125}$$

As derived in Section 9.6, the transverse rotational eigenstates are forced into the complex plane ($1 \pm i$). At the cavitation boundary, the total metric capacity (P_c) is partitioned equally between the real volumetric axis (longitudinal compression) and the imaginary antisymmetric axis (transverse shear). Because observable rest mass is generated by the real longitudinal axis, the maximum bounded mass state correlates to half the total structural load ($P_{load} \to \frac{1}{2}P_c$).

Substituting this orthogonal pressure partition into the compliance identity scales the structural yield limit of the tensor:

$$\alpha_{yield} = \left(1 - \frac{1}{2}\right)^{1/2} = \frac{1}{\sqrt{2}} \tag{126}$$

Because a stable defect cannot possess a mass density exceeding the structural compliance of the metric bounding it, this partition projects the real bounded state. This precise yield threshold ($\alpha_{yield} = 1/\sqrt{2}$) extracts the identical Planck mass limit at the intersection boundary ($M/M_P = \alpha_{yield} = 1/\sqrt{2} \implies M = M_P/\sqrt{2}$).

Because a stable defect cannot possess a mass density exceeding the structural compliance of the metric bounding it, this partition projects the real bounded state. By mapping the metric compliance to the mass ratio ($M/M_P = \alpha_{yield} = 1/\sqrt{2} \implies M = M_P/\sqrt{2}$), this threshold extracts the same Planck mass limit at the intersection boundary.

9. Electromagnetic Gauge & Unitary Evolution

Eigenstate analysis reveals the parametric coupling and structural mechanics of the metric. Structural decoupling and macroscopic quantization are extracted via the principal eigenvalues (λ).

The principal axes of strain are calculated by evaluating the characteristic determinant:

$$\det(\hat{U}_{\mu\nu} - \lambda\delta_{\mu\nu}) = 0 \tag{127}$$

Specific components of the metric are evaluated by isolating individual parameters. Aligning the primary axis of rotation to the z-coordinate ($\Omega_z = \Omega$) induces an orthogonal transverse slip across the $x - y$ plane (v_x, v_y). This structures the matrix:

$$\hat{U}_{\mu\nu} - \lambda\delta_{\mu\nu} = \begin{bmatrix} -\alpha_s^2 - \lambda & -\alpha\frac{v_x}{c} & -\alpha\frac{v_y}{c} & 0 \\ \alpha\frac{v_x}{c} & \alpha_s^{-2} - \lambda & -\alpha t_p\Omega & 0 \\ \alpha\frac{v_y}{c} & \alpha t_p\Omega & \alpha_s^{-2} - \lambda & 0 \\ 0 & 0 & 0 & \alpha_s^{-2} - \lambda \end{bmatrix} \tag{128}$$

Expanding the characteristic determinant isolates the invariant longitudinal axis (z-axis). Within the coupled transverse dimensions, the antisymmetric cross-terms algebraically cancel ($v_x v_y\Omega - v_y v_x\Omega = 0$). The rotation couples to the Pythagorean magnitude of the transverse slip ($v_\perp^2 = v_x^2 + v_y^2$). This cancellation proves that the eigenstates are invariant under continuous transverse rotation, establishing $O(2)$ gauge symmetry:

$$(\alpha_s^{-2} - \lambda)\Big[(-\alpha_s^2 - \lambda)\big((\alpha_s^{-2} - \lambda)^2 + (\alpha t_p\Omega)^2\big) + \Big(\alpha\frac{v_\perp}{c}\Big)^2(\alpha_s^{-2} - \lambda)\Big] = 0 \tag{129}$$

This polynomial expansion identifies the independent physical states of the boundary via its constituent roots.

9.1 The Longitudinal Axis (λ_1)

Extracting the decoupled spatial root ($\lambda_1 = \alpha_s^{-2}$) demonstrates invariant volumetric expansion along the primary axis of vorticity. Because this root factors completely from the transverse polynomial, the rotational axis is orthogonally independent from transverse shear. This decoupled strain constitutes the unperturbed inertial core.

9.2 Electromagnetic Cross-Coupling Equivalence

The remaining cubic polynomial restricts the temporal and transverse spatial states. Factoring the unperturbed volumetric strain roots from the active kinematic fields yields the following equality:

$$(\alpha_s^2 + \lambda)(\alpha_s^{-2} - \lambda)^2 = \Big(\alpha\frac{v_\perp}{c}\Big)^2(\alpha_s^{-2} - \lambda) - (\alpha_s^2 + \lambda)(\alpha t_p\Omega)^2 \tag{130}$$

The spatial eigenvalue ($\alpha_s^{-2} - \lambda$) scales the electric strain ($v_\perp/c$), while the temporal eigenvalue ($\alpha_s^2 + \lambda$) scales the magnetic strain ($t_p\Omega$). Solving this equality for transverse slip determines the governing ratio:

$$\Big(\alpha\frac{v_\perp}{c}\Big)^2 = (\alpha_s^2 + \lambda)(\alpha_s^{-2} - \lambda) + \frac{\alpha_s^2 + \lambda}{\alpha_s^{-2} - \lambda}(\alpha t_p\Omega)^2 \tag{131}$$

Variation in vorticity causes a proportional shift in the transverse slip, establishing the pointwise impedance required to support Maxwell's equations.

9.3 Orthogonal Stability (Eigenstate Degeneracy)

As the transverse roots tend towards the longitudinal root ($\lambda \to \alpha_s^{-2}$), the spatial multipliers tend to zero and the polynomial collapses to:

$$0 = -(\alpha_s^2 + \alpha_s^{-2})(\alpha t_p \Omega)^2 \tag{132}$$

Because the scalar capacity terms are positive, this equality is satisfied when vorticity evaluates to zero ($\Omega = 0$). The matrix excludes transverse (electromagnetic) roots from degenerating into the longitudinal spatial axis. This boundary preserves the orthogonal geometry of propagating waves, preventing spatial collapse provided that internal vorticity is present.

9.4 Classical Volume Preservation ($\alpha_s = 1$)

Evaluating the polynomial at full background capacity, where longitudinal dynamic pressure is zero ($P_{dyn} = 0 \implies \alpha_s = 1$), reduces the polynomial to:

$$(1+\lambda)(1-\lambda)^2 = \left(\alpha \frac{v_\perp}{c}\right)^2 (1-\lambda) - (1+\lambda)(\alpha t_p \Omega)^2 \tag{133}$$

Because the antisymmetric matrix is trace-free, transverse shear propagates as an isochoric (volume-preserving) strain that does not deplete the background capacity. To conserve the baseline Minkowski volume, the determinant of the perturbed matrix equates with the unperturbed determinant ($\det(\hat{U}) = \det(\eta) = -1$):

$$\det(\hat{U}_{\mu\nu}) = -1 - (\alpha t_p \Omega)^2 + \left(\alpha \frac{v_\perp}{c}\right)^2 = -1 \implies \left(\alpha \frac{v_\perp}{c}\right)^2 = (\alpha t_p \Omega)^2 \tag{134}$$

By structurally mapping the electric field ($E \propto v_\perp / c$) and magnetic field ($B \propto t_p \Omega$), this limit reproduces the impedance of free space (Z_0), and the proportional magnitude ($E = cB$).

9.5 Non-Linear Vacuum Polarization (The Schwinger Limit)

Classical electromagnetism assumes linear superposition, but the tensor restricts this to the weak-shear limit. For a coordinate free of longitudinal dynamic pressure ($\alpha_s = 1$) but subjected to a critical transverse shear limit (where transverse strain equals the volumetric strain, $\alpha t_p \Omega = 1$), the eigenvalues are forced into the complex plane ($\lambda = 1 \pm i$).

Substituting this eigenvalue into the pointwise impedance ratio from Section 9.2 determines the modulus of the vacuum:

$$|Z_{ratio}| = \left| \frac{1 + (1 \pm i)}{1 - (1 \pm i)} \right| = \frac{|2 \pm i|}{|\mp i|} = \sqrt{5} \tag{135}$$

Because the impedance ratio shifts from 1 to $\sqrt{5}$, classical $U(1)$ symmetry breaks. The tensor dictates that at this 45-degree shear limit, the vacuum becomes non-linear, providing a mechanism for vacuum birefringence at the Schwinger limit.

9.6 Continuous Unitary Evolution ($U(1)$ Symmetry)

Evaluating the characteristic polynomial at the real spatial ($\lambda = 1$) or real temporal ($\lambda = -1$) roots results in $\Omega = 0$ and $v_\perp = 0$, respectively.

Isolating the transverse spatial (2×2) sub-matrix from the primary tensor decomposes the strain into a symmetric scalar amplitude and an antisymmetric rotational generator:

$$\hat{U}_\perp = \begin{bmatrix} \alpha_s^{-2} & -\alpha t_p \Omega \\ \alpha t_p \Omega & \alpha_s^{-2} \end{bmatrix} = \alpha_s^{-2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \alpha t_p \Omega \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \tag{136}$$

A matrix possessing this structure ($a\mathbf{I} + b\mathbf{J}$, where $\mathbf{J}^2 = -\mathbf{I}$) is isomorphic to the complex plane ($a + ib$). The real scalar (α_s^{-2}) specifies the longitudinal volumetric amplitude. The antisymmetric generator ($\alpha t_p \Omega$) governs the rotational frequency.

Because the exponential of an antisymmetric generator structures the rotation group ($e^{\mathbf{J}\theta} = \cos\theta\mathbf{I} + \sin\theta\mathbf{J}$), the boundary operates as an acoustic standing wave. This connects the quantum mechanical $U(1)$ unitary phase rotation to the asymmetric strain components.

Evaluating the eigenvectors ($\mathbf{v}$) of this transverse sub-matrix for the complex conjugate roots ($\lambda = \alpha_s^{-2} \pm i\alpha t_p \Omega$) determines the spatial orientation of the strain:

$$\begin{bmatrix} \mp i\alpha t_p \Omega & -\alpha t_p \Omega \\ \alpha t_p \Omega & \mp i\alpha t_p \Omega \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = 0 \implies \mathbf{v}_{\pm} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ \mp i \end{bmatrix} \tag{137}$$

The geometric normalization of these orthogonal vectors scales by $1/\sqrt{2} = \cos(\pi/4)$. This projection supplies the 45-degree maximum Tresca shear plane required to yield the metric into a discrete mass boundary (Section 8.3). These complex orthogonal vectors recover the Jones vectors for circular polarization, deriving the gauge basis for spin-1 transverse wave propagation from the shear limits of the tensor.

9.7 The Generalized Determinant ($\det(\hat{U}) = 0$)

Setting the determinant to zero sets the condition where the 4D metric volume collapses:

$$\det(\hat{U}_{\mu\nu}) = -\alpha_s^{-4} - (\alpha t_p \Omega)^2 + \alpha_s^{-4} \left(\alpha \frac{v_\perp}{c}\right)^2 = 0 \tag{138}$$

Multiplying by the spatial capacity (α_s^4) isolates the hyperbolic boundary:

$$\left(\alpha \frac{v_\perp}{c}\right)^2 - \alpha_s^4 (\alpha t_p \Omega)^2 = 1 \tag{139}$$

This equality ($E^2 - \alpha_s^4 B^2 = 1$) unites distinct topological yield limits across the full evaluation of $\alpha_s \in [0, 1]$:

- **Hawking Radiation ($0 < \alpha_s < 1$):** As gravitational dynamic pressure increases, the magnetic term is suppressed by the fourth power of the scalar admittance (α_s^4). The topological boundary yields at exponentially lower energy thresholds, structuring spontaneous pair production near an event horizon.
- **The Kinematic Mass Limit ($\alpha_s \to 0$):** As dynamic pressure maximizes, the magnetic term vanishes ($\alpha_s^4 \to 0$). The boundary collapses when the transverse slip reaches the geometric limit ($E^2 = 1 \implies v_\perp = c/\alpha$). Because longitudinal causality restricts physical advection (group velocity) to $v_g < c$, this variable operates as the orthogonal phase velocity of the boundary ($v_p = c/\alpha$). This satisfies the de Broglie kinematic wave relation ($v_p \cdot v_g = c^2$) for a ground-state topological defect oscillating at $v_g = \alpha c$.

9.8 The Temporal Cavitation Limit ($\alpha_s \to 0$)

Evaluating the root conditions at the maximum thermodynamic yield capacity ($P_{dyn} = P_c \implies \alpha_s = 0$), the temporal multiplier ($-\alpha_s^2 - \lambda$) simplifies to ($-\lambda$). To satisfy the constraint, the temporal root equates to zero ($\lambda_0 = 0$).

This zero-root eliminates the time coordinate ($\hat{U}_{00} = 0$). Because the background capacity drops to zero, the coordinate space structurally freezes. The metric signature transitions from $(-,+,+,+)$ to $(0,+,+,+)$. Kinematic advection halts across this boundary, identifying both the macroscopic event horizon and the microscopic topological rest mass boundary.

9.9 Complex Strain & The Deborah Number

Separating the transverse spatial components from the primary tensor constructs a 2×2 submatrix ($\hat{U}_\perp$). Section 9.6 identifies this structure as isomorphic to the complex plane:

$$\hat{U}_\perp = \alpha_s^{-2}\mathbf{I} + \alpha t_p \Omega \mathbf{J} \equiv \alpha_s^{-2} + i(\alpha t_p \Omega) \tag{140}$$

This partitions the strain into two orthogonal bases:

- **Real Axis (Elastic Storage):** The symmetric diagonal (α_s^{-2}) quantifies longitudinal volumetric packing.
- **Imaginary Axis (Viscous Flow):** The antisymmetric generator ($i\alpha t_p \Omega$) tracks Clebsch vorticity.

1. Derivation of the Rheological Phase Angle
The ratio of the antisymmetric off-diagonals (viscous flow) to the symmetric spatial diagonals (elastic storage) determines the continuous rheological phase angle ($\tan\delta$):

$$\begin{aligned} \tan\delta &= \frac{\mathrm{Im}(\hat{U}_\perp)}{\mathrm{Re}(\hat{U}_\perp)} \\ \tan\delta &= \frac{\alpha t_p \Omega}{\alpha_s^{-2}} \\ \tan\delta &= \alpha_s^2 \alpha t_p \Omega \end{aligned} \tag{141}$$

2. The Deborah Number
The Deborah number (De) scales inversely with the loss tangent ($De \propto 1/\tan\delta$). The tensor extracts the Deborah number from the inverted ratio:

$$De = \frac{\mathrm{Re}(\hat{U}_\perp)}{\mathrm{Im}(\hat{U}_\perp)} = \frac{\alpha_s^{-2}}{\alpha t_p \Omega} = \frac{1}{\alpha_s^2 \alpha t_p \Omega} \tag{142}$$

This derivation governs the thermodynamic state of the vacuum metric:

- **Fluid Phase ($De \to 0$):** As internal vorticity increases or background capacity remains unperturbed ($\alpha_s \to 1$), the ratio approaches zero. The matrix models as a compliant fluid.
- **Glass Transition ($De \to \infty$):** As dynamic pressure increases ($P_{dyn} \to P_c$), scalar admittance collapses ($\alpha_s \to 0$). The denominator goes to zero, causing the ratio to diverge. The coordinate geometry hardens, demonstrating solid elastic behavior.

3. Complex Magnitude
The total strain applied to the boundary calculates as the magnitude of the complex vector:

$$|\hat{U}_\perp| = \sqrt{(\alpha_s^{-2})^2 + (\alpha t_p \Omega)^2} = \sqrt{\alpha_s^{-4} + \alpha^2 t_p^2 \Omega^2} \tag{143}$$

Because the structural components operate orthogonally in the complex plane, the metric sustains volumetric compression and rotational shear simultaneously. The tensor prevents infinite coordinate compression by distributing kinematic load across the imaginary axis.

4. The Kinematic Crossover Constraint
To establish the boundary, the Deborah number is derived from the parameters of the matrix. The Deborah number represents the ratio of transverse coupling capacity to proper temporal compliance.

In the asymmetric tensor, the transverse capacity is bounded by the Viscoelastic Modulus Ratio (α). The proper temporal compliance is given by the square root of the time-time diagonal ($\sqrt{-\hat{U}_{00}} = \alpha_s$). Dividing these tensor components formulates the macroscopic kinematic boundary:

$$De_{macro} = \frac{\alpha}{\sqrt{-\hat{U}_{00}}} = \frac{\alpha}{\alpha_s} \tag{144}$$

Because the inverse scalar admittance equates to the Lorentz factor ($\alpha_s^{-1} = \gamma$), this substitution produces the time-dilated kinematic limit:

$$De_{macro} = \alpha\gamma \tag{145}$$

Setting this macroscopic boundary (De_{macro}) equal to the localized extraction (Equation 142) produces:

$$\begin{aligned} \frac{\alpha}{\alpha_s} &= \frac{\alpha_s^{-2}}{\alpha t_p \Omega} \\ \alpha^2 (t_p \Omega) &= \alpha_s^{-1} \\ t_p \Omega &= \frac{1}{\alpha^2 \alpha_s} = \frac{\gamma}{\alpha^2} \end{aligned} \tag{146}$$

This intersection dictates the physical scaling of the topological boundary. As dynamic pressure saturates the metric capacity ($P_{dyn} \to P_c$), scalar admittance collapses ($\alpha_s \to 0$). This substitution requires that the normalized internal Clebsch vorticity diverges proportionally to the Lorentz factor ($\gamma \to \infty$). The metric maintains the constraint ($\nabla \cdot \mathbf{v} = 0$) by transferring infinite longitudinal compression into unbounded transverse rotational shear.

5. The Phenomenological Energy Limit

In relativistic kinematics, the total energy (E) of a boundary increases from its unperturbed rest mass energy ($E_0 = mc^2$) via the Lorentz factor:

$$\gamma = \frac{E}{mc^2} \tag{147}$$

Substituting this energy ratio into the macroscopic boundary ($De_{macro} = \alpha\gamma$) translates the Deborah number into a thermodynamic energy scale:

$$De = \alpha \left(\frac{E}{mc^2} \right) \tag{148}$$

This formulation converts the transverse shear of the complex tensor into the observable kinematic energy threshold of the particle.

10. Tensor Covariance & Non-Inertial Reference Frames

Extending the boundary states defined in Section 8, the metric maintains structural equilibrium across relative observer geometries. Comparing states ($\hat{U}_{\mu\nu}$) across arbitrary reference frames reveals the coordinate transformations between separate observers.

10.1 Kinematic Classification of Observer Frames

Translating the tensor from a resting observer to a transformed observer decomposes into two mechanical categories: Scalar Translation and Tensor Covariance.

1. Scalar Translation (Gradient Isomorphism):

Comparisons between stationary observers separated by a thermodynamic gradient resolve without kinematic shear. Because the matrix defines macroscopic gravity as the static pressure

gradient (∇P_{static}), comparing distinct gravitational reference frames operates via coordinate translation. The transformation matrices reduce to the Kronecker delta ($\Lambda^{\mu}_{\alpha} = \delta^{\mu}_{\alpha}$). Modifying the observer coordinate translates the matrix along the continuous gradient, shifting the spatial and temporal diagonals proportionately via the scalar admittance (α_s^2). This coordinate transformation governs stationary scalar variance (comparing gravitational elevations or distinct cosmological Hubble epochs), requiring no internal matrix cross-coupling.

2. Tensor Covariance (Kinematic Mixing):
Comparisons where one observer possesses relative velocity, applied rotation, or transverse shear demand matrix transformation. The tensor establishes equivalence between the observers via the Lorentz-Jacobian transformation matrix ($\Lambda^{\mu}_{\alpha} = \partial x^{\mu}/\partial x'^{\alpha}$) [38]:

$$\hat{U}'_{\alpha\beta} = \Lambda^{\mu}_{\alpha}\Lambda^{\nu}_{\beta}\hat{U}_{\mu\nu} \tag{149}$$

This operation directs the symmetric diagonals to cross-multiply against the antisymmetric transverse shear components, extracting the mechanics of relativistic and fictitious forces.

10.2 Active Kinematic Strain & The Invariant Phase Limit

When a topological boundary translates through the baseline metric, it generates a convective flow field (v_{rel}). Rather than a passive coordinate rotation, it represents active physical strain.

The convective velocity subjects the translating boundary to dynamic pressure ($P_{dyn} = \frac{1}{2}\rho_\tau v_{rel}^2$). By Bernoulli conservation [31] ($P_{static} = P_c - P_{dyn}$), the temporal diagonal ($\hat{U}_{00} = -\alpha_s^2$) yields the structural limit:

$$\begin{aligned}
\hat{U}'_{00} &= -\left(\frac{P_c - P_{dyn}}{P_c}\right) \\
\hat{U}'_{00} &= -\left(1 - \frac{\frac{1}{2}\rho_\tau v_{rel}^2}{\frac{1}{2}\rho_\tau c^2}\right) \\
\hat{U}'_{00} &= -\left(1 - \frac{v_{rel}^2}{c^2}\right)
\end{aligned} \tag{150}$$

As the relative velocity of the boundary approaches the acoustic limit ($v_{rel} \to c$), the dynamic pressure saturates ($P_{dyn} \to P_c$). The temporal diagonal collapses to zero ($\hat{U}'_{00} \to 0$), driving the symmetric spatial diagonals to diverge ($S'_{ii} \to \infty$).

Continuous temporal dilation and the invariant speed of light emerge from this operation. The tensor thus reveals Lorentz contraction as an active thermodynamic response which actively conserves the maximum static pressure capacity.

10.3 Rotational Covariance & Transverse Leakage

Comparing a stationary inertial observer to a rotating observer introduces multi-axis shear. For an observer rotating at angular velocity ω, the azimuthal coordinate transformation ($d\phi = d\phi' + \omega dt'$) defines the temporal Jacobian component [29]:

$$\Lambda^{\phi}_{0} = \frac{\partial \phi}{\partial t'} = \omega \tag{151}$$

Applying the transformation matrix (149), to evaluate the space-time off-diagonal ($\hat{U}'_{0i}$) in the rotating frame, produces the cross-coupling mechanism:

$$\begin{aligned}
\hat{U}'_{0i} &= \Lambda^{\alpha}_{0}\Lambda^{\beta}_{i}\hat{U}_{\alpha\beta} \\
\hat{U}'_{0i} &= \Lambda^{0}_{0}\Lambda^{\beta}_{i}\hat{U}_{0\beta} + \Lambda^{\phi}_{0}\Lambda^{\beta}_{i}\hat{U}_{\phi\beta} \\
\hat{U}'_{0i} &= (1)\Lambda^{\beta}_{i}\hat{U}_{0\beta} + (\omega)\Lambda^{\beta}_{i}\hat{U}_{\phi\beta}
\end{aligned} \tag{152}$$

The transformation causes the space-space off-diagonals (vorticity, $\hat{U}_{\phi\beta}$) to influence the space-time off-diagonals (transverse velocity, $\hat{U}'_{0i}$), scaled by the rotation rate (ω). The rotating coordinate frame results in apparent transverse velocity fields (Coriolis and Centrifugal strain) that resolve to zero for the resting frame. Fictitious forces manifest as the mechanical artifact of internal Clebsch vorticity across the rotating spatial boundary.

10.4 Conformal Scale Variance (The Micro-Macro Asymmetry)

The physical structure of the continuum breaks scale invariance. While the metric appears identical across arbitrary macroscopic scales, the generation of discrete topological mass (Section 8) sets a fixed physical boundary dimension ($r_c = \hbar/mc$).

Applying the dilation vector field ($\xi^\mu = \sigma x^\mu$) to compare a microscopic rest frame to a macroscopic observer exposes the broken symmetry. The Lie derivative resolves to a conformal factor ($\mathcal{L}_\xi \hat{U}_{\mu\nu} = 2\sigma \hat{U}_{\mu\nu}$), dictating that macroscopic scaling alters topology.

- **The Microscopic Observer:** The discrete spatial incompatibility of the topological boundary generates a structural gradient ($\sigma x^\alpha \partial_\alpha \hat{U}_{\mu\nu} \neq 0$), breaking conformal scaling.
- **The Macroscopic Observer:** Integrating the static pressure gradient across a macroscopic volume ($V \gg r_c^3$) causes the transverse shear components (A_{ij}) to sum to zero, homogenizing the gradient ($\int \partial_\alpha \hat{U}_{\mu\nu}\, dV = 0$) and recovering macroscopic scale invariance:

$$\iiint_V A_{ij}\, dV = 0 \implies \text{Metric Homogenization} \tag{153}$$

The phase rotations of the microscopic tensor homogenize over expanded integration volumes, recovering the macroscopic dynamics of General Relativity. This shows how invariance is distinct from the topological scale variance across the boundary.

10.5 The Equivalence Principle (Acceleration Isomorphism)

The equivalence principle postulates the observational equivalence between a uniform gravitational field and a uniformly accelerating reference frame [15]. The tensor derives this equivalence not as an independent assumption, but as a consequence of comparing distinct observer frames.

Evaluating the volumetric trace deformation ($\delta \hat{U}_{00} \propto \nabla P$) for both observers recovers the mechanical strain:

- **The Gravitational Observer:** A stationary frame submerged in a metric depression is governed by the ambient static pressure gradient (∇P_{static}).
- **The Accelerating Observer:** A non-inertial frame navigating a flat macroscopic vacuum increases its convective velocity. Substituting this kinematic load into the macroscopic metric generates a dynamic pressure gradient (∇P_{dyn}).

Because the continuous metric conserves total thermodynamic capacity ($P_{static} = P_c - P_{dyn}$):

$$\nabla P_{static} = -\nabla P_{dyn} \implies \mathbf{a}_{grav} = -\mathbf{a}_{kin} \tag{154}$$

Increasing the dynamic load extracts phase capacity from the identical structural ledger as an ambient gravitational deficit. Applying a continuous kinematic acceleration vector (∇P_{dyn}) and translating across a static gravitational potential (∇P_{static}) induce the identical volumetric deformation upon the Tensor.

10.6 Discrete Frame Inversion (CPT Symmetry & Antimatter)

Comparing a forward-time or spatially inverted observer partitions the discrete transformation invariants of the metric.

Part I: Time-Reversal (T-Symmetry)

Applying the time-reversal transformation matrix (Λ_T) inverts the temporal coordinate ($t \to -t$). The matrix is defined as $\Lambda_T = \text{diag}(-1, 1, 1, 1)$ [54]. Applying this operation to the tensor partitions orthogonal responses:

1. Symmetric Invariance (Mass/Gravity):

Transforming the temporal diagonal ($\hat{U}_{00}$) via the time-reversal matrix:

$$\hat{U}'_{00} = \Lambda^{\mu}_{0}\Lambda^{\nu}_{0}\hat{U}_{\mu\nu} = (-1)(-1)\hat{U}_{00} = \hat{U}_{00} \tag{155}$$

Because the negative inversion squares, the scalar diagonal is preserved. The inverted coordinate frame produces identical longitudinal inertia and gravitational attraction.

2. Antisymmetric Inversion (Kinematic Shear):

Transforming the space-time off-diagonal ($\hat{U}_{0i}$) defining transverse slip yields:

$$\hat{U}'_{0i} = \Lambda^{\mu}_{0}\Lambda^{\nu}_{i}\hat{U}_{\mu\nu} = (-1)(1)\hat{U}_{0i} = -\hat{U}_{0i} \tag{156}$$

The temporal index flips the sign of the transverse shear. The inverted coordinate frame outputs an identical phase with reversed transverse slip.

Executing these transformation operations verifies Time-Reversal (T) symmetry within the continuum.

Part II: Spatial Inversion (P-Symmetry) & Charge Conjugation (C)

To evaluate spatial Parity (P) and Charge Conjugation (C), the discrete spatial inversion matrix (Λ_P) is applied to the metric. The matrix inverts the spatial coordinates ($x^i \to -x^i$) and is defined as $\Lambda_P = \text{diag}(1, -1, -1, -1)$.

3. Volumetric Invariance (Mass/Pressure):

Transforming the spatial diagonals ($\hat{U}_{ii}$) yields:

$$\hat{U}'_{ii} = \Lambda^{\mu}_{i}\Lambda^{\nu}_{i}\hat{U}_{\mu\nu} = (-1)(-1)\hat{U}_{ii} = \hat{U}_{ii} \tag{157}$$

The symmetric spatial packing remains positive. The inverted coordinate frame produces an identical mass density and gravitational gradient.

4. Axial Inversion (Chirality & Parity):

Transforming the antisymmetric spatial off-diagonals ($\hat{U}_{ij}$) yields:

$$\hat{U}'_{ij} = \Lambda^{\mu}_{i}\Lambda^{\nu}_{j}\hat{U}_{\mu\nu} = (-1)(-1)\hat{U}_{ij} = \hat{U}_{ij} \tag{158}$$

While the spatial rotation is preserved as an axial vector, the spatial inversion of the coordinate axes reverses the chirality (geometric handedness) of the Clebsch vortex. The topological defect keeps its original flow magnitude but flips its spatial twist.

5. Polar Inversion (Winding Number & Charge):

Transforming the space-time off-diagonal ($\hat{U}_{0i}$) yields:

$$\hat{U}'_{0i} = \Lambda^{\mu}_{0}\Lambda^{\nu}_{i}\hat{U}_{\mu\nu} = (1)(-1)\hat{U}_{0i} = -\hat{U}_{0i} \tag{159}$$

The temporal-spatial cross term inverts as a polar vector. While classical kinematics treats this as boundary velocity ($v \to c$), this parameter acts as the directionality of the Clebsch phase. Charge polarity operates as the topological winding number of this boundary.

Because the metric preserves the axial rotation (U_{ij}) but inverts the transverse directionality (U_{0i}), the geometric handedness of the winding number is reversed. This permits spatial Parity inversion (P) to map to Charge Conjugation (C) for an isolated defect.

For the proposed direction for deriving CP violation, please see section 16.2.

11. Galactic Kinematics and the Cosmological Boundary

The continuous metric transitions from a collection of mass defects to the thermodynamic boundary of the expanding macroscopic background. Computing the total dynamic pressure state of the continuum secures a bounded geometric baseline for galactic kinematics.

11.1 The Total Dynamic Pressure State

The dynamic pressure depleted by a baryonic aggregate (mass) is parameterized as $P_{bar} = -\rho_\tau \Phi(r)$, where $\Phi(r)$ is the classical gravitational potential. Simultaneously, the continuum sustains the macroscopic load of longitudinal cosmological expansion. Attenuated by the causal horizon, this flow saturates exponentially: $v_0(r) = c(1 - e^{-r/R_h})$. The kinetic load equates to $P_{cosmo} = \frac{1}{2}\rho_\tau c^2(1 - e^{-r/R_h})^2$.

Because thermodynamic pressures superimpose as scalars, the total dynamic pressure sums to:

$$P_{dyn}(r) = -\rho_\tau \Phi(r) + \frac{1}{2}\rho_\tau c^2 \left(1 - e^{-r/R_h}\right)^2 \tag{160}$$

11.2 The Bounded Kinematic Baseline

In a continuous metric, orbital velocity responds to the spatial gradient of the dynamic pressure state. To maintain a circular orbit, inward centripetal acceleration balances the net continuum acceleration gradient ($g = -\frac{1}{\rho_\tau}\frac{\partial P_{dyn}}{\partial r}$). The kinematic state is modeled by the radial differential operator:

$$v^2 = -r\left(\frac{1}{\rho_\tau}\frac{\partial P_{dyn}}{\partial r}\right) \tag{161}$$

Executing the radial derivative across Equation 160 outputs the kinematic baseline of the continuum:

$$v_{base}^2(r) = r\frac{\partial \Phi}{\partial r} - \frac{c^2 r}{R_h}e^{-r/R_h}\left(1 - e^{-r/R_h}\right) \tag{162}$$

This derived baseline bounds the kinematic domain, correcting the coordinate divergences of the classical Schwarzschild-de Sitter (SdS) metric:

- **The Core Bound ($r \to 0$):** Inside a uniform baryonic bulge, the interior potential spatial derivative yields an outward gradient $\frac{\partial \Phi}{\partial r} \propto r$. Substituting this into the operator produces $v_{base}^2 \propto r^2$. As the cosmological gradient attenuates at local scales, the baseline reproduces the positive solid-body rotation profile ($v_{base} \propto r$) without generating imaginary roots.
- **The Finite Boundary (The Hill Sphere):** For exterior domains, the potential transitions to $\Phi_{out} = -GM/r$. Equation 162 evaluates to a finite root where inward baryonic acceleration equals outward cosmological advection ($v_{base}^2 = 0$). This marks the Jacobian capture radius, terminating the coordinate frame for bound rotational orbits.
- **The Macroscopic Asymptote ($r \to \infty$):** As the radial coordinate extends toward the macroscopic causal boundary, the longitudinal expansion velocity saturates ($v_0 \to c$). When the dynamic pressure maximizes to the capacity limit ($P_{dyn} \to P_c$) the orbital requirement drops to zero ($v_{base}^2 \to 0$). Geometric expansion is asymptotically inertial.

11.3 Isolating the Kinematic Discrepancy

The continuum requires a Keplerian decline terminating at a finite detachment limit. To map the geometric requirements of the missing mass anomaly, the unified kinematic state is formulated

as an empirically informed *ansatz*, rearranged to isolate the effective kinematic discrepancy (Γ_{anom}):

$$\Gamma_{anom}(r) = v_{obs}^2(r) - \left[\frac{GM_{bar}(r)}{r} - \frac{c^2 r}{R_h} e^{-r/R_h}\left(1 - e^{-r/R_h}\right)\right] \tag{163}$$

Where $v_{obs}^2(r)$ represents the empirical rotational velocity field. This separation bounds the structural parameters the anomaly must fulfill:

- **The Core Attenuation ($r \to 0$):** While High Surface Brightness (HSB) spirals exhibit baryon-dominated cores, Low Surface Brightness (LSB) and dwarf galaxies demand a non-zero anomaly at the origin. However, across all morphologies, empirical kinematics ($v_{obs}^2 \propto r^2$) require the enclosed anomalous mass to scale geometrically as $M_{anom}(r) \propto r^3$. Taking the spatial derivative dictates a constant volumetric density at the origin ($\rho \to$ const). Singular gravitational cusps are structurally prohibited; the anomaly is bounded to a flat-cored spatial profile.

- **The Effective Halo Plateau ($r \gg R_c$):** In the macroscopic halo, the observed rotation curve forms a plateau ($v_{obs}^2 \approx v_{flat}^2$). The ansatz isolates the discrepancy as $\Gamma_{anom}(r) \approx v_{flat}^2 - \frac{GM_{bar}(r)}{r}$. Sustaining this plateau requires a supplementary, positive dynamic pressure field scaling logarithmically ($P_{anom} \propto \ln(R_{max}/r)$) to dominate the decaying baryonic gradient.

- **The Detachment Limit (The Hill Sphere):** As established in Section 11.2, bound rotational orbits physically terminate at the Jacobian capture radius where total acceleration resolves to zero. Consequently, the logarithmic pressure well cannot extend to spatial infinity. The structural anomaly must truncate at this macroscopic coordinate boundary.

11.4 The Universal No-Go Theorem

A physical resolution to Γ_{anom} is constrained by two mechanical boundaries: it requires a supplementary pressure well ($P_{anom} \propto \ln(R_{max}/r)$) to provide a constant kinematic plateau, and its amplitude requires coupling to baryonic mass values to satisfy the Baryonic Tully-Fisher Relation ($M_{bar} \propto v_{flat}^4$).

By combining continuous geometric limitations with the causal coupling constraints of the ansatz, a Universal No-Go Theorem is constructed, falsifying current standard models:

- **Falsification of Modified Gravity (The Geometric Constraint):** In a continuous, empty vacuum topology ($\rho_{dyn} = 0$), the static pressure field satisfies Laplace's equation ($\nabla^2 P_{dyn} = 0$), permitting the Newtonian monopole solution in 3D space. Applying the 3D spherical Laplacian operator to a logarithmic well computes to $\nabla^2(-\ln r) = -1/r^2 \neq 0$. Utilizing non-linear modified operators violates the foundational axiom of the continuous tensor, which dictates that scalar dynamic pressures superimpose linearly.

- **Falsification of ΛCDM (The Causal Constraint):** While standard particulate mass (ΛCDM) supplies the required independent $1/r^2$ halo density, collisionless kinematic halos form hierarchically and independently of baryonic infall. They lack a dynamic thermodynamic mechanism to couple the required macroscopic logarithmic well to the baryonic mass, failing the causal constraints of the BTFR.

11.5 The Phenomenological Envelope

Consequently, the framework demonstrates that the missing mass anomaly cannot be a geometric illusion of empty space, nor can it be a decoupled primordial particulate halo. By restricting

convergence on existing standard models, the asymmetric tensor bounds the spacetime continuum.

To resolve the kinematic plateau, empirical observations specify that the continuum requires a structural modification exhibiting the following profile:

1. **Spatial Pressure Gradient:** A supplementary dynamic pressure field ($P_{anom} \propto \ln(R_{max}/r)$) generating the kinematic equivalent of a $1/r^2$ structural mass dispersion.
2. **Causal Coupling:** The amplitude of this pressure gradient requires a causal generation mechanism linked directly to the baryonic mass.
3. **Inertial Decoupling:** The gradient requires independent kinematic inertia to detach and coast ballistically during macroscopic cluster collisions.

The continuous metric derives the classical limit of General Relativity and extracts the phenomenological envelope of the kinematic anomaly. By satisfying the kinematic requirement for a $1/r^2$ structural mass dispersion while prohibiting the uncoupled variance of primordial ΛCDM particles, Dark Matter resolves as a dynamically spawned structural pressure gradient forged by thermodynamic strain, rather than a primordial cosmological remnant. The differential operators required to convert a centralized baryonic potential into this independent, stable pressure distribution remains an open problem for future works.

11.6 Proposed Resolution: Elastic Metric Tension (Scale-Invariant Vorticity Coupling)

As a resolution to satisfy the kinematic envelope without violating the density constraints, the rotational shear of the galactic disk translates as a continuous elastic tension upon the metric grid.

1. Microscopic Vorticity & The Baryon Count (N):
As derived in Section 8, a localized mass generates a cavitation boundary governed by internal vorticity ($\mathbf{\Omega}_{pr}$). The total baryonic mass (M_{bar}) of a galactic disk aggregates these localized defects ($N = M_{bar}/m_{pr}$).

2. Strain Polarization & Dimensional Reduction:
The macroscopic state of the metric is defined by the volumetric expectation value of the aggregate micro-states. Total thermodynamic randomness results in macroscopic cancellation ($\langle A_{\mu\nu} \rangle_V = 0$).

However, in a fully continuous coordinate space, macroscopic boundary conditions back-react upon microscopic states. A net galactic angular momentum ($\mathbf{J} \neq 0$) relative to the cosmic background applies a global rotational strain across the continuum. This global strain breaks the thermodynamic degeneracy of the localized defects, making alignment with the bulk rotation energetically preferential.

Consequently, the thermal fluctuations of the baryonic vorticity cannot achieve perfect isotropic cancellation. The global rotation forces a fractional alignment bias, establishing a non-zero macroscopic expectation value for the antisymmetric spatial tensor:

$$\langle A_{ij} \rangle = \frac{1}{V} \int_V A_{ij}^{(micro)} dV \neq 0 \tag{164}$$

Because this coherent residual is dictated by the conserved angular momentum ($\mathbf{J}$), the continuous medium accommodates the aggregate torque as a steady-state axial field. By definition, this steady-state rotation restricts the tensor macroscopically, rendering it translationally invariant along the rotational axis ($\partial_z \langle A_{ij} \rangle = 0$).

This translational invariance zeroes the off-plane shear components ($A_{xz} = A_{yz} = 0$). To accommodate this restricted state, the symmetric stress tensor decouples the axial component ($\sigma_{zz} = 0$).

Through the conservation of angular momentum, the macroscopic kinematic response of the continuum is restricted to a 2D cylindrical plane stress state (r, ϕ).

3. The 2D Laplacian & Logarithmic Shear:
The trace contraction of the tensor $(\hat{U}_{\mu\nu}\hat{U}^{\mu\nu})$ divides the kinetic energy. The antisymmetric trace filters the transverse shear stress $(P_{shear} = \frac{1}{4}P_c A_{\mu\nu}A^{\mu\nu})$. In a steady-state continuous medium, this scalar pressure field is bounded by the source-free Laplace equation $(\nabla^2 P_{shear} = 0)$. Applying the 2D polar Laplacian specifies the differential operator:

$$\nabla_{2D}^2 P_{shear} = \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial P_{shear}}{\partial r}\right) = 0 \tag{165}$$

Integrating this operator radially outward constructs the macroscopic scalar pressure field:

$$P_{shear}^{2D}(r) = k\ln\left(\frac{R_{max}}{r}\right) \tag{166}$$

The metric generates a logarithmic tension well, where the scalar amplitude (k) represents the net shear load of the galaxy.

4. The Elastic Metric Tension:
By the conservation laws of the thermodynamic budget, this shear load acts as an active transverse tension upon the Total Scalar Admittance (α_s^2). The symmetric temporal diagonal $(S_{00} = -\alpha_s^2)$ shifts to accommodate both the 3D isotropic Newtonian gravity and polarized 2D shear:

$$S_{00} = -\left(1 - \frac{P_{dyn}^{3D} + P_{shear}^{2D}}{P_c}\right) \tag{167}$$

5. Geodesic Gradient Evaluation:
The coordinate acceleration of a test mass evaluates via the spatial gradient of the time-time diagonal in the weak-field geodesic equation $(a^i \approx \frac{1}{2}c^2\nabla^i S_{00})$. Inserting the admittance components translates the scalar pressure loads into geometric curvature:

$$a^i = \frac{c^2}{2P_c}\left(\nabla P_{dyn}^{3D} + \nabla P_{shear}^{2D}\right) \tag{168}$$

Because the spatial gradient of a pressure well points inward $(\nabla P \propto -\hat{\mathbf{r}})$, taking the derivatives of both the 3D potential $(\rho_\tau G M_{bar}/r)$ and the 2D logarithmic shear $(k\ln(R_{max}/r))$ yields:

$$a^i = \frac{c^2}{2P_c}\left(-\frac{\rho_\tau G M_{bar}}{r^2} - \frac{k}{r}\right)\hat{\mathbf{r}} = \left(\frac{GM_{bar}}{r^2} + \frac{c^2 k}{2P_c r}\right)(-\hat{\mathbf{r}}) \tag{169}$$

Both components pull inward. The 3D component retrieves the Keplerian centripetal bound, while the 2D polarized component supplies the $1/r$ inward acceleration.

6. Macroscopic Plug Flow (Kinematic Plateau Extraction):
Equating the anomalous 2D spatial component to the required inward centripetal acceleration limit (v_{rot}^2/r) cancels the radial coordinate:

$$\frac{v_{rot}^2}{r} = \left(\frac{c^2}{2P_c}\right)\frac{k}{r} \implies v_{rot} = c\sqrt{\frac{k}{2P_c}} \tag{170}$$

The orbital velocity squares to a constant kinematic plateau. In Bingham fluid dynamics, this region of constant velocity and zero internal dynamic shear represents a macroscopic plug flow. The continuum moves as a unified bulk, sustained by the elastic tension of the vacuum (P_c).

7. The Three-Layer Rheological Coupling & The Macroscopic Yield Limit:
The shear amplitude (k) evaluates via the mechanical energy balance. The macroscopic yield limit (a_0) is derived from the divergence equilibrium of the tensor components.

Transverse slip is parameterized by the space-time off-diagonal ($\hat{U}_{0i} = -\alpha v_{\perp,i}/c$). The spatial gradient of this component constructs the rotational acceleration ($\partial_j \hat{U}_{0i} \propto a_{rot}/c^2$). Simultaneously, macroscopic cosmological expansion applies a continuous temporal divergence to the volumetric spatial diagonals ($\partial_0 \hat{U}_{ii} \propto H_0/c$).

To maintain a dynamic slipping state, the applied transverse shear gradient must exceed the baseline temporal strain rate ($\partial_j \hat{U}_{0i} > \partial_0 \hat{U}_{ii}$). Equating these orthogonal derivatives isolates the macroscopic kinematic yield threshold ($a_{rot} = cH_0$). Mapping this 1D radial temporal strain onto the 2D azimuthal plane of the galactic disk computes the rheological limit: $a_0 = cH_0/2\pi$.

This equilibrium constructs a bounded, three-layer architecture for the galactic metric:

Zone 1: *The Inner Shear Layer.* In the inner regions of the galaxy, the transverse spatial gradient overpowers the temporal expansion threshold ($\partial_j \hat{U}_{0i} > \partial_0 \hat{U}_{ii}$). The coordinate layers slip against one another, maintaining a yielding free rotational state governed by Newtonian gravity ($1/r^2$).

Zone 2: *The Trapped Plug Flow (The 3D → 2D Coupling).* As radial acceleration decays, the spatial gradient intersects the temporal limit ($\partial_j \hat{U}_{0i} = \partial_0 \hat{U}_{ii}$). At this transition radius (r_t), the local gravitational shear equals the cosmological baseline strain.

By the continuity equations, when the transverse gradient cannot exceed the longitudinal divergence, the exterior derivative of the 3D dynamic slip evaluates to zero ($d\mathbf{v} = 0$). Because the applied stress is insufficient to cause a local geometric yield, the torque cannot be dissipated via local rotation. Instead, this unresolved stress propagates outward as static structural strain.

To conserve angular momentum across the boundary without dynamic slip, the symmetric stress tensor (σ_{ij}) decouples the axial component ($\sigma_{zz} = 0$), forcing the continuum into a state of plane stress. This condition collapses the kinematic geometry from a 3D isotropic divergence into a 2D cylindrical stress field (r, ϕ). Mechanically, this 2D planar stress bounding a 3D fluid volume operates as macroscopic surface tension. The metric locks into a coherent elastic membrane, satisfying the Bingham plug flow condition (Equation 170).

This outward propagation behaves as a continuous stress integrated over time—a sustained structural push across the coordinate grid. Consequently, the local coordinate frame ($\hat{U}_{\mu\nu}$) rotates with the aggregate mass. The outer stars do not experience centrifugal coordinate acceleration ($d^2x^i/d\tau^2 = 0$); they are embedded at rest within this rotating surface tension. The coordinate space itself carries the rotational momentum outward until it reaches the terminal boundary.

Isolating this inner acceleration boundary identifies the transition radius:

$$\frac{GM_{bar}}{r_t^2} = a_0 \implies r_t = \sqrt{\frac{GM_{bar}}{a_0}} \tag{171}$$

Zone 3: *The Outer Shear Layer (The Terminal Boundary).* The 2D plug flow does not extend infinitely. While the inner boundary (r_t) evaluates via an acceleration match, the outer boundary evaluates via a velocity match. The background universe undergoes 3D volumetric expansion, where outward radial velocity increases with distance ($v_{exp} = H_0 r$). At the terminal Jacobian Capture Radius (R_{max}), the outward expansion velocity of the universe equals the constant rotational velocity of the galactic plug flow:

$$H_0 R_{max} = v_{rot} \implies R_{max} = \frac{v_{rot}}{H_0} \tag{172}$$

At this radial limit, the kinematic slip of the galaxy matches the volumetric divergence of the cosmic background. The 2D planar tension of the trapped plug flow shears against the isotropic tension of the surrounding void. The metric decouples across this phase-locked boundary, releasing the rotational strain back into the quiescent 3D expansion of the intergalactic medium.

Within the plug flow (Zone 2), the kinematics couple to the 2D tension. The rotational velocity conforms to the yield acceleration at the inner transition interface ($v_{rot}^2/r_t = a_0 \implies$

$v_{rot}^2 = a_0 r_t$). Substituting this transition radius into the kinematic plateau extracts the scaling equivalent to the net shear amplitude:

$$v_{rot}^2 = a_0 \sqrt{\frac{GM_{bar}}{a_0}} = \sqrt{a_0 GM_{bar}}$$
$$v_{rot}^4 = a_0 GM_{bar} \tag{173}$$

The Baryonic Tully-Fisher relation emerges as the kinematic consequence of this Bingham plug flow.

8. The Macroscopic Resolution (Bypassing the No-Go Theorem):
To verify the physical viability of the metric degradation, the strain evaluates against macroscopic measurements. For a galaxy exhibiting a flat rotational velocity of $v_{rot} \approx 200$ km/s, the ratio of the continuous shear pressure to the baseline vacuum stiffness (P_c) is:

$$\frac{P_{shear}^{2D}}{P_c} = \frac{v_{rot}^2}{c^2} \approx \left(\frac{2 \times 10^5 \text{ m/s}}{3 \times 10^8 \text{ m/s}} \right)^2 \approx 4.4 \times 10^{-7} \tag{174}$$

The anomalous behavior registers as a continuous 10^{-7} macroscopic elastic strain upon the coordinate grid.

This 10^{-7} magnitude quantifies the amplitude of the plug flow. Because the asymmetric tensor couples transverse rotation to electromagnetic gauge fields (Section 9), this locked mechanical strain manifests as an orthogonal electromagnetic signature. Substituting a macroscopic kinematic strain of 10^{-7} into the antisymmetric spatial off-diagonals ($\hat{U}_{ij}$) evaluates to the micro-Gauss magnetic fields observed across spiral galaxies.

By routing the rotational shear back into the symmetric sector, the continuum accommodates the load. Flat rotation curves and galactic magnetic dynamos are generated as orthogonal projections of the macroscopic plug flow without requiring non-baryonic particulate mass.

9. Observational Lensing Predictions (Vacuum Photoelasticity):
If the anomalous mass constitutes a continuous 2D elastic tension field rather than a 3D cloud of spherical particles (ΛCDM), the gravitational lensing signature diverges. Lensing operates as vacuum photoelasticity (stress birefringence), bounded by the refractive index of the vacuum (the temporal diagonal, S_{00}). Because the P_{shear}^{2D} tension is cylindrical, photons passing through the galactic plane (viewing an edge-on galaxy) traverse a distinct gradient of metric strain compared to photons passing parallel to the z-axis (viewing a face-on galaxy). Consequently, weak lensing data from current generation surveys (e.g., the Euclid space telescope or the Rubin Observatory LSST) should observe a statistical bias in the shear amplitude based on the inclination angle of the foreground galaxy, and future CMB interferometric surveys should capture the gradient.

Because this 2D tension field acts as plug flow trapped between the inner galactic shear and the outer cosmic expansion, it terminates at the terminal Jacobian Capture Radius (R_{max}) where the plug flow shears against the background. This establishes that the weak lensing profile will not asymptotically fade into the background as predicted by collisionless dark matter halos. Instead, it exhibits a discontinuity behaving as an outer shear ring, where the optical strain abruptly dissipates.

To directly map the macroscopic metric capacity without relying on stellar kinematics, high-resolution Cosmic Microwave Background (CMB) interferometry offers a pathway for validation. Because the tensor dictates that the speed of light is locally retarded by dynamic pressure ($v_c = c\alpha_s^2$), the CMB wavefront undergoes phase-shifting as it propagates through galactic strain gradients. Until next-generation interferometry can achieve the spatial resolution required to measure this pixelated index of refraction directly, wide-field weak lensing surveys remain the primary mechanism for isolating the 2D geometric discontinuity.

12. The Non-Linear Affine Connection (Einstein-Cartan Isomorphism)

The asymmetric strain tensor ($\hat{U}_{\mu\nu} = S_{\mu\nu} + A_{\mu\nu}$) requires an affine connection ($\Gamma^{\lambda}_{\mu\nu}$) that propagates both longitudinal volumetric deformation and transverse rotational shear across the continuum.

1. Connection Decomposition:
The generalized affine connection separates into a symmetric Levi-Civita component ($\tilde{\Gamma}^{\lambda}_{\mu\nu}$) and an antisymmetric contortion tensor ($K^{\lambda}_{\mu\nu}$):

$$\Gamma^{\lambda}_{\mu\nu} = \tilde{\Gamma}^{\lambda}_{\mu\nu} + K^{\lambda}_{\mu\nu} \tag{175}$$

2. The Symmetric Connection (Gravitational Acceleration):
The Levi-Civita connection is formulated from the symmetric diagonals ($S_{\mu\nu}$). Expanding the temporal component relative to the static capacity ($S_{00} \propto -P_{static}/P_c$) produces the restorative acceleration driven by the static pressure gradient:

$$\tilde{\Gamma}^{i}_{00} \approx -\frac{1}{2}\eta^{ij}\partial_j S_{00} = +\frac{1}{\rho_\tau c^2}\nabla^i P_{static} \tag{176}$$

The geodesic equation dictates coordinate acceleration ($a^i \approx -c^2\tilde{\Gamma}^{i}_{00}$). Because the static gradient opposes the dynamic pressure gradient ($\nabla P_{static} = -\nabla P_{dyn}$), this symmetric operator governs the longitudinal transport of the coordinate boundary ($\mathbf{a}_{grav} = -\frac{1}{\rho_\tau}\nabla P_{static} = +\frac{1}{\rho_\tau}\nabla P_{dyn}$). Since the dynamic gradient points inward toward the defect, this replicates the attractive irrotational curvature of General Relativity.

3. Spacetime Torsion:
In Einstein-Cartan theory, spacetime torsion ($T^{\lambda}_{\mu\nu}$) and kinematic vorticity ($\Omega_{\mu\nu}$) are treated as distinct fields. However, if the medium *is* the spacetime grid, a kinematic vortex in the medium is physically identical to a geometric twist in the coordinate frame.

Following the established differential geometry of topological defects in continuous media [28, 26, 24], the spacetime torsion tensor is isomorphic to the physical dislocation density of the medium. In this standard defect-mapping, dislocation density is defined as the curl of the plastic distortion field. By identifying the asymmetric tensor's antisymmetric strain ($A_{\mu\nu}$) as the plastic distortion, the contortion tensor ($K^{\lambda}_{\mu\nu}$) is determined. Spacetime torsion is therefore derived as the covariant exterior derivative of the antisymmetric strain ($T^{\lambda}_{\mu\nu} = \nabla_{[\lambda}A_{\mu\nu]}$).

4. Mapping Clebsch Vorticity to Torsion:
Because the grid is the medium, geometric torsion is sourced by localized rotation. Multiplying the continuum torsion by the macroscopic tension limit (c^4/G) translates the geometric strain gradient into angular momentum density ($s^{\lambda}_{\mu\nu}$):

$$s^{\lambda}_{\mu\nu} = \frac{c^4}{8\pi G}T^{\lambda}_{\mu\nu} = \frac{c^4}{8\pi G}\alpha t_p \nabla^{\lambda}\Omega_{\mu\nu} \tag{177}$$

The spatial curl of the kinematic velocity field ($\Omega_{\mu\nu}$) induces the macroscopic torsion of the coordinate frame, unifying the kinematic defect with the geometric connection.

5. The Macroscopic Torsion Boundary (Decay Limit):
Because Clebsch vorticity ($\Omega_{\mu\nu}$) is confined by the cavitation boundary (r_c), the spacetime torsion undergoes spatial decay outside the defect core ($T^{\lambda}_{\mu\nu} \propto e^{-r/r_c}$). At macroscopic distances ($r \gg r_c$), the antisymmetric contortion components attenuate to zero ($K^{\lambda}_{\mu\nu} \to 0$). The affine connection reduces to the symmetric Levi-Civita component ($\Gamma^{\lambda}_{\mu\nu} \approx \tilde{\Gamma}^{\lambda}_{\mu\nu}$), validating the flat-space Minkowski divergence limit utilized in Section 5.

6. The Riemann Curvature Expansion:
Applying the asymmetric connection to the Riemann curvature tensor expands the derivative operator:

$$R^{\rho}_{\sigma\mu\nu} = \partial_{\mu}\Gamma^{\rho}_{\nu\sigma} - \partial_{\nu}\Gamma^{\rho}_{\mu\sigma} + \Gamma^{\rho}_{\mu\lambda}\Gamma^{\lambda}_{\nu\sigma} - \Gamma^{\rho}_{\nu\lambda}\Gamma^{\lambda}_{\mu\sigma} \tag{178}$$

Substituting the decomposed connection $(\tilde{\Gamma} + K)$ partitions the Riemann tensor into a symmetric curvature $(\tilde{R})$ and a torsion-dependent spin. Normalizing the trace contraction with the macroscopic tension limit $(c^4/8\pi G)$ determines the non-linear Lagrangian density:

$$\mathcal{L} = \frac{c^4}{8\pi G}\left(\tilde{R} + K_{\mu\nu\lambda}K^{\mu\lambda\nu} - K_{\mu\nu}{}^{\nu}K^{\mu\lambda}{}_{\lambda}\right) \tag{179}$$

This expansion recovers the Einstein-Cartan functional limit. The symmetric and antisymmetric traces partition the Lagrangian into the Einstein-Hilbert baseline $(\mathcal{L}_{EH})$ and the Spin-Spin contact interaction $(\mathcal{L}_{spin})$:

$$\mathcal{L} = \underbrace{\frac{c^4}{8\pi G}\tilde{R}}_{\mathcal{L}_{EH}} + \underbrace{\frac{c^4}{8\pi G}\left(K_{\mu\nu\lambda}K^{\mu\lambda\nu} - K_{\mu\nu}{}^{\nu}K^{\mu\lambda}{}_{\lambda}\right)}_{\mathcal{L}_{spin}(A_{\mu\nu})} \tag{180}$$

Macroscopic gravity (symmetric Riemannian curvature) and quantum spin (antisymmetric Clebsch torsion) propagate as coupled, orthogonal deformations.

13. Transverse Wave Mechanics, Chiral Induction, and Plasma Attenuation

The structural capacity of the metric is derived from the spatial Cauchy stress tensor $(\hat{T}^{ij})$. Applying the field equation against the symmetric volumetric strain (S^{ij}) produces the coordinate stress:

$$\hat{T}^{ij} = P_c\Big(S^{ij} - \delta^{ij}\Big) = P_c\left(\frac{1}{\alpha_s^2} - 1\right)\delta^{ij} \tag{181}$$

Taking the spatial divergence (∇_j) extracts the volumetric force density $(f^i = \nabla_j\hat{T}^{ij})$. Because the scalar admittance is bounded by dynamic pressure $(\alpha_s^2 = 1 - P_{dyn}/P_c)$, the continuous restorative force of the metric is governed by the inverse scalar capacity. This establishes a non-linear geometric manifold:

$$f^i = P_c\nabla^i\left(\frac{1}{1 - \frac{P_{dyn}}{P_c}}\right) \tag{182}$$

13.1 Transverse Shear and Strain Propagation (Light)

The continuous metric velocity field partitions into an irrotational compressive mode (dh) and a solenoidal transverse mode $(\mathbf{v}_{\perp})$.

While classical electrodynamics truncates the compressive scalar wave via gauge fixing, the asymmetric tensor preserves it within the symmetric spatial diagonals (S_{ii}). Because the unperturbed vacuum limit $(P_{dyn} \ll P_c)$ dictates that longitudinal dynamic pressure approaches zero, the continuous wave equation is derived from the isolated antisymmetric sector $(A_{\mu\nu})$ of the metric tensor.

The conservation of momentum across the spatial coordinates targets the 4D spacetime divergence of the antisymmetric matrix $(\partial_{\mu}A^{\mu i} = 0)$. Expanding the temporal and spatial components extracts the transverse momentum equivalent:

$$\frac{1}{c^2}\frac{\partial\mathbf{v}_{\perp}}{\partial t} = t_p(\nabla\cdot\mathbf{\Omega}) \tag{183}$$

To define the temporal evolution of the rotational phase, the tensor must satisfy the Bianchi identity ($\partial_{[\mu}A_{\nu\lambda]} = 0$). Expanding the cyclic permutation of the antisymmetric components results in the induction equation, where the temporal derivative of the 2-form equates to the exterior derivative of the transverse 1-form:

$$t_p \frac{\partial \mathbf{\Omega}}{\partial t} = -d\mathbf{v}_\perp \tag{184}$$

Taking the time derivative of the momentum equation and substituting the rotational induction constraint produces the second-order wave equation:

$$\frac{1}{c^2}\frac{\partial^2 \mathbf{v}_\perp}{\partial t^2} = t_p \nabla \cdot \left(\frac{\partial \mathbf{\Omega}}{\partial t}\right) = -\nabla \cdot (d\mathbf{v}_\perp) \tag{185}$$

Because transverse waves are divergence-free ($\nabla \cdot \mathbf{v}_\perp = 0$), the divergence of the exterior derivative ($\nabla \cdot d\mathbf{v}_\perp = \nabla(\nabla \cdot \mathbf{v}_\perp) - \nabla^2 \mathbf{v}_\perp$) simplifies to the Laplacian limit:

$$\frac{\partial^2 \mathbf{v}_\perp}{\partial t^2} = c^2 \nabla^2 \mathbf{v}_\perp \tag{186}$$

The phase velocity resolves to the acoustic limit of the continuum (c). Light manifests as a linearly polarized, coupled transverse-rotational strain propagating through the baseline metric.

13.2 Metric Depletion and Chiral Induction

When transverse strain propagates into a plasma, the ambient scalar admittance depletes ($\alpha_s^2 < 1$). The wave encounters an elevated dynamic pressure and a macroscopic background vorticity ($\mathbf{\Omega}_0$) generated by the collective motion of the embedded topological defects.

The kinematic evolution of the propagating wave is governed by the non-inertial covariance transformations derived in Section 10.3. For a wave propagating through a macroscopic background vorticity, the coordinate frame rotates relative to the transverse strain. By Equation 152, this non-inertial rotation couples the spatial vorticity to the transverse velocity vector.

This transformation dictates the temporal acceleration of the transverse slip coupled to the spatial wave propagation:

$$\frac{1}{c^2}\frac{\partial^2 \mathbf{v}_\perp}{\partial t^2} - \nabla^2 \mathbf{v}_\perp \propto \frac{1}{c^2}\left(\frac{\partial \mathbf{v}_\perp}{\partial t} \mathbf{\Omega}_0\right) \tag{187}$$

Because the inner product contraction of a 1-form against an antisymmetric 2-form yields an orthogonal vector, the temporal acceleration is orthogonal to the temporal derivative.

$$\frac{1}{c^2}\frac{\partial^2 \mathbf{v}_\perp}{\partial t^2} - \nabla^2 \mathbf{v}_\perp \propto \frac{1}{c^2}\left(\mathbf{\Omega}_0 \cdot \frac{\partial \mathbf{v}_\perp}{\partial t}\right) \tag{188}$$

13.3 Macroscopic Measurement Boundaries (Deterministic Eigenstates)

A quantum measurement apparatus constitutes an anisotropic boundary condition ($P_{dyn} \to P_c$), imposing an impedance gradient across the metric.

A defect ($r \to r_c, \alpha_s \to 0$) propagates with an extended transverse strain field ($A_{\mu\nu}$). Intercepting the boundary at an incident angle (θ) relative to the axis of minimum resistance ($\mathbf{n}$), the transverse slip vector ($\mathbf{v}_\perp$) undergoes orthogonal decomposition. The boundary projects the amplitude onto the allowed axis:

$$\mathbf{v}_\perp^{trans} = (\mathbf{v}_\perp \cdot \mathbf{n})\mathbf{n} = \mathbf{v}_\perp \cos\theta \tag{189}$$

The observable measurement corresponds to dynamic pressure (P_{dyn}). Substituting the restricted velocity into Equation 5 calculates the transmitted energy limit:

$$\begin{aligned} P_{dyn}^{trans} &= \frac{1}{2}\rho_\tau |\mathbf{v}_\perp \cos\theta|^2 \\ P_{dyn}^{trans} &= \left(\frac{1}{2}\rho_\tau v_\perp^2\right)\cos^2\theta \\ P_{dyn}^{trans} &= P_{dyn}^{(0)}\cos^2\theta \end{aligned} \tag{190}$$

This tensor contraction recovers Malus's Law. Furthermore, the anisotropic projection of $\mathbf{v}_\perp$ generates a spatial gradient across the defect's extended envelope. Satisfying the Bianchi identity ($\partial_{[\mu}A_{\nu\lambda]} = 0$), the restricted slip vector substitutes into the rotational induction equation (Section 13.1), determining the temporal evolution of the core vorticity 2-form ($\mathbf{\Omega}$):

$$t_p \frac{\partial \mathbf{\Omega}}{\partial t} = -d\left(\mathbf{v}_\perp \cos\theta\right) \tag{191}$$

Applying the exterior derivative product rule for a scalar multiplied by a 1-form expands the boundary condition:

$$\begin{aligned} t_p \frac{\partial \mathbf{\Omega}}{\partial t} &= -\left[\cos\theta(d\mathbf{v}_\perp) + d(\cos\theta)\wedge\mathbf{v}_\perp\right] \\ t_p \frac{\partial \mathbf{\Omega}}{\partial t} &= -\cos\theta(d\mathbf{v}_\perp) + \sin\theta(d\theta\wedge\mathbf{v}_\perp) \end{aligned} \tag{192}$$

The expanded equation dictates the transverse shear stress gradient applied to the core. Because the wave propagates transversely, the incident transverse slip ($\mathbf{v}_\perp$) is strictly orthogonal to the phase gradient ($d\theta$). Therefore, their wedge product evaluates to a maximized, non-zero area element ($d\theta \wedge \mathbf{v}_\perp \neq 0$).

Because the boundary is saturated ($P_{dyn} \to P_c$), the defect must reach a steady-state kinematic equilibrium where the temporal acceleration of the transverse slip vanishes ($d\mathbf{v}_\perp \to 0$). Once this primary term stabilizes, fractional angular states ($\sin\theta \neq 0$) generate an unbounded torque that violates the zero-derivative steady-state boundary ($\partial_t \mathbf{\Omega} = 0$). To satisfy this condition against the non-zero wedge product, the remaining angular strain multiplier evaluates to zero:

$$\sin\theta = 0 \implies \theta \in \{0, \pi\} \tag{193}$$

The topology resolves to spatial equilibrium at aligned or anti-aligned orientations. Bounded by the phase limit (c), this discontinuous bifurcation derives quantum eigenstate convergence (wavefunction collapse) from the orthogonal projection limits of the asymmetric tensor.

13.4 Metric Yielding and Attenuation

The spatial attenuation of the transverse slip vector ($v_{\perp,\mu}$) isolates the mechanism of Debye shielding. From Equation 182, the restorative force of the metric scales with the inverse scalar capacity. When a coordinate grid is saturated by background dynamic pressure ($P_{dyn} \to P_c$), the spatial diagonals of the symmetric tensor diverge ($S_{ii} \to \infty$).

The spatial decay of the transverse strain amplitude (A) relative to the radial propagation axis (r) is governed by the metric's stiffness constraint:

$$\nabla^2 A - \left(\frac{1}{\alpha_s^2 L_0^2}\right) A = 0 \tag{194}$$

This second-order differential equation operates as a continuous screened Poisson boundary. Solving this operator extracts the exponential decay envelope of the strain amplitude:

$$A(r) = A_0 \frac{\exp\left(-\frac{r}{\lambda_D}\right)}{r} \tag{195}$$

Where the effective propagation length (λ_D) equates to the inverse of the spatial coordinate expansion:

$$\lambda_D = L_0 \alpha_s = L_0 \sqrt{1 - \frac{P_{dyn}}{P_c}} \tag{196}$$

As dynamic pressure increases, the attenuation length proportionally shrinks ($\lambda_D < L_0$). The transverse wave is not energetically damped; rather, its spatial advection is confined by the diverging coordinate depth. This maps the relativistic length expansion formula directly to the exponential decay envelope of a plasma screening length, deriving macroscopic Debye shielding from continuous geometry rather than particulate Poisson-Boltzmann statistics.

The yield of the metric determines the macroscopic boundary for magnetic reconnection. The Deborah number dictates the rheological state:

$$De = \frac{1}{\alpha_s^2 \alpha t_p \Omega} \tag{197}$$

As dynamic pressure converges on the critical boundary ($P_{dyn} \to P_c$), the scalar admittance collapses ($\alpha_s^2 \to 0$) extracting the glass transition threshold:

$$\lim_{\alpha_s \to 0} De = \infty \tag{198}$$

The spatial stress ($\hat{T}_{ii}$) diverges to infinity, conserving the thermodynamic budget and decoupling the rotational Clebsch phase ($\mathbf{\Omega}$).

The plasma undergoes reconfiguration, lowering the shear stress (P_{shear}), and recovering scalar capacity. This cancels the infinite stress, deriving magnetic reconnection as a mechanical failure of the matrix rather than a fluid anomaly.

14. Continuous Pressure Dynamics (Fundamental Forces)

Macroscopic force interactions operate as geometric projections of a single continuous pressure gradient.

1. The Isochoric Velocity Scaling

The antisymmetric sector ($A_{\mu\nu}$) dictates the trace-free isochoric constraint for the continuous velocity field ($\nabla \cdot \mathbf{v} = 0$). To determine the scaling of this field across a boundary, the divergence operator expands in spherical coordinates (r, θ, ϕ):

$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r}(r^2 v_r) + \frac{1}{r \sin\theta} \frac{\partial}{\partial \theta}(v_\theta \sin\theta) + \frac{1}{r \sin\theta} \frac{\partial v_\phi}{\partial \phi} = 0 \tag{199}$$

For a spherical boundary, the transverse angular gradients cancel ($\partial_\theta = 0$, $\partial_\phi = 0$):

$$\frac{1}{r^2} \frac{\partial}{\partial r}(r^2 v_r) = 0 \tag{200}$$

Multiplying by the squared radius (r^2) and integrating with respect to r:

$$r^2 v_r = C \implies v_r(r) \equiv v(r) = \frac{C}{r^2} \tag{201}$$

Because continuous volumetric flux is conserved, the scalar velocity field ($v(r)$) scales inversely with the boundary area ($A = 4\pi r^2$).

Substituting this velocity into the Bernoulli capacity equation of the temporal diagonal (S_{00}) constructs the dynamic pressure ($P_{dyn} = \frac{1}{2}\rho_\tau v^2$):

$$P_{dyn}(r) = \frac{1}{2} \rho_\tau \left(\frac{C}{r^2} \right)^2 = \frac{\rho_\tau C^2}{2r^4} \tag{202}$$

2. The Geometric Action Boundary (Deriving the Scalar)

To maintain baseline capacity, the background exerts a restorative confinement pressure according to the dynamic deficit ($P(r) = P_{dyn}(r)$).

The geometric Action (S) of a Clebsch twist constitutes an invariant constant (κ). Action becomes the structural energy (E) integrated over the period (T):

$$S = E \cdot T = \kappa \tag{203}$$

The period represents the duration required for the transverse velocity field to complete one closed 2π rotation at the phase limit (c):

$$T = \frac{2\pi r}{c} \tag{204}$$

By integrating the restorative pressure acting across the boundary area to spatial infinity:

$$\begin{aligned} E &= \int_r^\infty F(r')\, dr' = \int_r^\infty P(r') \cdot 4\pi (r')^2\, dr' \\ E &= \int_r^\infty \left(\frac{\rho_\tau C^2}{2(r')^4} \right) 4\pi (r')^2\, dr' = 2\pi \rho_\tau C^2 \int_r^\infty \frac{1}{(r')^2}\, dr' \\ E &= 2\pi \rho_\tau C^2 \left[-\frac{1}{r'} \right]_r^\infty = \frac{2\pi \rho_\tau C^2}{r} \end{aligned} \tag{205}$$

Substituting the period and energy into the Action limit (κ) extracts the integration constant (C):

$$\begin{aligned} \kappa &= \left(\frac{2\pi \rho_\tau C^2}{r} \right) \left(\frac{2\pi r}{c} \right) \\ \kappa &= \frac{4\pi^2 \rho_\tau C^2}{c} \implies C^2 = \frac{\kappa c}{4\pi^2 \rho_\tau} \end{aligned} \tag{206}$$

The radial dimension cancels, revealing that the integration constant (C) is governed by the invariant action limit (κ) and the baseline density of the vacuum (ρ_τ).

Substituting back into the pressure equation yields the spatial confinement gradient:

$$\begin{aligned} P(r) &= \frac{\rho_\tau}{2r^4} \left(\frac{\kappa c}{4\pi^2 \rho_\tau} \right) \\ P(r) &= \frac{\kappa c}{8\pi^2 r^4} \end{aligned} \tag{207}$$

3. The Kinematic Mass Boundary ($\kappa \equiv h$)

Force operates as pressure projected across a bounding cross-section ($F = P \cdot A$). Substituting the area of the spherical boundary ($A = 4\pi r^2$) into the spatial confinement gradient yields:

$$F(r) = \left(\frac{\kappa c}{8\pi^2 r^4} \right) (4\pi r^2) = \frac{\kappa c}{2\pi r^2} \tag{208}$$

In order to determine the value of the action constant (κ), we calculate the mechanical energy (E) required to integrate this restorative force across the characteristic radius:

$$E(r) = \int_r^\infty F(r')\, dr' = \int_r^\infty \frac{\kappa c}{2\pi (r')^2}\, dr' = \frac{\kappa c}{2\pi r} \tag{209}$$

For a boundary to form a stable rest mass (m), this radial energy constructs the kinematic rest energy of the defect (mc^2). Equating this physical limit to the continuous force work isolates the radius (r):

$$\begin{aligned} mc^2 &= \frac{\kappa c}{2\pi r} \\ r &= \frac{\kappa}{2\pi mc} \end{aligned} \tag{210}$$

Because the geometric bounding radius of a defect is defined by the reduced Compton wavelength ($r = \hbar/mc$), equating the two limits reveals the identity of the scalar action constant:

$$\frac{\kappa}{2\pi mc} = \frac{\hbar}{mc} \implies \kappa = 2\pi\hbar = h \tag{211}$$

The invariant action limit (κ) is the unreduced Planck constant (h). Substituting this definition back into the continuum force equation absorbs the reduction factor:

$$F(r) = \frac{hc}{2\pi r^2} \equiv \frac{\hbar c}{r^2} \tag{212}$$

The force scaling of a localized defect inherently differs from the unscaled volumetric expansion of the manifold (Section 4.2) because the reduction of the $8\pi^2$ temporal integration limit by the 4π spatial surface area generates the boundary constant ($\hbar = h/2\pi$).

4. The Interaction Hierarchy (Force Projections)
The boundary interactions map to distinct coordinate radii.

I. Gravity (The Core Boundary):
Applying the spatial gradient at the geometric minimum of the acoustic cavity ($l_p = \sqrt{\hbar G/c^3}$) reproduces the maximum tension of the continuum:

$$F_{grav} = F(l_p) = \frac{\hbar c}{(\sqrt{\hbar G/c^3})^2} = \frac{c^4}{G} \tag{213}$$

II. The Weak Interaction (The Cavitation Yield):
Calculating the continuous pressure across the Compton boundary of the topological W-Boson ($R_W = \hbar/m_W c$) specifies the volumetric rupture limit:

$$F_{weak} = F(R_W) = \frac{\hbar c}{(\hbar/m_W c)^2} = \frac{m_W^2 c^3}{\hbar} \tag{214}$$

III. The Strong Interaction (The Acoustic Confinement):
Projecting the spatial gradient at the viscoelastic relaxation envelope ($L_{crit} = \alpha\hbar/m_e c$) establishes the static boundary for topological phase equilibrium:

$$F_{strong} = F(L_{crit}) = \frac{\hbar c}{(\alpha\hbar/m_e c)^2} = \frac{m_e^2 c^3}{\alpha^2\hbar} \tag{215}$$

IV. Electromagnetism (The Transverse Leakage):
Applying the spatial gradient at the classical Bohr radius ($a_0 = \hbar/\alpha m_e c$) resolves to the macroscopic electrostatic restoring tension:

$$F_{EM_{base}} = F(a_0) = \frac{\hbar c}{(\hbar/\alpha m_e c)^2} = \frac{\alpha^2 m_e^2 c^3}{\hbar} \tag{216}$$

Because the identical boundary governs the force, the geometric ratio between the macroscopic electromagnetic baseline and the strong boundary is α^4.

5. The Macroscopic Baseline (The Horizon Floor)
Projecting the spatial gradient at the macroscopic scale radius of the universe ($R_{h,0}$) defines the kinematic boundary condition of the cosmic vacuum:

$$F_{cosmic} = F(R_{h,0}) = \frac{\hbar c}{R_{h,0}^2} \equiv \frac{hc}{2\pi R_{h,0}^2} \tag{217}$$

To evaluate the kinematic acceleration floor this tension exerts, the force is divided by the characteristic topological mass of the horizon boundary derived via the unreduced action limit ($m_h = h/R_{h,0}c$):

$$\begin{aligned} a_0 &= \frac{F(R_{h,0})}{m_h} \\ a_0 &= \left(\frac{hc}{2\pi R_{h,0}^2}\right)\left(\frac{R_{h,0}c}{h}\right) = \frac{c^2}{2\pi R_{h,0}} \end{aligned} \tag{218}$$

The action scalar (h) cancels, producing the kinematic acceleration threshold (a_0) dictating galactic rotation curves.

6. Macroscopic Combinatorial Aggregates (Newtonian Gravity)

The unified spatial gradient specifies the continuous state of a singular defect. For a macroscopic object, the physical mass partitions as a combinatorial superposition of boundaries.

The baseline mass required to scale classical aggregate interactions is the Planck mass:

$$M_P = \sqrt{\frac{\hbar c}{G}} \tag{219}$$

The total number of interacting boundaries (N) within a mass boundary (M) equates to the ratio of the aggregate mass to this boundary ($N = M/M_P$). The macroscopic force (F_{macro}) between two aggregate masses (M_1 and M_2) separated by a coordinate radius (r) acts as the combinatorial product of their individual boundaries ($N_1 \cdot N_2$) applied to the unified continuum force:

$$\begin{aligned} F_{macro} &= (N_1 \cdot N_2) \cdot F(r) \\ F_{macro} &= \left(\frac{M_1}{M_P}\right)\left(\frac{M_2}{M_P}\right)\left(\frac{\hbar c}{r^2}\right) \end{aligned} \tag{220}$$

Substituting the squared mass boundary ($M_P^2 = \hbar c/G$) recovers the classical format:

$$\begin{aligned} F_{macro} &= \frac{M_1 M_2}{\left(\frac{\hbar c}{G}\right)}\left(\frac{\hbar c}{r^2}\right) \\ F_{macro} &= M_1 M_2 \left(\frac{G}{\hbar c}\right)\left(\frac{\hbar c}{r^2}\right) = \frac{G M_1 M_2}{r^2} \end{aligned} \tag{221}$$

Universal gravitation scales as the combinatorial accumulation of tension.

15. Macroscopic Lensing (The Optical Pressure Gradient)

The accelerated expansion signal of the universe operates as a longitudinal optical projection of the continuous pressure gradient.

1. Cosmological Redshift as a Metric Capacity Ratio

The nested scalar admittance (α_s) established in Section 2 scales to the macroscopic cosmological domain, providing a geometric derivation for redshift ($1 + z$) independent of metric spatial expansion.

The spatial line element of the symmetric metric dictates how the physical rest wavelength (λ_0) of a photon scales against the background coordinate geometry (dx):

$$\lambda_0^2 = S_{ii} dx^2 = \left(\frac{1}{\alpha_s^2}\right) dx^2 \tag{222}$$

Because the rest wavelength (λ_0) remains invariant, the coordinate wavelength ($\lambda \equiv dx$) scales proportionally to scalar admittance ($dx = \alpha_s \lambda_0$).

When a wave propagates from a saturated emitter coordinate ($\alpha_{emitter}$) into a less perturbed observer coordinate ($\alpha_{observer}$), the wavelength elongates. Redshift determines the geometric ratio of the respective capacities:

$$1 + z = \frac{\lambda_{observed}}{\lambda_{emitted}} = \frac{\alpha_{observer}}{\alpha_{emitter}} \tag{223}$$

Cosmological redshift thus maps to the relative capacity of the vacuum.

2. The Tensor Refractive Index

For a null wave, the coordinate phase velocity (v_c) must satisfy the null geodesic constraint of the asymmetric metric ($ds^2 = S_{00}c^2dt^2 + S_{ii}dr^2 = 0$). Isolating the phase velocity incorporates both the temporal and spatial tensor capacities:

$$v_c(r) = \frac{dr}{dt} = c\sqrt{\frac{-S_{00}(r)}{S_{ii}(r)}} \tag{224}$$

Because the symmetric diagonals map to inverse thermodynamic states ($S_{00} = -\alpha_s^2$ and $S_{ii} = 1/\alpha_s^2$), substituting the Bernoulli limit ($\alpha_s^2 = 1 - P_{dyn}(r)/P_c$) results in the coordinate velocity:

$$v_c(r) = c\sqrt{\frac{\alpha_s^2}{1/\alpha_s^2}} = c\alpha_s^2 = c\left(1 - \frac{P_{dyn}(r)}{P_c}\right) \tag{225}$$

The refractive index ($n(r)$) scales the ratio of the phase limit (c) to the coordinate velocity:

$$n(r) = \frac{c}{v_c(r)} = \left(1 - \frac{P_{dyn}(r)}{P_c}\right)^{-1} \tag{226}$$

Whenever the dynamic pressure (P_{dyn}) is non-zero, the refractive index exceeds unity, resulting in a continuous temporal retardation of the propagating wavefront.

3. The Macroscopic Pressure Substitution

As derived from the continuity constraints of an expanding volumetric manifold, the dynamic pressure approaches P_c toward the causal horizon ($r = ct$). While the boundary constraint governing the continuum limit remains an open derivation, inverting the observable magnitude constraints of the Pantheon+ dataset [7] to this pressure boundary produces the following curve:

$$P_{dyn}(r) = P_c\left(1 - e^{-r/ct}\right)^2 \tag{227}$$

Substituting this back into the refractive index equation yields the optical geometry of the background metric:

$$\begin{aligned} n(r) &= \left(1 - \frac{P_c(1 - e^{-r/ct})^2}{P_c}\right)^{-1} \\ n(r) &= \left(1 - (1 - e^{-r/ct})^2\right)^{-1} \end{aligned} \tag{228}$$

Expanding the binomial term ($1 - (1 - 2e^{-r/ct} + e^{-2r/ct})$) simplifies the operator:

$$\begin{aligned} n(r) &= \left(2e^{-r/ct} - e^{-2r/ct}\right)^{-1} \\ n(r) &= \frac{1}{e^{-r/ct}(2 - e^{-r/ct})} \end{aligned} \tag{229}$$

This function specifies the optical density of the expanding coordinate field.

4. The Longitudinal Path Integral (Shapiro Delay)
According to Fermat's Principle, the apparent optical distance (r_{opt}) a photon travels resolves to the integration of the refractive index along the coordinate path (r_{true}):

$$r_{opt} = \int_0^{r_{true}} n(r')dr' \tag{230}$$

Because the medium maintains $n(r) > 1$, the apparent optical path accumulates continuous delay, guaranteeing $r_{opt} > r_{true}$.

5. The Magnitude Anomaly (Optical Expansion)
When assuming an unperturbed background metric where $n(r) = 1$, the optical path gets equated with the geometric coordinate ($r_{opt} = r_{true}$). Because the underlying metric is an expanding volumetric manifold ($R_h = c \cdot t$), standard candles obey the $(1+z)^{-4}$ Tolman surface brightness scaling required by macroscopic energy conservation [49, 35].

However, the observable distance modulus (μ) calculates the luminosity attenuation across the physical boundary. The differential magnitude ($\Delta\mu$) between the standard linear expectation (μ_{linear}) and the refractive metric observation (μ_{obs}) determines optical divergence:

$$\begin{aligned}
\Delta\mu &= \mu_{obs} - \mu_{linear} \\
\Delta\mu &= [5\log_{10}(r_{opt}(1+z)) + 25] - [5\log_{10}(r_{true}(1+z)) + 25] \\
\Delta\mu &= 5\log_{10}\left(\frac{r_{opt}}{r_{true}}\right)
\end{aligned} \tag{231}$$

The magnitude anomaly ($\Delta\mu > 0$) reproduces the signal historically attributed to accelerated expansion (Dark Energy). Apparent acceleration is therefore not a kinetic property, but an optical projection of the continuous pressure gradient superimposed over the baseline volumetric expansion.

16. Discussion: Predictive Corollaries & Future Work

Applying the foundational limits established in the preceding sections yields the following secondary corollaries.

Scope Limitation: The following represent hypothesized physical equivalencies between the Asymmetric Metric-Tensor and advanced quantum/relativistic phenomena. While the macroscopic boundaries and geometric bases are derived from the tensor in the preceding sections, the complete mappings required to finalize these proofs fall outside the scope of this text. These entries are presented not as closed derivations, but as a theoretical roadmap for future work.

16.1 Symmetric Diagonals (Bulk Strain & Gravitation)

Formalism	Continuum Tensor Mapping
Lemaître-Tolman-Bondi	Models a radially expanding, pressure-free metric [32, 50, 6]. Substituting the macroscopic velocity field ($v_0(r)$) into the tensor diagonals dictates expanding spatial compliance independent of FLRW homogeneity constraints.

Formalism (Continued)	**Continuum Tensor Mapping**
Baryon Acoustic Oscillations	Originates via the spatial derivative of the time-time diagonal ($\hat{U}_{00}$). The physical depth of the continuum expands proportionally to the scalar admittance ($dx = \alpha_s ds$). Volumetric expansion scales as the determinant ($\det(S_{ij}) \propto \alpha_s^{-3}$), mapping to the inverse function of redshift ($V \propto (1+z)^3$), linking boundary acceleration with early cosmological scaling.
The Holographic Bound	Derived from the boundary condition of the symmetric trace ($S_{ii} \to \infty$) [3]. Because the core limit ($r = r_c$) acts as incompressible ($\nabla \cdot \mathbf{u} = 0$), volumetric displacement accumulates across the 2D Dirichlet boundary.
The Casimir Effect	Truncation of the symmetric trace's energy density field ($\hat{T}_{00}$) [10]. Parallel spatial boundaries cause the continuous integration of background pressure ($\int_0^d \hat{T}_{00}\, dz$) to evaluate as a discrete sum, providing the pathway to recover the $1/d^4$ boundary suction.
Wakefield Cavitation	Derived from the spatial gradient of the antisymmetric trace ($\nabla(A_{\mu\nu}A^{\mu\nu})$). An intense transverse shear wave (e.g., a laser pulse) generates a shear stress gradient ($-\nabla P_{shear}$). To conserve the total thermodynamic budget (P_c), this gradient repels the symmetric dynamic pressure (P_{dyn}), structuring the Ponderomotive Force as mass-repulsion yielding metric cavitation.
Alpha Decay	The rupture of a boundary ($P_{dyn} \geq P_c$). Rather than stochastic teleportation, quantum tunneling is governed by the WKB spatial path integral of the temporal strain diagonal ($T \approx \exp(-\int \sqrt{-\hat{U}_{00}}\, dr)$) across the mass gradient. This constructs a phase deficit, providing the basis to derive transmission probability from the compliance of the confining metric well.
Optimal Transport (Wasserstein Metric)	The Benamou-Brenier formulation of the W_2 Wasserstein distance minimizes the integral of kinematic energy ($\int \rho v^2\, dxdt$). Because the continuum tensor solves the integrand as dynamic pressure (P_{dyn}), the geodesic equation is an Optimal Transport plan. Gravitation functions as the minimization of the thermodynamic strain required to displace the metric.
Michelson-Morley Null Result	Volumetric constraint. Because dynamic pressure collapses scalar admittance (α_s^2), the physical coordinate path dynamically compresses by the fraction ($L_\parallel = L_0\alpha_s$). This enforces isotropic acoustic phase invariance ($t_\parallel = t_\perp$), undetectable to internal interferometry. This affirms the perspective of Lorentz invariance, where relativistic contraction originates from physical continuum forces rather than kinematic postulates [5, 8, 51].
Shapiro Time Delay	Setting the spacetime interval to zero ($ds^2 = 0$) for a radial transit path demonstrates that the apparent phase velocity of light drops proportionally to the squared scalar admittance ($dr/dt = c\alpha_s^2$). Integrating this differential transit time over the path recovers the Shapiro delay as continuous optical refraction through the gradient.

Formalism (Continued)	**Continuum Tensor Mapping**
Newton's First Law (Inertia)	Under uniform translation, the dynamic pressure field (P_{dyn}) is symmetric across the longitudinal axis. By the Divergence Theorem, integrating the spatial Cauchy stress gradient over the volume ($\int \nabla P_{dyn}\, dV$) equates to zero. This trace equilibrium recovers d'Alembert's Paradox, conserving momentum.
Newton's Third Law (Action/Reaction)	Bernoulli conservation dictates that an applied kinematic action (∇P_{dyn}) generates an equal and opposite static pressure gradient ($-\nabla P_{static}$). Applying the equilibrium limit ($\partial_t \hat{T}^{0i}_A = -\partial_t \hat{T}^{0i}_B$) transfers this gradient balance into kinematic momentum exchange across the boundary.

16.2 Antisymmetric Off-Diagonals (Transverse Shear & Vorticity)

Formalism	Continuum Tensor Mapping
Gravitoelectromagnetism	The transverse slip ($v_\perp$) maps to the Gravitoelectric field. Metric vorticity ($\mathbf{\Omega}$) functions as the Gravitomagnetic field. This structures frame-dragging as metric vorticity.
Dirac Equation & Clifford Algebra	Because the asymmetric tensor operates as a Clifford geometric product ($S + A \equiv u \cdot v + u \wedge v$), spacetime inherently functions as a multivector. The symmetric tensor ($S_{\mu\nu}$) applies the inner product, supplying a moving frame via the tetrad formalism (e_a^μ) to produce the invariant Minkowski signature (η_{ab}) required to span the Clifford algebra ($C\ell_{1,3}$) and construct the anticommuting Dirac vector matrices ($\{\gamma^a, \gamma^b\} = 2\eta^{ab}$). Correspondingly, the 6 antisymmetric off-diagonals ($A_{\mu\nu}$) apply the wedge product, mapping exactly to the 6 bivector spin generators ($\sigma_{\mu\nu}$). At the quantum boundary, a 360° (2π) spatial rotation returns the symmetric volumetric identity to its original state. However, because the defect remains physically tethered to the background via the continuous Clebsch rotational phase ($d\lambda_\Omega \wedge d\beta_\Omega$), this 2π rotation imparts a residual chiral twist into the antisymmetric off-diagonals. A full 720° (4π) rotation "untangles" the geometric tethers and resets the metric state. This mechanical divergence between the symmetric inner product and the antisymmetric wedge product generates the Spin$(1,3)$ double-cover, producing the 4-component Dirac bi-spinor (ψ).
Mott Scattering & Helicity	The symmetric diagonals generate the $1/r$ scalar GRIN lens, providing the basis for macroscopic $1/Q^4$ scattering cross-sections. Furthermore, transverse waves interacting with the antisymmetric vorticity matrix (A_{ij}) structurally conserve kinematic helicity, establishing the geometric baseline to evaluate the scattering amplitude via the tensor contraction ($\frac{d\sigma}{d\Omega} \propto |\langle \mathbf{v}_f | \hat{U}_{\mu\nu} | \mathbf{v}_i \rangle|^2$), capturing the $\cos^2(\theta/2)$ spin-dependent Mott scattering limit.
The Poynting Vector (Convective Energy Flux)	The convective acceleration expands via the vector identity $(\mathbf{v} \cdot \nabla)\mathbf{v} = \frac{1}{2}\nabla(v^2) - \mathbf{v} \cdot d\mathbf{v}$. By Bernoulli conservation, transverse slip ($\mathbf{v}_\perp$) generates a proportional static pressure deficit ($P_{static} = P_c - \frac{1}{2}\rho_\tau v_\perp^2$) causing the surrounding metric (P_c) to generate a restorative spatial gradient ($-\nabla P_{static}$). Transport is controlled by the tensor contraction of the transverse slip 1-form against the vorticity 2-form ($\mathbf{v}_\perp \cdot \mathbf{\Omega}$), defining the Lamb vector. This breaks the spatial symmetry of the pressure gradient, driving the directional transport of kinematic strain. The inner product of the space-time off-diagonals (electric slip) and space-space off-diagonals (magnetic vorticity) maps directly to the Poynting vector ($S^i \propto A^{0j} A_j{}^i$), deriving electromagnetic energy flux as convective metric momentum.

Formalism (Continued)	**Continuum Tensor Mapping**
Quantum Entanglement & Decoherence	Bell's Theorem relies on combining expectation values from mutually exclusive experimental runs onto a 1D scalar ledger to formulate its inequalities ($S = \|E(a,b) - E(a,b') + \dots\|$), mandating a universal Joint Probability Distribution (JPD). This is a geometric category error. An entangled emission does not create independent point-particles governed by a static scalar variable (λ); it establishes a continuous loop. Polarizers establish the geometric constraints for this unified state. Dynamically altering a detector boundary ($b \to b'$) redefines the integration limits of the governing partial differential equations. The integration probability space (Λ) is therefore contextual to the boundaries ($\Lambda_{ab} \neq \Lambda_{ab'}$), invalidating the assumption of statistical independence. Furthermore, this dynamic alteration rotates the relative coordinate basis of the shared space. By the laws of tensor covariance, scalar projections evaluated in rotated reference frames cannot be added or subtracted without applying the requisite transformation matrices. Because the boundary forces the rotational eigenstates into the complex plane (Section 9.6), the correlated measurement evaluates not as a classical statistical integral, but as the sesquilinear inner product of these complex principal eigenvectors against the boundary normals ($E(a,b) \propto \mathbf{v}_A^\dagger \hat{U}_{\mu\nu} \mathbf{v}_B$). This covariant contraction solves the interference cross-terms, deriving correlated measurement. Finally, because the asymmetric metric accommodates viscosity (Section 9.9), interaction with environmental shear dissipates these shared boundary constraints, providing a physical explanation for quantum decoherence.
CP Violation & Baryogenesis	As established in Section 10.6, Parity inversion maps to Charge conjugation ($P \to C$) for an isolated defect. However, the coupling energy between a charged defect ($\mathbf{v}_\perp$) and a chiral background ($\mathbf{\Omega}_0$) solves as the spatial dot product ($E_{CP} = \mathbf{v}_\perp \cdot \mathbf{\Omega}_0$). Under a combined CP transformation, the defect inverts ($\mathbf{v}_\perp \to -\mathbf{v}_\perp$), but the background vacuum remains un-inverted ($\mathbf{\Omega}_0 \to \mathbf{\Omega}_0$). The coupling energy shifts ($E_{CP} \to -E_{CP}$). Because the energy state is not invariant, perfect reversibility ($CP = I$) is prohibited.

16.3 Cross-Coupled & Non-Linear Dynamics

Formalism	Continuum Tensor Mapping
Almansi Finite Strain	Near the cavitation limit ($r \to r_c$), deformations require non-linear dynamics. Mapping the spatial diagonals to the Eulerian Almansi strain tensor ($e_{ij} = \frac{1}{2}(\delta_{ij} - S_{ij}^{-1})$) models strain-hardening, reproducing Asymptotic Freedom.
SU(3) Symmetry & The Strong Interaction	Isolating the 3×3 spatial sub-matrix of $\hat{U}_{\mu\nu}$ assesses continuous deformation via the Cauchy strain tensor (ε_{ij}). For an incompressible core ($\nabla \cdot \mathbf{v} = 0$), the volumetric rate of change vanishes, isolating the continuous geometry to the deviatoric strain tensor (ε'_{ij}), which is trace-free ($\mathrm{Tr}(\varepsilon') = 0$). This leaves 5 real symmetric deviatoric components and 3 antisymmetric rotational components (A_{ij}). As derived in Section 9.6, at this yield boundary, the transverse rotational eigenstates are forced into the complex plane ($\lambda = 1 \pm i$). The 3 antisymmetric matrices align with the imaginary axis (iA_{ij}). The superposition of these 5 real symmetric matrices and 3 imaginary antisymmetric matrices spans the 8-dimensional space of traceless Hermitian matrices. This provides the isomorphism to the 8 geometric generators of the $\mathfrak{su}(3)$ Lie algebra.
The Lorentz Force	Contracting the antisymmetric off-diagonals against a velocity vector ($F_\mu \propto A_{\mu\nu}v^\nu$) produces the Lorentz force. This translates to the Magnus force exerted on a Clebsch vortex advecting through rotational shear.
Z-Pinch & Sausage Instability	Emerges from trace equilibrium under macroscopic rotational shear, generating azimuthal Clebsch vorticity (Ω_ϕ) elevating P_{shear}. To conserve the thermodynamic limit ($P_{static} = P_c - P_{shear}$), the scalar admittance (α_s^2) collapses. The symmetric spatial diagonals ($1/\alpha_s^2$) contract the radial geometry. If boundary pressure exceeds the restoring capacity, the boundary undergoes topological yield and a physical "sausage" fracture.
Acoustic Covariance	As $v \to c$, dynamic pressure drives longitudinal contraction ($L = L_0/\gamma$). Because the macroscopic boundary remains compressible ($\nu < 0.5$), transverse axes do not expand. The symmetric trace constraint forces the total volume ($V = V_0/\gamma$) to shrink, increasing internal frequency (relativistic mass). The zero-sum tensor trace couples longitudinal translational strain directly to transverse rotational strain. Internal Clebsch circulation accelerates proportionally. Kinematic acceleration bounds when the longitudinal axis compresses to the Planck length (l_p).

Formalism (Continued)	**Continuum Tensor Mapping**
Wake Detachment	Resolves the "Bullet Cluster" anomaly [12]. During high-velocity collisions, entrained wakes (β_ϕ) strip from baryonic anchors. Relativistic time variations between interacting frames generate spatial curvature gradients ($\nabla \hat{U}_{00} \neq 0$). Normalizing the antisymmetric transverse components (A_{ij}) against the Total Scalar Admittance (α_s) causes the macroscopic tension gradient to attenuate the amplitude of unanchored rotational shear. This structural cross-coupling stabilizes the detached GRIN lens.
U(1) Gauge Invariance	Translates gauge symmetry via the Clebsch parametrization ($\mathbf{v} = \nabla h + \mathbf{A}$). An arbitrary shift in the irrotational phase ($h \to h + \chi$) leaves the metric invariant if and only if the transverse field simultaneously transforms ($\mathbf{A} \to \mathbf{A} - \nabla\chi$). This Eulerian identity connects to $U(1)$ phase-rotation symmetry.
Viscoelastic Phase Angle	The ratio of the antisymmetric off-diagonals (transverse elastic shear) to the symmetric spatial diagonals (viscous bulk compliance) outputs the rheological phase angle ($\tan\delta$).
Electromagnetic Decoupling (Running Couplings)	Physical measurement of the gauge field isolates the mixed tensor ($A^\mu{}_\nu = S^{\mu\lambda}A_{\lambda\nu}$). Raising the index applies the inverse symmetric metric ($S^{00} = -1/\alpha_s^2$ and $S^{ii} = \alpha_s^2$), separating the effective coupling strength into orthogonal energy-dependent limits. For time-space interactions (Electric Slip), the mixed tensor scales inversely with scalar admittance ($A^0{}_i = (\alpha/\alpha_s^2)v_\perp/c$). As dynamic pressure approaches the metric capacity ($P_{dyn} \to P_c \implies \alpha_s \to 0$), the effective coupling diverges, structuring QED Vacuum Polarization. For space-space interactions (Magnetic Vorticity), the mixed tensor scales proportionally with scalar admittance ($A^i{}_j = (\alpha\alpha_s^2)t_p\Omega$). As dynamic pressure maximizes, the magnetic coupling bounds to zero. This sets the geometric decoupling of electric and magnetic fields across high-density topological boundaries.
Fourier Curvature Decomposition (QFT Momentum States)	Transforming the d'Alembertian wave operator ($\Box\hat{U}_{\mu\nu} = 0$) from coordinate space into momentum space via multidimensional Fourier transform translates spatial curvature (∇^2) as squared wavenumbers ($-k^2$) and temporal evolution (∂_t^2) as squared frequencies ($-\omega^2$). The asymmetric matrix requires this spectral decomposition to partition orthogonally into a symmetric longitudinal spectrum (gravitational potential) and an antisymmetric transverse spectrum (Clebsch vorticity). Because sharp spatial curvature (high k^2) mandates high kinematic energy states (ω^2), this decomposition provides the structural precursor for Quantum Field Theory, deriving discrete momentum states from continuous metric strain.

References

[1] Aharonov, Y., & Bohm, D. (1959). Significance of electromagnetic potentials in the quantum theory. *Physical Review*, 115(3), 485-491.

[2] Bair-Moshe, B. T. (2026). "The Cosmological Lensing Effect." *Research Square*, Preprint. https://doi.org/10.21203/rs.3.rs-8896448/v1

[3] Bekenstein, J. D. (1973). Black holes and entropy. *Physical Review D*, 7(8), 2333-2346.

[4] Bell, J. S. (1964). On the Einstein Podolsky Rosen paradox. *Physics Physique Fizika*, 1(3), 195-200.

[5] Bell, J. S. (1976). "How to teach special relativity," *Progress in Scientific Culture*, vol. 1, no. 2, pp. 135–144. (Reprinted in *Speakable and Unspeakable in Quantum Mechanics*, Cambridge University Press, 1987).

[6] Bondi, H. (1947). Spherically symmetrical models in general relativity. *Monthly Notices of the Royal Astronomical Society*, 107(5-6), 410-425.

[7] Brout, D., et al. (2022). The Pantheon+ Analysis: Cosmological Constraints. *The Astrophysical Journal*, 938(2), 110.

[8] Brown, H. R. (2005). *Physical Relativity: Space-time structure from a dynamical perspective*. Oxford University Press.

[9] Călugăreanu, G. (1961). Sur les classes d'isotopie des noeuds tridimensionnels et leurs invariants. *Czechoslovak Mathematical Journal*, 11(4), 588-625.

[10] Casimir, H. B. G. (1948). On the attraction between two perfectly conducting plates. *Proceedings of the Royal Netherlands Academy of Arts and Sciences*, 51, 793-795.

[11] Clebsch, A. (1859). Ueber die Integration der hydrodynamischen Gleichungen. *Journal für die reine und angewandte Mathematik*, 56, 1-10.

[12] Clowe, D., Bradač, M., Gonzalez, A. H., Markevitch, M., Randall, S. W., Jones, C., & Zaritsky, D. (2006). A direct empirical proof of the existence of dark matter. *The Astrophysical Journal Letters*, 648(2), L109.

[13] Cosserat, E., & Cosserat, F. (1909). *Théorie des corps déformables*. Hermann et Fils.

[14] Einstein, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Annalen der Physik*, 322(8), 549-560.

[15] Einstein, A. (1907). Über das Relativitätsprinzip und die aus demselben gezogenen Folgerungen. *Jahrbuch der Radioaktivität und Elektronik*, 4, 411-462.

[16] Eringen, A. C. (1999). *Microcontinuum Field Theory I: Foundations and Solids*. Springer.

[17] Feynman, R. P. (1955). Application of quantum mechanics to liquid helium. *Progress in Low Temperature Physics*, 1, 17-53.

[18] Fine, A. (1982). Hidden Variables, Joint Probability, and the Bell Inequalities. *Physical Review Letters*, 48(5), 291-295.

[19] François, J., Ravera, L., & Berghofer, P. (2024). The dressing field method for diffeomorphisms: a relational framework. *Journal of Physics A: Mathematical and Theoretical*, 57(30).

[20] Giacomini, F., Castro-Ruiz, E., & Brukner, Č. (2019). Quantum mechanics and the covariance of physical laws in quantum reference frames. *Nature Communications*, 10(1), 494.

[21] Gordon, W. (1923). Zur Lichtfortpflanzung nach der Relativitätstheorie. *Annalen Physics*, 377(22), 421-456.

[22] Gribov, V. N. (1978). Quantization of Non-Abelian Gauge Theories. *Nuclear Physics B*, 139(1-2), 1-19.

[23] Gurtin, M. E. (1981). *An Introduction to Continuum Mechanics*. Academic Press.

[24] Katanaev, M. O., & Volovich, I. V. (1992). Theory of defects in solids and complex structures on Riemann surfaces. *Annals of Physics*, 216(1), 1-28.

[25] Kibble, T. W. B. (1961). Lorentz invariance and the gravitational field. *Journal of Mathematical Physics*, 2(2), 212-221.

[26] Kleinert, H. (2008). *Multivalued Fields in Condensed Matter, Electromagnetism, and Gravitation*. World Scientific.

[27] Kröner, E. (1958). *Kontinuumstheorie der Versetzungen und Eigenspannungen*. Springer.

[28] Kröner, E. (1981). Continuum theory of defects. In *Physics of defects* (pp. 215-315). North-Holland.

[29] Landau, L. D., & Lifshitz, E. M. (1971). *The Classical Theory of Fields* (Course of Theoretical Physics, Volume 2) (3rd ed.). Pergamon Press.

[30] Landau, L. D., & Lifshitz, E. M. (1986). *Theory of Elasticity* (Course of Theoretical Physics, Volume 7) (3rd ed.). Butterworth-Heinemann.

[31] Landau, L. D., & Lifshitz, E. M. (1987). *Fluid Mechanics* (Course of Theoretical Physics, Volume 6) (2nd ed.). Butterworth-Heinemann.

[32] Lemaître, G. (1933). L'Univers en expansion. *Annales de la Société Scientifique de Bruxelles*, A53, 51-85.

[33] Lense, J., & Thirring, H. (1918). Über den Einfluss der Eigenrotation der Zentralkörper auf die Bewegung der Planeten und Monde nach der Einsteinschen Gravitationstheorie. *Physikalische Zeitschrift*, 19, 156-163.

[34] Lovelock, D. (1971). The Einstein tensor and its generalizations. *Journal of Mathematical Physics*, 12(3), 498-501.

[35] Lubin, L. M., & Sandage, A. (2001). The Tolman Surface Brightness Test for the Reality of the Expansion. IV. A Measurement of the Tolman Signal and the Luminosity Evolution of Early-Type Galaxies. *The Astronomical Journal*, 122(3), 1084-1103.

[36] Lüders, G. (1954). On the Equivalence of Invariance under Time Reversal and under Particle-Antiparticle Conjugation for Relativistic Field Theories. *Kongelige Danske Videnskabernes Selskab, Matematisk-Fysiske Meddelelser*, 28(5).

[37] Milgrom, M. (1983). A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *The Astrophysical Journal*, 270, 365-370.

[38] Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman.

[39] Nye, J. F. (1953). Some geometrical relations in dislocated crystals. *Acta Metallurgica*, 1(2), 153-162.

[40] Onsager, L. (1949). Statistical hydrodynamics. *Il Nuovo Cimento (1943-1954)*, 6(2), 279-287.

[41] Pauli, W. (1955). Exclusion Principle, Lorentz Group and Reflection of Space-Time and Charge. In *Niels Bohr and the Development of Physics* (pp. 30-51). Pergamon Press.

[42] Planck Collaboration (2020). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641, A6.

[43] Podolsky, B. (1942). A Generalized Electrodynamics Part I: Non-Quantum. *Physical Review*, 62(1-2), 68.

[44] Rovelli, C. (1996). Relational Quantum Mechanics. *International Journal of Theoretical Physics*, 35(8), 1637-1678.

[45] de Saint-Venant, A. J. C. B. (1864). Mémoire sur l'équilibre des corps solides, dans les limites de leur élasticité, et sur les conditions de leur résistance, quand les déplacements éprouvés par leurs points ne sont pas très-petits. *Comptes Rendus de l'Académie des Sciences*, 24, 16-21.

[46] Sciama, D. W. (1964). The physical structure of general relativity. *Reviews of Modern Physics*, 36(1), 463.

[47] Singer, I. M. (1978). Some Remarks on the Gribov Ambiguity. *Communications in Mathematical Physics*, 60(1), 7-12.

[48] Stelle, K. S. (1977). Renormalization of higher-derivative gravity. *Physical Review D*, 16(4), 953.

[49] Tolman, R. C. (1930). On the Estimation of Distances in a Curved Universe with a Non-Static Line Element. *Proceedings of the National Academy of Sciences*, 16(7), 511-520.

[50] Tolman, R. C. (1934). Effect of Inhomogeneity on Cosmological Models. *Proceedings of the National Academy of Sciences*, 20(3), 169-176.

[51] Volovik, G. E. (2003). *The Universe in a Helium Droplet*. Clarendon Press.

[52] Volterra, V. (1907). Sur l'équilibre des corps élastiques multiplement connexes. *Annales scientifiques de l'École Normale Supérieure*, 24, 401-517.

[53] Wald, R. M. (1984). *General Relativity*. University of Chicago Press.

[54] Weinberg, S. (1995). *The Quantum Theory of Fields, Volume 1: Foundations*. Cambridge University Press.