



离散数学

Discrete Mathematics

for Computer Science

计算机学院计科系
薛思清 xuesiqing@cug.edu.cn

Discrete Mathematics, 6 Relation



第6讲 关系 Relation (3)

Good order is the foundation of all things.
—Edmund Burke (1729–1797)

Discrete Mathematics, 6 Relation

Outline

几个特殊的二元关系

- 等价关系
- 偏序关系
- 函数

Discrete Mathematics, 6 Relation

In mathematics, an **equivalence relation** is a binary relation that is at the same time a reflexive relation, a symmetric relation and a transitive relation. As a consequence of these properties an equivalence relation provides a partition of a set into equivalence classes.

| Contents [hide] | |
|-----------------|--|
| 1 | Notation |
| 2 | Definition |
| 3 | Examples |
| 3.1 | Simple example |
| 3.2 | Equivalence relations |
| 3.3 | Relations that are not equivalences |
| 4 | Connections to other relations |
| 5 | Well-definedness under an equivalence relation |
| 6 | Equivalence class, quotient set, partition |
| 6.1 | Equivalence class |
| 6.2 | Quotient set |
| 6.3 | Projection |
| 6.4 | Equivalence kernel |
| 6.5 | Partition |
| 6.5.1 | Counting possible partitions |
| 7 | Fundamental theorem of equivalence relations |
| 8 | Comparing equivalence relations |
| 9 | Generating equivalence relations |
| 10 | Algebraic structure |
| 10.1 | Group theory |
| 10.2 | Categories and groupoids |
| 10.3 | Lattices |
| 11 | Equivalence relations and mathematical logic |
| 12 | Euclidean relations |
| 13 | See also |
| 14 | Notes |
| 15 | References |
| 16 | External links |

Discrete Mathematics, 6 Relation

关于等价关系——Why these three properties

In math you may want to ask why these three properties have been selected to represent equivalence between sets? If you are indeed curious, try to think of other properties common to all equivalence relations you may think of. I can only offer a very lame excuse: over many many years mathematicians agreed that the above three are the simplest and the commonest.

Discrete Mathematics, 6 Relation

关于等价关系——Why these three properties


Somehow, by the age of five, we know that



Discrete Mathematics, 6 Relation

关于等价关系——Why these three properties


while



Discrete Mathematics, 6 Relation

关于等价关系——Why these three properties

But then again



Discrete Mathematics, 6 Relation


关于等价关系

Fuzzy equivalence relations

Fuzzy equivalence relations were introduced by Zadeh [37] as a generalization of the concept of an equivalence relation. They have been since widely studied as a way to measure the degree of indistinguishability or similarity between the objects of a given universe of discourse, and they have shown to be useful in different contexts such as fuzzy control, approximate reasoning, fuzzy cluster analysis, etc. Depending on the authors and the context in which they appeared, they have received other names such as similarity relations (original Zadeh's name [37]), indistinguishability operators ([36], [23], [24], [6], [25], [14], [15]), \mathcal{F} -equivalences ([10], [11]), many-valued equivalence relations ([12], [13]), etc.


The first definition of a fuzzy partition was given by Ruspini [33], and it has played a significant role in many studies in fuzzy cluster analysis. Butnariu [1] proposed another definition which was originally based on the Lukasiewicz t-norm, and later defined for an arbitrary t-norm. But, none of these definitions of a fuzzy partition gives a bijective

Discrete Mathematics, 6 Relation



Discrete Mathematics, 6 Relation

关于等价关系



Discrete Mathematics, 6 Relation

软件等价类测试



A set of stamps partitioned in bundles: No stamp is in two bundles, and no bundle is empty.

| 序号 | [A, B, C] | 覆盖等价类 | 输出 |
|----|-----------|--|-------|
| 1 | [1, 4, 5] | (1), (2), (3), (4), (5), (6) | 一般三角形 |
| 2 | [1, 1, 2] | (7) | |
| 3 | [1, 6, 2] | (8) | |
| 4 | [1, 2, 6] | (9) | |
| 5 | [1, 2, 3] | (10) | |
| 6 | [1, 3, 2] | (11) | |
| 7 | [1, 1, 2] | (12) | |
| 8 | [3, 3, 4] | (13), (23), (33), (43), (53), (63), (73) | |
| 9 | [3, 4, 4] | (14), (24), (34), (44), (54), (64), (74) | |
| 10 | [3, 4, 3] | (15), (25), (35), (45), (55), (65), (75) | |
| 11 | [3, 4, 5] | (16), (26), (36), (46), (56), (66), (76) | |
| 12 | [3, 3, 3] | (17), (27), (37), (47), (57), (67), (77) | |
| 13 | [3, 4, 4] | (18), (28), (38), (48), (58), (68), (78) | |
| 14 | [3, 4, 3] | (19), (29), (39), (49), (59), (69), (79) | |
| 15 | [3, 3, 4] | (20), (30), (40), (50), (60), (70), (80) | |

Discrete Mathematics, 6 Relation


关于等价关系

RSA Cryptography



Equivalent code (compilers)

Compiler optimisation of source code by determination and utilization of the equivalence of algebraic expressions in the source code



Modulo equivalence

Discrete Mathematics, 6 Relation

1 等价类

等价关系?

等价类

设 R 是 $A \neq \emptyset$ 上等价关系, $\forall x \in A$, 令 $[x]_R = \{ y \mid y \in A \wedge xRy \}$, 称 $[x]_R$ 为 x 关于 R 的等价类, 简称 x 的等价类, 可简记为 $[x]$.

示例 设 $A = \{a, b, c, d, e\}$, A 上的关系

$\rho = \{(a, a), (a, b), (b, a), (b, b), (c, c), (d, d), (d, e), (e, d), (e, e)\}$,

确定由集合 A 中的元素产生的等价类。

Discrete Mathematics, 6 Relation

1 等价类

示例 设 $A = \{1, 2, 3, 4, 5, 8\}$, 求


$R_3 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \equiv y \pmod{3} \}$ 的等价类, 画出 R_3 的关系图。

解:

$[1] = [4] = \{1, 4\}$,

$[2] = [5] = [8] = \{2, 5, 8\}$,

$[3] = \{3\}$.



Discrete Mathematics, 6 Relation

2 等价关系

等价关系(Equivalence)?

设 $R \subseteq A \times A$; 且 $A \neq \emptyset$, 若 R 是**自反的**, **对称的**, **传递的**

示例: 判断是否等价关系(A 是某班学生):

$R_1 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \text{ 与 } y \text{ 同年生} \}$

$R_2 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \text{ 与 } y \text{ 同姓} \}$

$R_3 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \text{ 的年龄不比 } y \text{ 小} \}$ **\emptyset 是 A 上等价关系吗?**

$R_4 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \text{ 与 } y \text{ 选修同门课程} \}$

$R_5 = \{ \langle x, y \rangle \mid x, y \in A \wedge x \text{ 的体重比 } y \text{ 重} \}$

Discrete Mathematics, 6 Relation

2 等价关系

示例

- 1) 在一群人的集合上年龄相等的关系是等价关系, 而朋友关系不一定是等价关系, 因为它可能不是传递的。
- 2) 命题公式间的逻辑等价关系是等价关系。
- 3) 集合上的恒等关系 I_A 和普遍关系 U_A 都是等价关系。
- 4) 在同一平面上直线之间的平行关系, 三角形之间的相似关系都是等价关系。

Discrete Mathematics, 6 Relation

2 等价关系

练习

假设给定了正整数的序偶集合 A , 在 $A \times A$ 上定义二元关系 R 如下:
 $((x, y), (u, v)) \in R$, 当且仅当 $x * v = y * u$, $*$ 为一般乘法, 请证明 R 是一个等价关系。

Discrete Mathematics, 6 Relation

2 等价关系

示例 设 $n \in \{2, 3, 4, \dots\}$, $x, y \in \mathbb{Z}$, 则 x 与 y 模 n 同余 (be congruent modulo n) $\Leftrightarrow x \equiv y \pmod{n} \Leftrightarrow n \mid (x - y) \Leftrightarrow x - y = kn$ ($k \in \mathbb{Z}$), 称 x, y 具有**同余关系** (Congruence)。

同余等价类


$$[0] = \{ kn \mid k \in \mathbb{Z} \},$$

$$[1] = \{ 1 + kn \mid k \in \mathbb{Z} \},$$

$$[2] = \{ 2 + kn \mid k \in \mathbb{Z} \},$$

...

$$[n-1] = \{ (n-1) + kn \mid k \in \mathbb{Z} \}.$$



Discrete Mathematics, 6 Relation

2 等价关系

注意到, 由等价关系确定的等价类具有如下性质:

$$[x]_R \neq \emptyset;$$


$$xRy \Rightarrow [x]_R = [y]_R;$$

$$\neg xRy \Rightarrow [x]_R \cap [y]_R = \emptyset;$$

$$\bigcup \{ [x]_R \mid x \in A \} = A.$$

$$\begin{aligned} A &= \bigcup \{ [x]_R \mid x \in A \} \\ &\subseteq \bigcup \{ [x]_R \mid x \in A \} \\ &\subseteq \bigcup \{ A \mid x \in A \} \\ &= A. \\ \therefore \bigcup \{ [x]_R \mid x \in A \} &= A. \end{aligned}$$

Discrete Mathematics, 6 Relation



Discrete Mathematics, 6 Relation

3 商集与划分

商集: 设 R 是 $A \neq \emptyset$ 上等价关系,

$$A/R = \{ [x]_R \mid x \in A \}$$

称为 A 关于 R 的商集, 简称 A 的商集.

$$\text{显然 } \bigcup A/R = A.$$

Discrete Mathematics, 6 Relation

3 商集与划分

设 $A \neq \emptyset$, 则

(1) R 是 A 上等价关系 \Rightarrow 商集 A/R 是 A 的划分

(2) Π 是 A 的划分 $\Rightarrow R_\Pi$ 是 A 上等价关系, 其中

$$x R_\Pi y \Leftrightarrow \exists z (z \in \Pi \wedge x \in z \wedge y \in z)$$

称为由划分 Π 所定义的等价关系 (同块关系).

Discrete Mathematics, 6 Relation

An important fact about equivalence relations is expressed by the following

Theorem

Let \sim be an equivalence relation defined between elements of a set A . Then the set A can be written as a union $\bigcup A_i$ of pairwise disjoint subsets $A_i \subset A$ such that

$$a \sim b \text{ iff there exists } A_i \text{ such that } a \in A_i \text{ and } b \in A_i$$

There are several ways to express the fact that $a \sim b$. For example:

- a is equivalent to b .
- a and b are equivalent.
- a and b belong to the same equivalence class.

Discrete Mathematics, 6 Relation

3 商集与划分

示例 $A=\{a,b,c\}$, 求A上全体等价关系

解: A上不同划分共有5种:


$$R_1 = U_A,$$

$$R_2 = I_A \cup \{ \langle b, c \rangle, \langle c, b \rangle \},$$

$$R_3 = I_A \cup \{ \langle a, c \rangle, \langle c, a \rangle \},$$


$$R_4 = I_A \cup \{ \langle a, b \rangle, \langle b, a \rangle \},$$

$$R_5 = I_A.$$




Discrete Mathematics, 6 Relation

3 商集与划分



Set partitions 4; Hasse



The 52 partitions of a set with 5 elements

From Wikimedia

Discrete Mathematics, 6 Relation

Bell triangle

| | | | | | | | | | |
|-----|------|------|------|------|------|------|------|--|--|
| 1 | | | | | | | | | |
| 1 | 2 | | | | | | | | |
| 2 | 3 | 5 | | | | | | | |
| 5 | 7 | 10 | 15 | | | | | | |
| 15 | 20 | 27 | 37 | 52 | | | | | |
| 52 | 67 | 87 | 114 | 151 | 203 | | | | |
| 203 | 255 | 322 | 409 | 523 | 674 | 877 | | | |
| 877 | 1080 | 1335 | 1657 | 2066 | 2589 | 3263 | 4140 | | |
| | | | | | | | | | |

Discrete Mathematics, 6 Relation

Discrete Mathematics, 6 Relation