

NanoGT Match Report: Chronic neurovisceral acid sphingomyelinase deficiency

Disease: Chronic neurovisceral acid sphingomyelinase deficiency (ORPHA:618891)

Primary gene: SMPD1

Gene CDS: 1890 bp

Inheritance: Autosomal recessive

Target tissues scored: liver, spleen, lung, CNS

Interpretation

- At least one high-confidence precedent was found, but this is still a precedent match rather than a clinical-trial recommendation.
- Main review flags: Multi-system disease; define a primary therapeutic target tissue before selecting route/vector; Vector does not naturally cover all annotated disease tissues: spleen; Cell-autonomous protein across multiple tissues; high transduction coverage may be required.

Disease Mechanism Evidence

Molecular mechanism: loss of function

Mechanistic detail: Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS

Gene-addition compatibility: conditional

Preferred modality class: liver or cns gene addition or cross correction

Evidence level/status: direct / needs_user_fact_check

Evidence summary: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery

Evidence source: OMIM SMPD1 607616

Study-Level Limitations

- Catalog-relative ranking: current catalog contains 21 precedent programs and 8 vectors, so absence of a strong match is not proof that no therapy is possible.
 - Modality coverage is limited mainly to AAV and lentiviral precedents; dual-AAV, LNP/mRNA, genome editing, ASO, and transplant-enabling strategies are not fully represented.
 - Endpoint readiness: liver/metabolic targets may have biochemical biomarkers, but biomarker correction must be linked to clinical benefit.
 - Endpoint risk: CNS/neurodevelopmental outcomes may require natural-history data, age-stratified endpoints, and long follow-up because short-term clinical change can be hard to interpret.
 - Endpoint risk: multi-system disease may need a hierarchy of primary and secondary endpoints; one tissue response may not equal whole-disease benefit.
-

Top 5 GT Precedent Matches

Rank	Program	Vector	Score	Confidence	Approval
1	BMN 307	AAV5	8.2/10	High	phase2
2	OAV101-IT	AAV9	7.9/10	High	approved

Rank	Program	Vector	Score	Confidence	Approval
3	Zolgensma	AAV9	7.9/10	High	approved
4	Hemgenix	AAV5	7.8/10	High	approved
5	Libmeldy	LV	7.7/10	High	approved

Match #1: BMN 307

Precedent disease: Phenylketonuria

Vector: AAV5

Tissue target: liver

Composite score: 8.2 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	2.00	2.0	Vector naturally reaches disease target tissue
Protein class	1.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	1.00	2.0	Same or related biological pathway
Modality compatibility	1.50	2.0	Disease mechanism supports gene-addition precedent
Inheritance compatibility	1.00	1.0	AR/XL loss-of-function pattern match
Approval precedent	0.60	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	1.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	0.90	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	1.00	1.0	Established delivery route to target tissue
Organelle targeting	1.00	1.0	Nuclear AAV delivery reaches correct subcellular compartment
TOTAL (normalised)	8.19	10.0	Raw sum / 21×10 (v2: 14 dimensions)

Rationale

- Gene CDS 1353bp / cargo 4700bp (29% utilized)
- Vector tropism plus precedent target match: liver

- Both intracellular proteins
- Inheritance match (Autosomal recessive <-> AR)
- Unknown pathway — neutral score
- Disease mechanism: loss of function — Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS
- Gene-addition modality compatibility: conditionally supports gene addition; review disease-specific constraints
- Mechanism evidence: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery
- Mechanism source: OMIM SMPD1 607616 (<https://omim.org/entry/607616>)
- Approval status: phase2
- Vector immunogenicity (AAV5): low (~9%) — most patients eligible; minimal screening burden
- Moderate therapeutic window — progressive disease with childhood onset; early intervention strongly recommended; newborn screening integration beneficial
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: high privilege — blood-brain barrier severely limits T-cell access; durable expression expected
- Promoter availability: ApoE/hAAT, TBG, transthyretin, albumin — extensively validated; used in Hemgenix, Roctavian, DTX301
- Route of administration: IV systemic — established, minimally invasive; used in all hepatic GT programs
- ORGANELLE TARGETING: COMPATIBLE — SMPD1 standard subcellular localisation; nuclear AAV delivery directly produces functional protein at the correct cellular compartment with no additional organelle-import steps required.

Manual Review Flags

- Multi-system disease; define a primary therapeutic target tissue before selecting route/vector
- Vector does not naturally cover all annotated disease tissues: spleen
- Cell-autonomous protein across multiple tissues; high transduction coverage may be required
- Mechanism evidence is conditionally compatible with gene addition; review the listed disease-specific constraints before treating vector precedent as transferable
- Gene annotation incomplete; packaging, protein class, and localization scores need manual confirmation
- AAV tropism is species- and route-dependent; confirm human target-cell biodistribution rather than relying on animal tropism alone

Match #2: OAV101-IT

Precedent disease: Spinal Muscular Atrophy

Vector: AAV9

Tissue target: CNS/spinal cord

Composite score: 7.9 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity

Dimension	Score	Max	What it measures
Tissue tropism	2.00	2.0	Vector naturally reaches disease target tissue
Protein class	1.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	1.00	2.0	Same or related biological pathway
Modality compatibility	1.50	2.0	Disease mechanism supports gene-addition precedent
Inheritance compatibility	1.00	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	1.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	1.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	0.90	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	1.00	1.0	Established delivery route to target tissue
Organelle targeting	1.00	1.0	Nuclear AAV delivery reaches correct subcellular compartment
TOTAL (normalised)	7.90	10.0	Raw sum / 21×10 (v2: 14 dimensions)

Rationale

- Gene CDS 891bp / cargo 4700bp (19% utilized)
- Vector tropism plus precedent target match: cns
- Both intracellular proteins
- Inheritance match (Autosomal recessive <-> AR)
- Unknown pathway — neutral score
- Disease mechanism: loss of function — Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS
- Gene-addition modality compatibility: conditionally supports gene addition; review disease-specific constraints
- Mechanism evidence: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery
- Mechanism source: OMIM SMPD1 607616 (<https://omim.org/entry/607616>)
- Approval status: approved
- Vector immunogenicity (AAV9): high (~22%) — substantial patient exclusion expected; immunodepletion protocols may be needed
- Moderate therapeutic window — progressive disease with childhood onset; early intervention strongly recommended; newborn screening integration beneficial
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more

targeted dose

- Immune privilege: high privilege — blood-brain barrier severely limits T-cell access; durable expression expected
- Promoter availability: ApoE/hAAT, TBG, transthyretin, albumin — extensively validated; used in Hemgenix, Roctavian, DTX301
- Route of administration: IV systemic — established, minimally invasive; used in all hepatic GT programs
- ORGANELLE TARGETING: COMPATIBLE — SMPD1 standard subcellular localisation; nuclear AAV delivery directly produces functional protein at the correct cellular compartment with no additional organelle-import steps required.

Manual Review Flags

- Multi-system disease; define a primary therapeutic target tissue before selecting route/vector
- Vector does not naturally cover all annotated disease tissues: lung, spleen
- Cell-autonomous protein across multiple tissues; high transduction coverage may be required
- Mechanism evidence is conditionally compatible with gene addition; review the listed disease-specific constraints before treating vector precedent as transferable
- Gene annotation incomplete; packaging, protein class, and localization scores need manual confirmation
- AAV tropism is species- and route-dependent; confirm human target-cell biodistribution rather than relying on animal tropism alone

Match #3: Zolgensma

Precedent disease: Spinal Muscular Atrophy

Vector: AAV9

Tissue target: CNS/motor neuron

Composite score: 7.9 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	2.00	2.0	Vector naturally reaches disease target tissue
Protein class	1.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	1.00	2.0	Same or related biological pathway
Modality compatibility	1.50	2.0	Disease mechanism supports gene-addition precedent
Inheritance compatibility	1.00	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	1.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	1.50	2.0	Can GT be given before irreversible damage?

Dimension	Score	Max	What it measures
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	0.90	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	1.00	1.0	Established delivery route to target tissue
Organelle targeting	1.00	1.0	Nuclear AAV delivery reaches correct subcellular compartment
TOTAL (normalised)	7.90	10.0	Raw sum / 21×10 (v2: 14 dimensions)

Rationale

- Gene CDS 891bp / cargo 4700bp (19% utilized)
- Vector tropism plus precedent target match: cns
- Both intracellular proteins
- Inheritance match (Autosomal recessive <-> AR)
- Unknown pathway — neutral score
- Disease mechanism: loss of function — Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS
- Gene-addition modality compatibility: conditionally supports gene addition; review disease-specific constraints
- Mechanism evidence: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery
- Mechanism source: OMIM SMPD1 607616 (<https://omim.org/entry/607616>)
- Approval status: approved
- Vector immunogenicity (AAV9): high (~22%) — substantial patient exclusion expected; immunodepletion protocols may be needed
- Moderate therapeutic window — progressive disease with childhood onset; early intervention strongly recommended; newborn screening integration beneficial
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: high privilege — blood-brain barrier severely limits T-cell access; durable expression expected
- Promoter availability: ApoE/hAAT, TBG, transthyretin, albumin — extensively validated; used in Hemgenix, Roctavian, DTX301
- Route of administration: IV systemic — established, minimally invasive; used in all hepatic GT programs
- ORGANELLE TARGETING: COMPATIBLE — SMPD1 standard subcellular localisation; nuclear AAV delivery directly produces functional protein at the correct cellular compartment with no additional organelle-import steps required.

Manual Review Flags

- Multi-system disease; define a primary therapeutic target tissue before selecting route/vector
- Vector does not naturally cover all annotated disease tissues: lung, spleen

- Cell-autonomous protein across multiple tissues; high transduction coverage may be required
- Mechanism evidence is conditionally compatible with gene addition; review the listed disease-specific constraints before treating vector precedent as transferable
- Gene annotation incomplete; packaging, protein class, and localization scores need manual confirmation
- AAV tropism is species- and route-dependent; confirm human target-cell biodistribution rather than relying on animal tropism alone

Match #4: Hemgenix

Precedent disease: Hemophilia B

Vector: AAV5

Tissue target: liver

Composite score: 7.8 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	2.00	2.0	Vector naturally reaches disease target tissue
Protein class	0.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	1.00	2.0	Same or related biological pathway
Modality compatibility	1.50	2.0	Disease mechanism supports gene-addition precedent
Inheritance compatibility	0.70	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	1.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	0.90	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	1.00	1.0	Established delivery route to target tissue
Organelle targeting	1.00	1.0	Nuclear AAV delivery reaches correct subcellular compartment
TOTAL (normalised)	7.76	10.0	Raw sum / 21×10 (v2: 14 dimensions)

Rationale

- Gene CDS 1383bp / cargo 4700bp (29% utilized)
- Vector tropism plus precedent target match: liver

- Protein class mismatch
- LOF inheritance — compatible for gene replacement
- Unknown pathway — neutral score
- Disease mechanism: loss of function — Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS
- Gene-addition modality compatibility: conditionally supports gene addition; review disease-specific constraints
- Mechanism evidence: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery
- Mechanism source: OMIM SMPD1 607616 (<https://omim.org/entry/607616>)
- Approval status: approved
- Vector immunogenicity (AAV5): low (~9%) — most patients eligible; minimal screening burden
- Moderate therapeutic window — progressive disease with childhood onset; early intervention strongly recommended; newborn screening integration beneficial
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: high privilege — blood-brain barrier severely limits T-cell access; durable expression expected
- Promoter availability: ApoE/hAAT, TBG, transthyretin, albumin — extensively validated; used in Hemgenix, Roctavian, DTX301
- Route of administration: IV systemic — established, minimally invasive; used in all hepatic GT programs
- ORGANELLE TARGETING: COMPATIBLE — SMPD1 standard subcellular localisation; nuclear AAV delivery directly produces functional protein at the correct cellular compartment with no additional organelle-import steps required.

Manual Review Flags

- Multi-system disease; define a primary therapeutic target tissue before selecting route/vector
- Vector does not naturally cover all annotated disease tissues: spleen
- Cell-autonomous protein across multiple tissues; high transduction coverage may be required
- Mechanism evidence is conditionally compatible with gene addition; review the listed disease-specific constraints before treating vector precedent as transferable
- Gene annotation incomplete; packaging, protein class, and localization scores need manual confirmation
- AAV tropism is species- and route-dependent; confirm human target-cell biodistribution rather than relying on animal tropism alone

Match #5: Libmeldy

Precedent disease: Metachromatic leukodystrophy

Vector: LV

Tissue target: hematopoietic/CNS

Composite score: 7.7 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity

Dimension	Score	Max	What it measures
Tissue tropism	1.50	2.0	Vector naturally reaches disease target tissue
Protein class	0.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	1.00	2.0	Same or related biological pathway
Modality compatibility	1.50	2.0	Disease mechanism supports gene-addition precedent
Inheritance compatibility	1.00	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	1.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	0.90	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	1.00	1.0	Established delivery route to target tissue
Organelle targeting	1.00	1.0	Nuclear AAV delivery reaches correct subcellular compartment
TOTAL (normalised)	7.67	10.0	Raw sum / 21×10 (v2: 14 dimensions)

Rationale

- Gene CDS 1521bp / cargo 8000bp (19% utilized)
- Precedent target match: cns
- Protein class mismatch
- Inheritance match (Autosomal recessive <-> AR)
- Unknown pathway — neutral score
- Disease mechanism: loss of function — Acid sphingomyelinase deficiency causes sphingomyelin accumulation in liver, spleen, lung, and CNS
- Gene-addition modality compatibility: conditionally supports gene addition; review disease-specific constraints
- Mechanism evidence: SMPD1 is a secreted lysosomal enzyme; cross-correction may be feasible for visceral disease but CNS involvement complicates single-compartment delivery
- Mechanism source: OMIM SMPD1 607616 (<https://omim.org/entry/607616>)
- Approval status: approved
- Vector immunogenicity (LV): low (~2%) — most patients eligible; minimal screening burden
- Moderate therapeutic window — progressive disease with childhood onset; early intervention strongly recommended; newborn screening integration beneficial
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose

- Immune privilege: high privilege — blood-brain barrier severely limits T-cell access; durable expression expected
- Promoter availability: ApoE/hAAT, TBG, transthyretin, albumin — extensively validated; used in Hemgenix, Roctavian, DTX301
- Route of administration: IV systemic — established, minimally invasive; used in all hepatic GT programs
- ORGANELLE TARGETING: COMPATIBLE — SMPD1 standard subcellular localisation; nuclear AAV delivery directly produces functional protein at the correct cellular compartment with no additional organelle-import steps required.

Manual Review Flags

- Multi-system disease; define a primary therapeutic target tissue before selecting route/vector
- Vector does not naturally cover all annotated disease tissues: cns, liver, lung, spleen
- Cell-autonomous protein across multiple tissues; high transduction coverage may be required
- Mechanism evidence is conditionally compatible with gene addition; review the listed disease-specific constraints before treating vector precedent as transferable
- Gene annotation incomplete; packaging, protein class, and localization scores need manual confirmation