

NanoGT Match Report: Leber hereditary optic neuropathy

Disease: Leber hereditary optic neuropathy (ORPHA:104)

Primary gene: MT-ND4

Gene CDS: 1377 bp

Inheritance: Mitochondrial inheritance

Target tissues: retina, CNS

Top 5 GT Precedent Matches

Rank	Program	Vector	Score	Confidence	Approval
1	CPCB-RPE1	AAV8	6.9/10	[YELLOW] Medium	phase2/3
2	Skysona	LV	6.8/10	[YELLOW] Medium	approved
3	Luxturna	AAV2	6.6/10	[YELLOW] Medium	approved
4	Libmeldy	LV	6.3/10	[YELLOW] Medium	approved
5	Hemgenix	AAV5	6.0/10	[YELLOW] Medium	approved

Match #1: CPCB-RPE1

Precedent disease: Achromatopsia

Vector: AAV8

Tissue target: retina/photoreceptor

Composite score: 6.9 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	1.50	2.0	Vector naturally reaches disease target tissue
Protein class	1.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	2.00	2.0	Same or related biological pathway
Inheritance compatibility	0.30	1.0	AR/XL loss-of-function pattern match
Approval precedent	0.70	1.0	Regulatory approval / trial stage
Immunogenicity	1.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	0.50	2.0	Can GT be given before irreversible damage?

Dimension	Score	Max	What it measures
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	1.00	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	0.80	1.0	Established delivery route to target tissue
TOTAL (normalised)	6.94	10.0	Raw sum / 18 × 10

Rationale

- Gene CDS 1377bp / cargo 4700bp (29% utilized)
- Precedent target match: retina
- Both membrane proteins
- Inheritance mismatch (dominant/mitochondrial — higher complexity)
- Pathway match: retinal phototransduction
- Approval status: phase2/3
- Vector immunogenicity (AAV8): high (~30%) — substantial patient exclusion expected; immunodepletion protocols may be needed
- Very narrow therapeutic window — congenital or rapidly fatal early onset; in utero or immediate neonatal GT required; substantially increases trial complexity
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: highest immune privilege — blood-retinal barrier + FasL + TGF-β2; minimal T-cell clearance risk
- Promoter availability: VMD2, RPGR, GRK1, CRX, IRBP — multiple validated retinal promoters used in Luxturna, GS010, CPCB-RPE1
- Route of administration: Subretinal/intravitreal injection — specialist ophthalmic procedure; established in Luxturna and GS010

Match #2: Skysona

Precedent disease: Cerebral adrenoleukodystrophy

Vector: LV

Tissue target: hematopoietic/CNS

Composite score: 6.8 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	1.50	2.0	Vector naturally reaches disease target tissue

Dimension	Score	Max	What it measures
Protein class	1.50	2.0	Same se-creted/lysosomal/membrane/intracellular class
Pathway similarity	0.50	2.0	Same or related biological pathway
Inheritance compatibility	0.30	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	0.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	1.00	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	0.80	1.0	Established delivery route to target tissue
TOTAL (normalised)	6.83	10.0	Raw sum / 18 × 10

Rationale

- Gene CDS 1377bp / cargo 8000bp (17% utilized)
- Precedent target match: cns
- Both membrane proteins
- Inheritance mismatch (dominant/mitochondrial — higher complexity)
- Different pathway (retinal_phototransduction vs peroxisomal)
- Approval status: approved
- Vector immunogenicity (LV): low (~2%) — most patients eligible; minimal screening burden
- Very narrow therapeutic window — congenital or rapidly fatal early onset; in utero or immediate neonatal GT required; substantially increases trial complexity
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: highest immune privilege — blood-retinal barrier + FasL + TGF-β2; minimal T-cell clearance risk
- Promoter availability: VMD2, RPGR, GRK1, CRX, IRBP — multiple validated retinal promoters used in Luxturna, GS010, CPCB-RPE1
- Route of administration: Subretinal/intravitreal injection — specialist ophthalmic procedure; established in Luxturna and GS010

Match #3: Luxturna

Precedent disease: Leber congenital amaurosis type 2

Vector: AAV2

Tissue target: retina/RPE

Composite score: 6.6 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	2.00	2.0	Vector naturally reaches disease target tissue
Protein class	0.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	2.00	2.0	Same or related biological pathway
Inheritance compatibility	0.30	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	0.50	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	0.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	1.00	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	0.80	1.0	Established delivery route to target tissue
TOTAL (normalised)	6.56	10.0	Raw sum / 18×10

Rationale

- Gene CDS 1377bp / cargo 4700bp (29% utilized)
- Vector tropism plus precedent target match: retina
- Protein class mismatch
- Inheritance mismatch (dominant/mitochondrial — higher complexity)
- Pathway match: retinal_visual_cycle
- Approval status: approved
- Vector immunogenicity (AAV2): very high (~55%) — majority of patients may be ineligible; major trial design challenge
- Very narrow therapeutic window — congenital or rapidly fatal early onset; in utero or immediate neonatal GT required; substantially increases trial complexity

- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: highest immune privilege — blood-retinal barrier + FasL + TGF- β 2; minimal T-cell clearance risk
- Promoter availability: VMD2, RPGR, GRK1, CRX, IRBP — multiple validated retinal promoters used in Luxturna, GS010, CPCB-RPE1
- Route of administration: Subretinal/intravitreal injection — specialist ophthalmic procedure; established in Luxturna and GS010

Match #4: Libmeldy

Precedent disease: Metachromatic leukodystrophy

Vector: LV

Tissue target: hematopoietic/CNS

Composite score: 6.3 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	1.50	2.0	Vector naturally reaches disease target tissue
Protein class	0.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	0.50	2.0	Same or related biological pathway
Inheritance compatibility	0.30	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	0.50	2.0	Can GT be given before irreversible damage?
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	1.00	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	0.80	1.0	Established delivery route to target tissue
TOTAL (normalised)	6.28	10.0	Raw sum / 18×10

Rationale

- Gene CDS 1377bp / cargo 8000bp (17% utilized)
- Precedent target match: cns
- Protein class mismatch
- Inheritance mismatch (dominant/mitochondrial — higher complexity)
- Different pathway (retinal_phototransduction vs leukodystrophy)
- Approval status: approved
- Vector immunogenicity (LV): low (~2%) — most patients eligible; minimal screening burden
- Very narrow therapeutic window — congenital or rapidly fatal early onset; in utero or immediate neonatal GT required; substantially increases trial complexity
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: highest immune privilege — blood-retinal barrier + FasL + TGF- β 2; minimal T-cell clearance risk
- Promoter availability: VMD2, RPGR, GRK1, CRX, IRBP — multiple validated retinal promoters used in Luxturna, GS010, CPCB-RPE1
- Route of administration: Subretinal/intravitreal injection — specialist ophthalmic procedure; established in Luxturna and GS010

Match #5: Hemgenix

Precedent disease: Hemophilia B

Vector: AAV5

Tissue target: liver

Composite score: 6.0 / 10

Score Breakdown

Dimension	Score	Max	What it measures
Packaging fit	2.00	2.0	Gene CDS size vs vector cargo capacity
Tissue tropism	1.00	2.0	Vector naturally reaches disease target tissue
Protein class	0.50	2.0	Same secreted/lysosomal/membrane/intracellular class
Pathway similarity	0.50	2.0	Same or related biological pathway
Inheritance compatibility	0.30	1.0	AR/XL loss-of-function pattern match
Approval precedent	1.00	1.0	Regulatory approval / trial stage
Immunogenicity	2.00	2.0	Pre-existing NAb seroprevalence for this vector
Therapeutic window	0.50	2.0	Can GT be given before irreversible damage?

Dimension	Score	Max	What it measures
Cross-correction	0.20	1.0	Can transduced cells rescue untransduced neighbours?
Immune privilege	1.00	1.0	Immunological protection of target tissue
Promoter availability	1.00	1.0	Validated tissue-specific promoters exist
Route of administration	0.80	1.0	Established delivery route to target tissue
TOTAL (normalised)	6.00	10.0	Raw sum / 18×10

Rationale

- Gene CDS 1377bp / cargo 4700bp (29% utilized)
- Vector tropism overlaps cns, but precedent target is liver
- Protein class mismatch
- Inheritance mismatch (dominant/mitochondrial — higher complexity)
- Different pathway (retinal phototransduction vs coagulation)
- Approval status: approved
- Vector immunogenicity (AAV5): low (~9%) — most patients eligible; minimal screening burden
- Very narrow therapeutic window — congenital or rapidly fatal early onset; in utero or immediate neonatal GT required; substantially increases trial complexity
- Intracellular or membrane-bound protein — no cross-correction possible; each target cell must individually receive the vector; requires high transduction efficiency and therefore a higher or more targeted dose
- Immune privilege: highest immune privilege — blood-retinal barrier + FasL + TGF- β 2; minimal T-cell clearance risk
- Promoter availability: VMD2, RPGR, GRK1, CRX, IRBP — multiple validated retinal promoters used in Luxturna, GS010, CPCB-RPE1
- Route of administration: Subretinal/intravitreal injection — specialist ophthalmic procedure; established in Luxturna and GS010