

# Therapeutic Exercise is Effective in Reducing the Intensity of Nonspecific Low Back Pain in Children and Adolescents: A Systematic Review and Network Meta-analysis

Author links open overlay panel José Manuel García-Moreno PhD <sup>a</sup>, Inmaculada Calvo-Muñoz PhD <sup>b</sup>, Antonia Gómez-Conesa PhD <sup>c</sup>, José Antonio López-López PhD <sup>c d</sup>

Show more

Add to Mendeley

Share

Cite

<https://doi.org/10.1016/j.apmr.2024.11.002> [Get rights and content](#)

Abstract

Objective

To compare the different physiotherapy treatments and determine the most effective treatment to reduce the nonspecific low back pain (NSLBP) intensity in children and adolescents.

Data Sources

Eight databases (Cochrane Library, MEDLINE, PEDro, Web of Science, LILACS, IBECs, PsycINFO, and SCOPUS), and 2 health-specialized journals (*BMJ* and *Spine*) were searched from inception to May 2023, with no language restriction.

Study Selection

Individuals aged 6-18 years with NSLBP were selected, and physical therapy treatments were considered. Studies were required to be controlled by clinical trials with pretest and posttest evaluations, and to report pain intensity.

Data Extraction

Data extraction and risk of bias assessment were performed independently by 2 reviewers.

## Data Synthesis

A meta-analysis of 11 controlled trials with 827 participants found that physiotherapy treatments effectively reduced NSLBP intensity on posttest measurement ( $d_+ = 0.75$ ; 95% confidence interval [CI], 0.30-1.20) and 6-month follow-up ( $d_+ = 0.35$ ; 95% CI, -0.72 to 1.40). Network meta-analysis showed both therapeutic exercise ( $d_+ = 1.11$ ; 95% CI, 0.48-1.74) and a combination of therapeutic exercise and manual therapy ( $d_+ = 1.45$ ; 95% CI, 0.40-2.49) were effective compared to no treatment. There were no significant differences between therapeutic exercise and the combination of therapeutic exercise and manual therapy.

## Conclusions

Physical exercise has proven to be the most effective treatment for addressing the intensity of NSLBP in children and adolescents. While combining it with manual therapy may yield even better results, it is crucial to emphasize that physical exercise should serve as the cornerstone in the physiotherapeutic approach to managing NSLBP intensity in this age group.

## Introduction

In recent years, an increase in the prevalence and severity of low back pain (LBP) in children and adolescents has been observed,<sup>1</sup> reaching about 39% of lifetime prevalence,<sup>2,3</sup> rising to 73.6% in some studies.<sup>4</sup> Furthermore, LBP in adolescence is associated with LBP in adulthood.<sup>1</sup> The effects of LBP may lead to activity limitations and restricted social participation, with an emphasis on school absenteeism and gym or sports participation,<sup>2</sup> and this is considered a concern that needs to be addressed.<sup>1</sup> Nonspecific LBP (NSLBP) is the most common form of LBP, as the specific cause is rarely identified.<sup>5,6</sup>

Physiotherapists employ a variety of modalities to treat NSLBP in children and adolescents. Among them, therapeutic exercise stands out as a well-studied and highly recommended treatment to control the intensity and disability of NSLBP in all age groups.<sup>7</sup> In addition, education emerges as a valuable tool to reduce the intensity of NSLBP in children and adolescents.<sup>8</sup> The combination of exercise and education is advocated, as the synergistic benefits outweigh those achieved when these interventions are employed individually in all age groups.<sup>9</sup> In addition, manual therapy is another effective intervention employed by physical therapists to alleviate the intensity of NSLBP in all age groups.<sup>10</sup> Some researchers propose the involvement of parents or caregivers in the patient's treatment as a beneficial approach.<sup>11</sup>

There is a large difference between the research activity in the treatment of NSLBP in children and adolescents and in adults.<sup>12</sup> In adults, the various therapeutic approaches for NSLBP have undergone extensive examination, resulting in the formulation of clinical

practice guidelines. In contrast, insufficient information is available for children and adolescents, making it difficult to develop corresponding clinical guidelines for NSLBP in this demographic group.<sup>13</sup> In recent years, a guideline has been published on the treatment of back pain in children and adolescents, but not specific to NSLBP.<sup>14</sup> Therefore, substantial uncertainty remains as to which treatment option should be preferred for NSLBP.

Network meta-analysis (NMA) is a statistical technique that allows the results of more than 2 interventions to be compared.<sup>15,16</sup> This method has been used in systematic reviews on the effectiveness of interventions aimed at reducing NSLBP intensity in adults<sup>17,18</sup> but not in children and adolescents.

Given these considerations, updating the knowledge base is essential, as a considerable amount of time has passed since the last meta-analysis, during which new clinical trials have been published. Furthermore, no NMA has been conducted in this field and population, which suggests that this new analysis could offer insights not captured by traditional meta-analyses.<sup>16</sup>

Therefore, this study aimed to investigate whether physiotherapy treatment could decrease the intensity of NSLBP in children and adolescents and to determine the most effective treatments to achieve this outcome.

Access through your organization

Check access to the full text by signing in through your organization.

Section snippets

Methods

The current meta-analysis was carried out following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement guidelines in the NMA extension<sup>19</sup> and registered with PROSPERO (CRD42021267640).

Results

The search strategy collected 25,696 results (fig 1). After removing duplicates, 20,227 were screened by title and abstract, from which 28 were rated as potentially relevant papers. The articles that were excluded after a review of the body are listed in supplemental appendix S3, with the reasons for exclusion. Finally, 11 studies were included in the meta-analysis.<sup>35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45</sup>

Discussion

A systematic review with pairwise and NMA was conducted to compare different physiotherapy treatments on the intensity of NSLBP in children and adolescents. In the past decade, both systematic reviews about spinal manual therapy in pediatric patients<sup>46,47</sup> and traditional meta-analyses about physical therapy treatments in NSLBP in children and adolescents<sup>48,49</sup> without follow-up analysis had been published. Since then, several new primary studies have been published. This is the first NMA

## Conclusion

Therapeutic exercise has been proven to be effective, and it is recommended that physical therapists prioritize therapeutic exercise as the cornerstone of their treatment approach for reducing pain intensity in cases of NSLBP in children and adolescents. Among therapeutic exercises, lumbar strengthening and stretching exercises are widely recognized as the most common and beneficial.

## Acknowledgments

The authors are grateful to Mitchell Selhorst, PhD and Neta Vitman, MsC, for providing additional information on their studies.

## Ethical approval

Not applicable.

## References (52)

- J Hartvigsen *et al.*

[What low back pain is and why we need to pay attention](#)

Lancet

(2018)

- SJ Kamper *et al.*

[The prevalence, risk factors, prognosis and treatment for back pain in children and adolescents: an overview of systematic reviews](#)

Best Pract Res Clin Rheumatol

(2016)

- JA Hayden *et al.*

[Some types of exercise are more effective than others in people with chronic low back pain: a network meta-analysis](#)

J Physiother

(2021)

- MJ Page *et al.*

[Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement](#)

J Clin Epidemiol

(2021)

- R Brignardello-Petersen *et al.*

[Advances in the GRADE approach to rate the certainty in estimates from a network meta-analysis](#)

J Clin Epidemiol

(2018)

- GL Fanucchi *et al.*

[Exercise reduces the intensity and prevalence of low back pain in 12-13 year old children: a randomised trial](#)

Aust J Physiother

(2009)

- J Hwang *et al.*

Low back pain in children: a rising concern

Eur Spine J

(2019)

- I Calvo-Muñoz *et al.*

Risk factors for low back pain in childhood and adolescence: a systematic review

Clin J Pain

(2018)

- I Calvo-Muñoz *et al.*

Prevalence of low back pain in children and adolescents: a meta-analysis

BMC Pediatr

(2013)

- J Vidal-Conti *et al.*

Prevalence of low back pain among school-aged children between 10 and 12 years

Sustainability

(2021)

- B Minghelli

Low back pain in childhood and adolescent phase: consequences, prevalence and risk factors - a revision

J Spine

(2017)

- HB Vaegter *et al.*

Impaired exercise-induced hypoalgesia in individuals reporting an increase in low back pain during acute exercise

Eur J Pain

(2021)

- P O'Sullivan *et al.*

Understanding adolescent low back pain from a multidimensional perspective: implications for management

J Orthop Sports Phys Ther

(2017)

- KC Jones *et al.*

Pain education with therapeutic exercise in chronic nonspecific low back pain rehabilitation: a critically appraised topic

J Sport Rehabil

(2020)

- JY Hong *et al.*

An updated overview of low back pain management in primary care

Asian Spine J

(2017)

- R Kordi *et al.*

Low back pain in children and adolescents: an algorithmic clinical approach

Iran J Pediatr

(2011)

- SZ George *et al.*

Interventions for the management of acute and chronic low back pain: revision 2021

J Orthop Sports Phys Ther

(2021)

- M Frosch *et al.*

Treatment of unspecific back pain in children and adolescents: results of an evidence-based interdisciplinary guideline

Children (Basel)

(2022)

- DL Belavy *et al.*

Network meta-analysis for comparative effectiveness of treatments for chronic low back pain disorders: systematic review protocol

BMJ Open

(2021)

- CL Arden *et al.*

Implementing the 27 PRISMA 2020 Statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and Sports science) guidance

Br J Sports Med

(2022)

- PJ Owen *et al.*

Which specific modes of exercise training are most effective for treating low back pain?

Network meta-analysis

Br J Sports Med

(2020)

- B Hutton *et al.*

The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations

Ann Intern Med

(2015)

- MW Lipsey

Identifying interesting variables and analysis opportunities

- J Cohen

Statistical power analysis for the behavioral sciences

(1988)

- JAC Sterne *et al.*

RoB 2: a revised tool for assessing risk of bias in randomised trials

BMJ

(2019)

- M Rubio-Aparicio *et al.*

A methodological review of meta-analyses of the effectiveness of clinical psychology treatments

Behav Res Methods

(2018)