





Calculus and Analysis > Differential Equations > Ordinary Differential Equations >

Modified Spherical Bessel Differential Equation

The modified spherical Bessel differential equation is given by the spherical Bessel differential equation with a negative separation constant,

$$r^{2} \frac{d^{2} R}{d r^{2}} + 2 r \frac{d R}{d r} - [r^{2} + n (n+1)] R = 0.$$

The solutions are called modified spherical Bessel functions of the first and second kinds.

SEE ALSO

Modified Spherical Bessel Function of the First Kind, Modified Spherical Bessel Function of the Second Kind, Spherical Bessel Differential Equation

EXPLORE WITH WOLFRAMIALPHA



Solve your math problems and get step-by-step solutions



More things to try:

= adjoint =

= 1.05 * 12,000

= curvilinear asymptote

REFERENCES

Abramowitz, M. and Stegun, I. A. (Eds.). §10.2.1 in *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing.* New York: Dover, pp. 374-377, 1972.

Zwillinger, D. Handbook of Differential Equations, 3rd ed. Boston, MA: Academic Press, p. 121, 1997.

REFERENCED ON WOLFRAM|ALPHA

Modified Spherical Bessel Differential Equation

CITE THIS AS:

Weisstein, Eric W. "Modified Spherical Bessel Differential Equation." From *MathWorld--*A Wolfram Resource. https://mathworld.wolfram.com/ModifiedSphericalBesselDifferentialEquation.html

SUBJECT CLASSIFICATIONS

Calculus and Analysis > Differential Equations > Ordinary Differential Equations >







About MathWorld | MathWorld Classroom | Contribute | MathWorld Book

13,278 Entries | Last Updated: Tue Oct 21 2025 | ©1999-2025 Wolfram Research, Inc. | Terms of Use

wolfram.com | Wolfram for Education | Created, developed and nurtured by Eric Weisstein at Wolfram Research