

## Limits of measurable functions are measurable


**Proposition.** If  $\{f_n\}$  are measurable functions  $(X, \mathcal{M}) \rightarrow \mathbb{R}$ , then

$$\sup_n f_n(x), \inf_n f_n(x), \limsup_{n \rightarrow \infty} f_n(x), \text{ and } \liminf_{n \rightarrow \infty} f_n(x)$$

are all measurable.

$$\limsup_{n \rightarrow \infty} f_n(x) = \inf_{n \geq 1} (\sup_{k \geq n} f_k(x))$$

$$\liminf_{n \rightarrow \infty} f_n(x) = \sup_{n \geq 1} (\inf_{k \geq n} f_k(x))$$



## Limits of measurable functions are measurable

**Proposition.** If  $\{f_n\}$  are measurable functions  $(X, \mathcal{M}) \rightarrow \mathbb{R}$ , then

$$\sup_n f_n(x), \inf_n f_n(x), \limsup_{n \rightarrow \infty} f_n(x), \text{ and } \liminf_{n \rightarrow \infty} f_n(x)$$

are all measurable. And if  $\lim_{n \rightarrow \infty} f_n(x)$  exists, it is measurable.

**Proof of  $\sup_n$ :** Define  $g(x) = \sup_n f_n(x)$ . We want to show  $g$  is measurable, so we want to show that for each  $a \in \mathbb{R}$ ,  $\{x : g > a\} \in \mathcal{M}$ .

Claim:  $\{x : g(x) > a\} = \bigcup_{n=1}^{\infty} \{x : f_n(x) > a\}$ . (Why?)

Since each  $f_n$  is a measurable function, each  $\{x : f_n(x) > a\} \in \mathcal{M}$ .

Therefore  $\{x : g(x) > a\} \in \mathcal{M}$ . Since this works for all  $a$ ,  $\sup_n f_n(x)$  is a measurable function.

Trick question: is the limit of integrable functions integrable?

## Bounded Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space with  $\mu(X) < \infty$ . Let  $\{f_n\}$  be a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $\exists M$  such that  $|f_n(x)| \leq M$  for all  $n$  (and a.e.  $x$ ), then  $f$  is integrable and

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

# Bounded Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space with  $\mu(X) < \infty$ . Let  $\{f_n\}$  be a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $\exists M$  such that  $|f_n(x)| \leq M$  for all  $n$  (and a.e.  $x$ ), then  $f$  is integrable and

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

*Example of why we need boundedness.*

Let  $X = [0, 1]$ ,  $\mu = m$ , and  $f_n = n\chi_{(0, 1/n)}$ .

$$\int \lim_{n \rightarrow \infty} f_n = 0 \neq 1 = \lim_{n \rightarrow \infty} \int f_n$$

## Bounded Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space with  $\mu(X) < \infty$ . Let  $\{f_n\}$  be a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $\exists M$  such that  $|f_n(x)| \leq M$  for all  $n$  (and a.e.  $x$ ), then  $f$  is integrable and

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

*Example of why we need finiteness.*

Let  $X = [0, \infty)$ ,  $\mu = m$ , and  $f_n = \chi_{[n, n+1]}$ . Then  $f_n \rightarrow 0$  pointwise.

$$\int \lim_{n \rightarrow \infty} f_n(x) \, dm = 0 \text{ but } \lim_{n \rightarrow \infty} \int f_n \, dm = 1.$$

# Dominated Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space,  $\{f_n\}$  a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $\exists g \in L^1(X, \mu)$  such that  $|f_n(x)| \leq g$  for all  $n$  and a.e.  $x$ , then  $f \in L^1(X, \mu)$  and

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$



# Monotone Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a  $\sigma$ -finite measure space,  $\{f_n\}$  a sequence of non-negative measurable functions such that  $f_1 \leq f_2 \leq \dots$  a.e. and  $f_n \rightarrow f$  a.e. Then

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

# Monotone Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a  $\sigma$ -finite measure space,  $\{f_n\}$  a sequence of non-negative measurable functions such that  $f_1 \leq f_2 \leq \dots$  a.e. and  $f_n \rightarrow f$  a.e. Then

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

*Why we need non-negativity*

Let  $f_n : [0, \infty) \rightarrow \mathbb{R}$  be defined by  $f_n = (-1/n)\chi_{[n, 2n]}$ . Then  $f_n$  increases pointwise to  $f = 0$ , but

$$\int \lim_{n \rightarrow \infty} f_n = \int f = 0 \not\geq -1 = \lim_{n \rightarrow \infty} \int f_n.$$

# Monotone Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a  $\sigma$ -finite measure space,  $\{f_n\}$  a sequence of non-negative measurable functions such that  $f_1 \leq f_2 \leq \dots$  a.e. and  $f_n \rightarrow f$  a.e. Then

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

*Why we need  $f_n$  increasing to  $f$*

Let  $f_n = n\chi_{(0,1/n)}$ . Then  $f_n$  converges pointwise to  $f = 0$ , but

$$\int \lim_{n \rightarrow \infty} f_n = \int f = 0 \not\leq -1 = \lim_{n \rightarrow \infty} \int f_n.$$

# Monotone Convergence Theorem

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a  $\sigma$ -finite measure space,  $\{f_n\}$  a sequence of non-negative measurable functions such that  $f_1 \leq f_2 \leq \dots$  a.e. and  $f_n \rightarrow f$  a.e. Then

$$\int f = \lim_{n \rightarrow \infty} \int f_n.$$

*Monotone convergence does not hold for Riemann integrals*

Recall our sequence on  $[0, 1]$  whose limit was  $\mathbf{1}_{\mathbb{Q}}$  on  $[0, 1]$ .

## Fatou's Lemma

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space,  $\{f_n\}$  a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $f_n \geq 0$ , then

$$\int f \leq \liminf_{n \rightarrow \infty} \int f_n.$$

# Fatou's Lemma

**Theorem.** Let  $(X, \mathcal{M}, \mu)$  be a measure space,  $\{f_n\}$  a sequence of measurable functions with  $f_n \rightarrow f$  pointwise a.e. If  $f_n \geq 0$ , then

$$\int f \leq \liminf_{n \rightarrow \infty} \int f_n.$$

*Why we need non-negativity*

Let  $f_n : [0, \infty) \rightarrow \mathbb{R}$  be defined by  $f_n = (-1/n)\chi_{[n, 2n]}$ . Then  $f_n$  converges to the zero function, but

$$\int \lim_{n \rightarrow \infty} f_n = \int f = 0 \not\geq -1 = \liminf_{n \rightarrow \infty} \int f_n.$$

## Littlewood's Three Principles

- ★ *Every measurable set is nearly a finite union of intervals.*
- ★ *Every measurable function is nearly continuous.*
- ★ *Every convergent sequence of measurable functions is nearly uniform convergent.*

# Littlewood's Three Principles

- ★ *Every measurable set is nearly a finite union of intervals.*
- ★ *Every measurable function is nearly continuous.*
- ★ *Every convergent sequence of measurable functions is nearly uniform convergent.*

The first principle refers to the definition of outer measure, based on approximating measurable sets with unions of intervals.



# Littlewood's Three Principles

The second principle (*every measurable function is nearly continuous*) is captured by **Lusin's Theorem**:

**Theorem.** Let  $\mu$  be a finite Borel measure on  $X \subseteq \mathbb{R}^n$ , and let  $f : X \rightarrow \mathbb{R}$  be measurable. For all  $\epsilon > 0$ , there exists a closed set  $F \subseteq X$  such that  $f : F \rightarrow \mathbb{R}$  is continuous and  $\mu(X \setminus F) < \epsilon$ .

*Lusin's Theorem might not do what you expect*

Let  $X = [0, 1]$ ,  $f = \chi_{\mathbb{Q} \cap [0, 1]}$ . Let  $E$  be an open cover of  $\mathbb{Q} \cap [0, 1]$  with  $m(E) < \epsilon$ , and let  $F = E^C$ . Then  $m(F^C) < \epsilon$  and  $F$  is closed. Since  $f$  is identically 1 on  $F$ ,  $f : F \rightarrow \mathbb{R}$  is continuous.

But of course  $f : [0, 1] \rightarrow \mathbb{R}$  isn't continuous *anywhere*, including the points of  $F$ . So Lusin's Theorem might trim the domain in a weird way—it doesn't necessarily find a subset on which the original function is continuous.

## Littlewood's Three Principles

The third principle (*every convergent sequence of measurable functions is nearly uniform convergent*) is probably a reference to **Egorov's Theorem**:

**Theorem.** Let  $\mu$  be a finite measure on a metric space  $X$ . Let  $f_n : X \rightarrow \mathbb{R}$  be  $\mu$ -measurable functions that converge to  $f$   $\mu$ -a.e. Then for all  $\delta > 0$  there exists a closed set  $A$  so  $\mu(A^C) < \delta$  and  $f_n \rightarrow f$  uniformly on  $A$ .