

the 1990s, the incidence of *S. flexneri* infections has increased in the United Kingdom [10]. In the United States, the incidence of *S. flexneri* infections has increased in the 1990s, but the incidence of *S. flexneri* infections in the United States is still lower than that in the United Kingdom [11].

There is a need to develop a vaccine against *S. flexneri* infections. The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.

The development of a vaccine against *S. flexneri* infections is a complex task. The development of a vaccine against *S. flexneri* infections requires the identification of the antigens that are most likely to elicit a protective immune response. The development of a vaccine against *S. flexneri* infections also requires the identification of the antigens that are most likely to be conserved among the different strains of *S. flexneri*.