

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [14].

The purpose of this study was to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in the United Kingdom.