

41903 Pset 3

Andrew McKinley, Lauren Mostrom, Pietro Ramella, Francisco Ruela, and Bohan Yang

May 11, 2022

Question 1

Question 2

(a)

```
#setwd("C:/Users/17036/OneDrive/Documents/GitHub/metrics3-zombie-boards/Psets/3")
murder <- read.delim("PS3Data/MURDER_RAW.txt", header=FALSE)

colnames(murder) <- c('id','state','year','mrdrtc','exec','unem','d90','d93',
                     'cmrdrtc','cexec','cunem','cexec1','cunem1')

# Use only data for 1990 and 1993
data <- murder %>%
  filter(year==90 | year==93)

reg_pooled <- lm(mrdrtc~d93+exec+unem, data=data)
### calculate standard errors
# standard homoskedastic standard errors
se.homo_pooled <- sqrt(diag(vcov(reg_pooled)))
# robust standard errors
HCV.coef_pooled <- vcovHC(reg_pooled, type = 'HC1')
se.robust_pooled <- sqrt(diag(HCV.coef_pooled))
# clustered standard errors
CLCV.coef_pooled <- cluster.vcov(reg_pooled,data$state)
se.cluster_pooled <- sqrt(diag(CLCV.coef_pooled))
```

In the table below we report the IID standard errors, heteroskedasticity-robust standard errors, and clustered standard errors. Robust SEs were calculated as follows:

$$\hat{\Omega} = \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{it} \right)^{-1} \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{it} \hat{\epsilon}_{it}^2 \right) \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{it} \right)^{-1}$$

Where $\hat{\epsilon}$ is the POLS residual. We also include clustered standard errors; however, if we suspect the observations are serially correlated (as would be implied by the use of clustered SEs) we should not be using POLS anyway. Nevertheless, clustered SEs were calculated as follows:

$$\hat{\Omega} = \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{it} \right)^{-1} \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{i,90} \hat{\epsilon}_{it} \hat{\epsilon}_{i,90} + x_{it} x'_{i,93} \hat{\epsilon}_{it} \hat{\epsilon}_{i,93} \right) \left(\frac{1}{NT} \sum_{i,t} x_{it} x'_{it} \right)^{-1}$$

Table 1: Pooled OLS			
	IID	Robust	Clustered
(Intercept)	-5.2780 (4.4278)	-5.2780 (5.3868)	-5.2780 (6.6760)
d93	-2.0674 (2.1446)	-2.0674 (1.9981)	-2.0674 (1.3066)
exec	0.1277 (0.2632)	0.1277 (0.1342)	0.1277 (0.1678)
unem	2.5289** (0.7817)	2.5289** (1.1076)	2.5289** (1.5047)
R ²	0.1016	0.1016	0.1016
Adj. R ²	0.0741	0.0741	0.0741
Num. obs.	102	102	102

*** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$

```
texreg(list(reg_pooled, reg_pooled, reg_pooled), digits=4, caption.above=TRUE,
  override.se = list(se.homo_pooled, se.robust_pooled, se.cluster_pooled),
  custom.model.names=c("IID", "Robust", "Clustered"),
  caption = "Pooled OLS")
```

(b)

In this setting FE and FD are numerically identical because there are only two time periods, 1990 and 1993. To see this, recall the fixed effect model:

$$\begin{aligned}
y_{i1} - \bar{y}_i &= (x_{i1} - \bar{x}_i)' \beta + \epsilon_{i1} - \bar{\epsilon}_i \\
y_{i1} - \left(\frac{y_{i1} + y_{i0}}{2} \right) &= \left(x_{i1} - \frac{x_{i1} + x_{i0}}{2} \right)' \beta + \epsilon_{i1} - \left(\frac{\epsilon_{i1} + \epsilon_{i0}}{2} \right) \\
\frac{y_{i1} - y_{i0}}{2} &= \left(\frac{x_{i1} - x_{i0}}{2} \right)' \beta + \frac{\epsilon_{i1} - \epsilon_{i0}}{2} \\
y_{i1} - y_{i0} &= (x_{i1} - x_{i0})' \beta + \epsilon_{i1} - \epsilon_{i0}
\end{aligned}$$

Which is exactly the first differences model.

In the table below we report only the results for FD. We chose FD because it reduces the procedure down to a cross-sectional dataset and only requires heteroskedasticity-robust standard errors, rather than maintaining the panel data structure and requiring clustered standard errors. Clustered standard errors are reported in the table below, but only to illustrate that they are identical to robust in the first differences method when we only have two periods.

There does not appear to be any deterrent effect of capital punishment; based on these results we fail to reject the null hypothesis that the coefficient on *exec* is equal to zero.

(c)

First Differences

```
data_90 <- data %>%
  filter(year==90)
data_93 <- data %>%
  filter(year==93)
```

```

data_fd <- left_join(data_90,data_93,by=c("id"))
data_fd$delta_mrdрте <- data_fd$mrdрте.y-data_fd$mrdрте.x
data_fd$delta_exec <- data_fd$exec.y-data_fd$exec.x
data_fd$delta_unem <- data_fd$unem.y-data_fd$unem.x
reg_fd <- lm(delta_mrdрте~delta_exec+delta_unem, data=data_fd)
### calculate standard errors
# standard homoskedastic standard errors
se.homo_fd <- sqrt(diag(vcov(reg_fd)))
# robust standard errors
HCV.coef_fd <- vcovHC(reg_fd, type = 'HC1')
se.robust_fd <- sqrt(diag(HCV.coef_fd))
# clustered standard errors
CLCV.coef_fd <- cluster.vcov(reg_fd,data_fd$state.x)
se.cluster_fd <- sqrt(diag(CLCV.coef_fd))

```

Table 2: First Differences			
	IID	Robust	Clustered
(Intercept)	0.4133 (0.2094)	0.4133 (0.2000)	0.4133 (0.2000)
delta_exec	-0.1038* (0.0434)	-0.1038* (0.0170)	-0.1038* (0.0170)
delta_unem	-0.0666 (0.1587)	-0.0666 (0.1469)	-0.0666 (0.1469)
R ²	0.1097	0.1097	0.1097
Adj. R ²	0.0727	0.0727	0.0727
Num. obs.	51	51	51

*** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$

```
texreg(list(reg_fd, reg_fd, reg_fd), digits=4, caption.above=TRUE,
         override.se = list(se.homo_fd, se.robust_fd, se.cluster_fd),
         custom.model.names=c("IID", "Robust", "Clustered"),
         caption = "First Differences")
```

As explained above, we used heteroskedasticity-robust SEs because the assumptions required for IID standard errors are too strong, and in a first differences model with only two periods, clustered and robust SEs are identical (as demonstrated in the table above).

From these results the coefficient on *exec* is negative and statistically significant, so it does appear that there is a deterrent effect of capital punishment.

(d)

FD is preferred in this context because POLS requires the assumption that there is no serial correlation between observations. This is an unrealistic assumption in this setting because there could be many persistent features of states that are correlated both with executions and with the murder rate.

Question 3

Question 4

(a)

Assumption: after controlling for *Log of state nonfarm employment*, *state fixed effect* (all the non-observable state-level factors influencing both outcome and the implication of policy, and don't change over time) and *year fixed effect* (all the non-observable year-level macroeconomics factors influencing both outcome and the implication of policy), implication of policy is independent on the error term.

Implication: the implication of contract exceptions increases THS employment by 12.8%, while this is not statistically significant.

```
df <- read.table("PS3Data/autor_out2.txt", header = TRUE,
                 quote="\"", comment.char="")

a <- feols(lnths ~ mico + lnemp | s + t, cluster = 's', df)
out <- etable(a, tex = TRUE)

knitr::asis_output(c("\\begin{center}", out, "\\end{center}"))
```

Dependent Variable:	lnths
Model:	(1)
<i>Variables</i>	
mico	0.1280 (0.0888)
lnemp	2.014*** (0.4236)
<i>Fixed-effects</i>	
s	Yes
t	Yes
<i>Fit statistics</i>	
Observations	850
R ²	0.97270
Within R ²	0.13857
<i>Clustered (s) standard-errors in parentheses</i>	
<i>Signif. Codes: ***: 0.01, **: 0.05, *: 0.1</i>	

(b)

Assumption: after controlling for *Log of state nonfarm employment*, *state fixed effect* (all the non-observable state-level factors influencing both outcome and the implication of policy, and don't change over time), *year fixed effect* (all the non-observable year-level macroeconomics factors influencing both outcome and the implication of policy) and *state-level time trends* (states can have their own linear time trends over the years), implication of policy is independent on the error term.

Implication: the implication of contract exceptions increases THS employment by 14.5%, which is statistically significant.

```
b <- feols(lnths ~ mico + lnemp | s + t + s[t], cluster = 's', df)
out <- etable(b, tex = TRUE)

knitr::asis_output(c("\\begin{center}", out, "\\end{center}"))
```

Dependent Variable:	lnths
Model:	(1)
<i>Variables</i>	
mico	0.1451** (0.0566)
lnemp	1.500*** (0.4139)
<i>Fixed-effects</i>	
s	Yes
t	Yes
<i>Varying Slopes</i>	
t (s)	Yes
<i>Fit statistics</i>	
Observations	850
R ²	0.98848
Within R ²	0.08042
<i>Clustered (s) standard-errors in parentheses</i>	
<i>Signif. Codes: ***: 0.01, **: 0.05, *: 0.1</i>	

(c)

```
c <- feols(lnths ~ 1 | s + t | mico ~ lnemp, cluster = 's', df)
out <- etable(c, tex = TRUE)
```

```
knitr::asis_output(c("\\begin{center}", out, "\\end{center}"))
```

Dependent Variable:	lnths
Model:	(1)
<i>Variables</i>	
mico	-10.18 (23.04)
<i>Fixed-effects</i>	
s	Yes
t	Yes
<i>Fit statistics</i>	
Observations	850
R ²	-1.6737
Within R ²	-83.358
<i>Clustered (s) standard-errors in parentheses</i>	
<i>Signif. Codes: ***: 0.01, **: 0.05, *: 0.1</i>	

(d)

```
df <- panel(df, panel.id = c('s', 't'))

d_a <- feols(lnths ~ mico + lnemp +
             l(mico, 1) + l(mico, 2) + l(mico, 3) + l(mico, 4) +
```



```

      f(mico, 1) + f(mico, 2) + f(mico, 3) + f(mico, 4)
      | s + t, cluster = 's', df)
d_b <- feols(lnths ~ mico + lnemp +
      l(mico, 1) + l(mico, 2) + l(mico, 3) + l(mico, 4) +
      f(mico, 1) + f(mico, 2) + f(mico, 3) + f(mico, 4)
      | s + t + s[t], cluster = 's', df)
out <- etable(d_a, d_b, tex = TRUE)

knitr::asis_output(c("\\begin{center}", out, "\\end{center}"))

```

Dependent Variable:	lnths	
Model:	(1)	(2)
<i>Variables</i>		
mico	0.1579** (0.0622)	0.0754 (0.0668)
lnemp	1.834*** (0.4722)	2.253*** (0.5627)
l(mico,1)	0.0471 (0.0441)	0.0341 (0.0618)
l(mico,2)	0.0461 (0.0357)	0.0454 (0.0510)
l(mico,3)	-0.1076* (0.0551)	-0.1114 (0.0703)
l(mico,4)	0.0439 (0.0660)	0.0497 (0.0602)
f(mico,1)	-0.0662 (0.0455)	-0.0251 (0.0788)
f(mico,2)	0.0528 (0.0525)	-0.0362 (0.0784)
f(mico,3)	0.1167* (0.0691)	0.1282 (0.0867)
f(mico,4)	-0.2371** (0.1084)	-0.0874 (0.1045)
<i>Fixed-effects</i>		
s	Yes	Yes
t	Yes	Yes
<i>Varying Slopes</i>		
t (s)		Yes
<i>Fit statistics</i>		
Observations	450	450
R ²	0.98112	0.99192
Within R ²	0.11847	0.12288
<i>Clustered (s) standard-errors in parentheses</i>		
<i>Signif. Codes: ***: 0.01, **: 0.05, *: 0.1</i>		