

CO142 - Discrete Structures

9th October 2018

Recommended Books

- K.H. Rosen. *Discrete Mathematics and its Applications*
- J.L. Gersting. *Mathematical Structures for Computer Science*
- J.K. Truss. *Discrete Mathematics for Computer Science*
- R. Johnsonbaugh. *Discrete Mathematics*
- C. Schumacher. *Fundamental Notions of Abstract Mathematics*

However, these books don't cover the same content. Learn his notation.

Logical Formula, and Notation

This notation will be shared with **CO140**.

$A \wedge B$	A and B both hold
$A \vee B$	A or B holds (or both)
$\neg A$	A does not hold
$A \Rightarrow B$	if A holds, then so does B
$A \Leftrightarrow B$	A holds if and only if B holds
$\forall x(A)$	the predicate A holds for all x
$\exists x(A)$	the predicate A holds for some x
$a \in A$	the object a is in the set A (a is an element of A)
$a \notin A$	the object a is not in the set A
$=_A$	tests whether two elements of A are the same

Sets

Sets are like data types in Haskell: Haskell data type declaration;

```
data Bool = False | True
```

Set of Boolean values: $\{\text{false}, \text{true}\}$

List of Boolean values: $[\text{true}, \text{false}, \text{true}, \text{false}]$

Set equality: $\{\text{false}, \text{true}\} = \{\text{true}, \text{false}\}$ note order doesn't matter

A set is a collection of objects from a pool of objects. Each object is an *element*, or a *member* of the set. A set *contains* its elements. Sets can be defined in the following ways;

$\{a_1, \dots, a_2\}$	as a collection of n distinct elements
$\{x \in A \mid P(x)\}$	for all the elements in A , where P holds
$\{x \mid P(x)\}$	for all elements, where P holds (dangerous - Russel's paradox)

Use of "triangleq"

The use of \triangleq is for "is defined by". Hence the empty set, $\emptyset \triangleq \{\}$. The difference between \triangleq and $=$, is that the former cannot be proven, it is fact, whereas the latter takes work to prove.

Russel's paradox

Not everything we write as $\{x \mid P(x)\}$ is automatically a set. Assume $R = \{X \mid X \notin X\}$ is a set, the set of all sets which don't contain themselves. As R is a set, then $R \in R$, or $R \notin R$ (law of excluded middle), and thus we can do a case by case analysis.

- Assume $R \in R$. By the definition of R , it then follows that $R \notin R$ (if $R \in R$, then it doesn't satisfy the definition of R) - which is a contradiction.
- Assume $R \notin R$. It then follows that $R \in R$, as it follows the definition of R , hence it is another contradiction.

As both assumptions lead to contradictions, it's possible to write sets which aren't defined. We should only select from a set that we know is defined; $\{x \in A \mid P(x)\}$ - where A is a set.