

CO145 - Mathematical Methods

Prelude

The content discussed here is part of CO145 - Mathematical Methods (Computing MEng); taught by Michael Huth, and Mario Berta, in Imperial College London during the academic year 2018/19. The notes are written for my personal use, and have no guarantee of being correct (although I hope it is, for my own sake). This should be used in conjunction with the lecture notes. This module differs as there isn't as much new content, but it requires practice - as such, I will likely be including worked examples for my own benefit (which are probably incorrect).

Also, the formatting of this document will be a mess. I'd waste too much time trying to figure out how to adjust line spacing otherwise.

Sequences

Formal Definition of a Limit

A sequence a_n , for $n \geq 1$, converges to some limit $l \in \mathbb{R}$ if, and only if, we can prove $\forall \epsilon > 0 [\exists N_\epsilon \in \mathbb{N} [\forall n > N_\epsilon [|a_n - l| < \epsilon]]]$.

To show convergence for the sequence $a_n = \frac{1}{n}$, we need to first make a guess for the limit - suppose $l = 0$. We can now attempt to find some N_ϵ . As $\frac{1}{n} - 0$ is positive for all $n \in \mathbb{N}$, we can drop the absolute, thus it's sufficient to find n such that $\frac{1}{n} < \epsilon$. Since both are positive (hence non-zero), we can take reciprocals on both sides, to get $n > \frac{1}{\epsilon}$. However, we are restricted by the fact that n must be an integer, hence it follows $N_\epsilon = \lceil \frac{1}{\epsilon} \rceil$. For any value of ϵ , we can get some N_ϵ with the function, thus it proves that a limit exists.

Common Converging Sequences

Note that for all of these, we are implicitly saying $\lim_{n \rightarrow \infty}$, and that $a_n \rightarrow 0$.

a_n	condition	N_ϵ
$\frac{1}{n^c}$	for some $c \in \mathbb{R}^+$	$\lceil \frac{1}{\epsilon^c} \rceil$
$\frac{1}{c^n}$	for some $c \in \mathbb{R}$, such that $ c > 1$	$\lceil \log_c(\frac{1}{\epsilon}) \rceil$
c^n	for some $c \in \mathbb{R}$, such that $ c < 1$	$\lceil \log_c(\epsilon) \rceil$
$\frac{1}{n!}$	$n > 1$	$\lceil e^{\frac{1}{\epsilon}} \rceil$
$\frac{1}{\ln(n)}$		

Combining Sequences

Suppose that $a_n \rightarrow a$, and $b_n \rightarrow b$, as $\lim_{n \rightarrow \infty}$;

- $\lim_{n \rightarrow \infty} \lambda a_n = \lambda a$ given $\lambda \in \mathbb{R}$
- $\lim_{n \rightarrow \infty} (a_n + b_n) = a + b$
- $\lim_{n \rightarrow \infty} (a_n b_n) = ab$
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$ given $b \neq 0$

For example, the sequence $a_n = \frac{4n^2+3n}{7n^2+3n-2}$, it's trivial to find the limit as $n \rightarrow \infty$ by inspection as $\frac{4}{7}$. However, if we divide every term by n^2 , we end up with $a_n = \frac{4+\frac{3}{n}}{7+\frac{3}{n}-\frac{2}{n^2}}$, which we can break into $a_n = \frac{b_n}{c_n}$, where $b_n = 4 + \frac{3}{n}$, and $c_n = 7 + \frac{3}{n} - \frac{2}{n^2}$. Using the rules from above, we can further break down the sequences (but I really cannot be bothered to do so), to a point where we get $a = \frac{4+0}{7+0-0} = \frac{4}{7}$.

Sandwich Theorem

In the sandwich theorem, where we want to prove that $\lim_{n \rightarrow \infty} a_n = l$, we need two sequences that form upper, and lower bounds for a_n , namely u_n , and l_n . If such sequences exist, and satisfy $\exists N \in \mathbb{N} [\forall n \geq N [l_n \leq a_n \leq u_n]]$, and both $\lim_{n \rightarrow \infty} u_n = \lim_{n \rightarrow \infty} l_n = l$, then we get $\lim_{n \rightarrow \infty} a_n = l$.

For example, consider the sequence $a_n = \frac{\cos(n)}{n}$. We know that $-1 \leq \cos(n) \leq 1$, therefore $l_n = -\frac{1}{n} \leq a_n \leq \frac{1}{n} = u_n$. However, as both $u_n \rightarrow 0$, and $l_n \rightarrow 0$, when $n \rightarrow \infty$, it follows that $\lim_{n \rightarrow \infty} a_n = 0$.

The sandwich theorem can be proven by finding $N_{\epsilon l}$, and $N_{\epsilon u}$ for l_n , and u_n respectively. As they both converge to the same limit, we can justify that for some $N_\epsilon = \max(N_{\epsilon l}, N_{\epsilon u})$, any $n > N_\epsilon$, we have $|l_n - l| < \epsilon$, and $|u_n - l| < \epsilon$. By removing the modulus signs, we get $-\epsilon < l_n - l < \epsilon$, and $-\epsilon < u_n - l < \epsilon$. By rearranging this, and knowing $l_n < u_n$ by definition, we have $l - \epsilon < l_n < u_n < l + \epsilon$. In order to apply the sandwich theorem, we have to assume $l_n \leq a_n \leq u_n$, hence it follows $l - \epsilon < l_n \leq a_n \leq u_n < l + \epsilon$. This can then be arranged to get $-\epsilon < a_n - l < \epsilon$, thus for all $n > N_\epsilon$, $|a_n - l| < \epsilon$.

Ratio Tests

The follow tests allow us to check if a sequence a_n converges to 0, or diverges. If we want to verify a sequence $\lim_{n \rightarrow \infty} b_n = l$, where $l \neq 0$, we can reformulate it as $a_n = b_n - l$, and prove it for a_n .

If $|\frac{a_{n+1}}{a_n}| \leq c < 1$, for $c \in \mathbb{R}$, where n is sufficiently large, then $\lim_{n \rightarrow \infty} a_n = 0$. Conversely, if $|\frac{a_{n+1}}{a_n}| \geq c > 1$, then a_n diverges.

Suppose $a_n = 2^{-n}$, in order to show it converges, we want to find some c , such that $|\frac{2^{-(n+1)}}{2^{-n}}| \leq c < 1$. By arithmetic, we can say $|\frac{2^{-(n+1)}}{2^{-n}}| = \frac{1}{2} \leq c < 1$. As there exists real $c = \frac{1}{2} < 1$, we can conclude that a_n converges to 0.

However, we cannot prove that $a_n = \frac{1}{n!}$ converges, despite it being a fairly straightforward proof with just the standard ration tests. For example, we would end up with $|\frac{a_{n+1}}{a_n}| = \frac{1}{n+1}$. While this is fairly conclusive that it's less than 1, we don't get a constant c which sits between it, and 1.

Instead of analysing the behaviour of consecutive terms, we take the limit as $r = \lim_{n \rightarrow \infty} |\frac{a_{n+1}}{a_n}|$. If $r < 1$, it converges to 0, but if $r > 1$, it then diverges. Applying this to the previous factorial example, we can clearly see that $\lim_{n \rightarrow \infty} \frac{1}{n+1} = 0$, therefore a_n converges to 0.

Manipulating Absolute Values

A useful decomposition with absolute limits is $|x| < a \Leftrightarrow -a < x < a$. Further useful properties are;

- $|xy| = |x| \cdot |y|$
- $|\frac{x}{y}| = \frac{|x|}{|y|}$
- $|x + y| \leq |x| + |y|$ triangle inequality
- $|x - y| \geq ||x| - |y||$

Properties of Real Numbers

Suppose we have some set S of real numbers, then the following properties hold;

- u is an upper bound on S if $\forall s \in S [u \geq s]$
- l is a lower bound on S if $\forall s \in S [l \leq s]$
- S is bounded above if it has an upper bound, bounded below if it has a lower bound, and bounded if it has both
- a set may have no bounds, or many, therefore we say $\sup(S)$ is the least upper bound of S , and $\inf(S)$ to denote the greatest upper bound of S

- if a set has no upper bound, then $\sup(S) = \infty$, and similarly if there is no lower bound, then $\inf(S) = -\infty$

The fundamental axiom of analysis states that if an increasing sequence of reals is bounded above, then it must converge.

Series

A series is a sum of a sequence - we consider an infinite series as a summation in the form $S = \sum_{n=1}^{\infty} a_n$.

We can also take a partial sum, where we sum from 1 to n instead of ∞ as $S_n = \sum_{i=1}^n a_i$.

Since we're dealing with finite numbers at this point, we can consider S_n as a sequence - thus whether a series converges, or diverges, depends on whether the sequence of partial sums converge, or diverge. If we know that $\forall i \geq 1 [a_i > 0]$, then the partial sum is an increasing sequence where $S_1 < S_2 < S_3 < \dots$, if this is bounded above, then it must converge (from the axiom of analysis).

For a series to have a chance of converging, we require $\lim_{n \rightarrow \infty} a_n = 0$. However, just proving that is insufficient for concluding a series converges. It's important to note that we don't really care about the first values of a series, only how it behaves on large numbers, as it tends to infinity.

If S converges, or diverges, then $\sum_{n=N}^{\infty} a_n$ also converges, or diverges, as it's equivalent to $S - S_{N-1}$.

Geometric Series

To determine whether a limit G exists for the geometric series, we need to determine that the partial sums converge. Hence we do the proof on the right hand side. It's important to note that the only term that changes based on n is x^{n+1} , therefore we can conclude that G must converge when $|x| < 1$ (from earlier results).

Let us represent the geometric series as $G = \sum_{n=1}^{\infty} x^n$. Should a limit G exist - see the left side;

$$\begin{aligned}
 G &= \sum_{n=1}^{\infty} x^n \\
 &= x + \sum_{n=2}^{\infty} x^n \\
 &= x + x \sum_{n=1}^{\infty} x^n \\
 &= x + xG \\
 &= \frac{x}{1-x}
 \end{aligned}
 \qquad
 \begin{aligned}
 G_n &= \sum_{i=1}^n x^i \\
 &= x + \sum_{i=2}^n x^i \\
 &= x + x \sum_{i=1}^{n-1} x^i \\
 &= x + x(G_n - x^n) \\
 &= \frac{x - x^{n+1}}{1-x}
 \end{aligned}$$

Harmonic Series

The harmonic series is often used in comparison tests to prove that another series diverges.

Let harmonic series be written as $S = \sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \underbrace{\left(\frac{1}{3} + \frac{1}{4}\right)}_{> \frac{1}{4} + \frac{1}{4}} + \underbrace{\left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right)}_{> \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}} + \dots > 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \dots$

Therefore, with partial sums, we get $S_n = \sum_{i=1}^{2^n} \frac{1}{i} > 1 + \frac{n}{2}$. As S_n clearly diverges, S must also diverge.

Inverse Squares

Suppose we have the sum of squares $S = \sum_{n=1}^{\infty} \frac{1}{n^2}$, and an auxiliary partial sum $T_n = \sum_{i=1}^n \frac{1}{i(i+1)}$.

To prove that S converges, we should consider $T_n = 1 - \frac{1}{n+1}$ (write out the first few terms and a pattern should be spotted). Hence $\lim_{n \rightarrow \infty} T_n = 1$. Using this, we can argue how the partial sums of the series of inverse squares are bounded. It's trivial to say the following inequality; $\frac{1}{i(i+1)} < \frac{1}{i^2} < \frac{1}{i(i-1)}$.

If we sum this inequality from $i = 2$ to n , we can then justify the following - it's important to note that the sums on the left, and right, differ by only the first, and last term (therefore we can reduce it to a partial sum);

$$\begin{aligned} \frac{1}{2(3)} + \frac{1}{3(4)} + \frac{1}{4(5)} + \dots + \frac{1}{n(n+1)} &< \sum_{i=2}^n \frac{1}{i^2} < \frac{1}{1(2)} + \frac{1}{2(3)} + \frac{1}{3(4)} + \dots + \frac{1}{n(n-1)} \\ \Leftrightarrow T_n - \frac{1}{2} &< \sum_{i=2}^n \frac{1}{i^2} < T_{n-1} \\ \Leftrightarrow T_n + \frac{1}{2} &< \sum_{i=1}^n \frac{1}{i^2} < T_{n-1} + 1 \\ \Leftrightarrow \frac{3}{2} + \frac{1}{n+1} &< S_n < 2 + \frac{1}{n} \end{aligned}$$

Therefore, it follows that the partial sums are bounded by $\frac{3}{2}$, and 2. The existence of an upper bound suggests the sequence converges, therefore the series must also converge.

Important Series

These examples will be useful for the comparison tests covered later on. We can use these results to aid us in proving whether some other series converges, or diverges.

name	S	condition	diverges or converges
harmonic series	$\sum_{n=1}^{\infty} \frac{1}{n}$		diverges
harmonic primes	$\sum_{p:\text{prime}} \frac{1}{p}$		diverges
geometric series	$\sum_{n=1}^{\infty} x^n$	$ x \geq 1$	diverges
geometric series	$\sum_{n=1}^{\infty} x^n$	$ x < 1$	converges
inverse squares series	$\sum_{n=1}^{\infty} \frac{1}{n^2}$		converges
???	$\sum_{n=1}^{\infty} \frac{1}{n^c}$	$c > 1$	converges

Convergence Tests

For brevity, suppose we have the following;

- $S = \sum_{i=1}^{\infty} a_i$ a series we want to reason about, and $a_i \geq 0$
- $\sum_{i=1}^{\infty} c_i$ a series we have already established converges to c
- $\sum_{i=1}^{\infty} d_i$ a series we have already established diverges

In the **comparison test**, we have some $\lambda > 0$, and a $N \in \mathbb{N}$. If $\forall i > N [a_i \leq \lambda c_i]$, then S converges. On the other hand, if $\forall i > N [a_i \geq \lambda d_i]$, then S diverges.

To avoid having to find such a λ , we can use the **limit comparison test**, in which we can say if $\lim_{i \rightarrow \infty} \frac{a_i}{c_i}$ exists, then S converges, but if $\lim_{i \rightarrow \infty} \frac{d_i}{a_i}$ exists, then S diverges.

Another useful test for finding convergence, without needing a pre-established series, is **D'Alembert's ratio test**. Once again, we consider this from some point $N \in \mathbb{N}$, where the cases are;

1. $\forall i \geq N[\frac{a_{i+1}}{a_i} \geq 1]$, then S diverges
2. there exists a k such that $\forall i \geq N[\frac{a_{i+1}}{a_i} \leq k < 1]$, then S converges

This is similar to the comparison test for sequences, and once again the requirement of finding some k between the ratio, and 1 can be avoided by taking limits. This leads to **D'Alembert's limit ratio test**, in which we have the following cases;

1. $\lim_{i \rightarrow \infty} \frac{a_{i+1}}{a_i} > 1$, then S diverges
2. $\lim_{i \rightarrow \infty} \frac{a_{i+1}}{a_i} = 1$, then it is inconclusive
3. $\lim_{i \rightarrow \infty} \frac{a_{i+1}}{a_i} < 1$, then S converges

The second case, in which it is inconclusive, can be illustrated by setting $a_n = 1$, which obviously causes S to diverge, or by setting $a_n = \frac{1}{n^2}$ which causes S to converge. However, both lead to a limit test of 1.

The **integral test** for a sequence, where $a_n = f(n)$ is a decreasing function relies on the following idea;

$$\sum_{n=1}^{\infty} a_{n+1} < \int_1^{\infty} f(x) dx < \sum_{n=1}^{\infty} a_n$$

In practice, if the integral were to diverge once evaluated, the series would also diverge by the right hand inequality, but if it were to converge, then the series would converge with the left hand inequality.

Absolute Convergence

Consider a series with negative a_n values, and another series $S' = \sum_{n=1}^{\infty} |a_n|$. If S' converges, so does S .

However, the same isn't true the other way around, for example, when we have $a_n = (-1)^{n-1} \frac{1}{n}$, S converges to $\ln(2)$, however, doing this with the absolute series, S' , it would diverge, as if it were the normal harmonic series. In order to test for **absolute** convergence - we have the following cases;

1. if $\lim_{n \rightarrow \infty} |\frac{a_{n+1}}{a_n}| > 1$ then $\sum_{n=1}^{\infty} a_n$ diverges
2. if $\lim_{n \rightarrow \infty} |\frac{a_{n+1}}{a_n}| = 1$ then $\sum_{n=1}^{\infty} a_n$ can converge, or diverge
3. if $\lim_{n \rightarrow \infty} |\frac{a_{n+1}}{a_n}| < 1$ then $\sum_{n=1}^{\infty} a_n$ converges absolutely, and therefore also converges

Power Series, and Radius of Convergence

We can represent a function as a power series; $f(x) = \sum_{n=0}^{\infty} a_n x^n$.

Consider the case where $f(x) = \frac{x}{1-x}$, which has the expansion $\sum_{n=1}^{\infty} x^n$, which only converges for $|x| < 1$.

Take the power series $S = \sum_{n=1}^{\infty} n^2 x^n$, and apply the **D'Alembert ratio test**, we'd get the following;

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(n+1)^2 x^{n+1}}{n^2 x^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| x \left(1 + \frac{1}{n} \right)^2 \right| \\ &= |x| \end{aligned}$$

As we want to establish convergence, we set the result $|x| < 1$, therefore it follows that the ratio of convergence for this particular power series is 1.

Power Series

As previously mentioned, a function can be represented in a series expansion on x ; $f(x) = \sum_{n=0}^{\infty} a_n x^n$.

Maclaurin Series

Let some function $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$, then we can easily work out a_0 , by setting $x = 0$, such that $f(0) = a_0$. By differentiating the function, we get $f'(x) = a_1 + 2a_2x + 3a_3x^2 + \dots$, therefore it follows that $f'(0) = a_1$. By differentiating again, we get $f''(x) = 2a_2 + 2 \cdot 3a_3x + \dots$, therefore it follows that $\frac{f''(0)}{2!} = a_2$. Hence, generally $a_n = \frac{f^{(n)}(0)}{n!}$, where $n \geq 0$.

A **Maclaurin series** is the representation of $f(x) = \sum_{i=0}^{\infty} a_i x^i$.

This function can then be differentiated n times to get $f^{(n)}(x) = \sum_{i=n}^{\infty} a_i (i-1)\dots(i-n+1)x^{i-n}$.

From this, we can see that $f^{(n)}(0) = n!a_n$, hence $f(x) = \sum_{n=0}^{\infty} f^{(n)}(0) \frac{x^n}{n!}$.

For example, if we take $f(x) = \ln(1+x)$, and differentiate repeatedly;

$$\begin{array}{ll} f(x) &= \ln(1+x) & f(0) &= 0 \\ f'(x) &= (1+x)^{-1} & f'(0) &= 1 \\ f''(x) &= (-1)(1+x)^{-2} & f''(0) &= -1 \\ f'''(x) &= (-2)(-1)(1+x)^{-3} & f'''(0) &= 2 \\ f^{(n)}(x) &= \frac{(-1)^{n-1}(n-1)!}{(1+x)^n} & f^{(n)}(0) &= (-1)^{n-1}(n-1)! \\ a_n &= \frac{(-1)^{n-1}}{n} \end{array}$$

Therefore, we can justify that $f(x) = \ln(1+x) = \frac{x}{1} - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$. By setting $x = -1$, we'd get $-\infty$, and the power series would become the harmonic series, which we know to diverge. However, by setting $x = 1$, we get $\ln(2)$, is the convergence of the alternating harmonic series (as previously mentioned, in the ratio test example).

Taylor Series

When we're interested in values around a certain point, we can use a more generalised version of a **Maclaurin series**. The general result, where we apply the series around a is;

$$f(x) = f(a) + \frac{f^{(1)}(a)}{1!}(x-a)^1 + \frac{f^{(2)}(a)}{2!}(x-a)^2 + \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$$

For example, if we take $f(x) = \ln(x)$ around a , and differentiate repeatedly;

$$\begin{array}{ll} f(x) &= \ln(x) & f(2) &= \ln(2) \\ f'(x) &= \frac{1}{x} & f'(2) &= \frac{1}{2} \\ f''(x) &= (-1)\frac{1}{x^2} & f''(2) &= -\frac{1}{4} \\ f'''(x) &= (-2)(-1)\frac{1}{x^3} & f'''(2) &= \frac{1}{4} \\ f^{(n)}(x) &= \frac{(-1)^{n-1}(n-1)!}{x^n} & f^{(n)}(2) &= \frac{(-1)^{n-1}(n-1)!}{2^n} \\ a_n &= \frac{(-1)^{n-1}}{n2^n} \end{array}$$

For example, we can approximate $\ln(x)$ around 2 as; $\ln(x) = \ln(2) + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n2^n}(x-2)^n$.

It's important to note the bounds for this, as we start from $n = 1$, and manually state the constant term, since that doesn't fall within the "pattern" we had. We use the same technique to calculate the

radius of convergence, as follows;

$$\begin{aligned}\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{\frac{(x-2)^{n+1}}{(n+1)2^{n+1}}}{\frac{(x-2)^n}{n2^n}} \right| \\ &= \lim_{n \rightarrow \infty} \frac{n}{n+1} \cdot \frac{|x-2|}{2} \\ &= \frac{|x-2|}{2}\end{aligned}$$

And then, by setting $\frac{|x-2|}{2} < 1$, we have a radius of convergence of 2, around $x = 2$. Generally, the radius is limited by the closest singularity (where a point is not defined, or isn't well-behaved).

Taylor Series Error Term

In order to ensure termination, instead of summing to ∞ , we often truncate at some arbitrary k^{th} term.

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = \sum_{n=0}^k \frac{f^{(n)}(a)}{n!} (x-a)^n + \underbrace{\frac{f^{(k+1)}(c)}{(k+1)!} (x-a)^{k+1}}_{\text{Lagrange error term}}$$

The Lagrange error term replaces the tail of the infinite series, from, which would otherwise be a sum from $k+1$ to ∞ . The constant c lies between x , and a . By taking the bound $a < c < x$, or $x < c < a$, it can be used to generate the worst-case error.

The mean value theorem states that there is some point c between x , and a , such that $f'(c) = \frac{f(x)-f(a)}{x-a}$, therefore we can get $f(x) = f(a) + (x-a)f'(c)$.

Cauchy Error Term

Define the set of partial sums of the Taylor series as $F_k(t) = \sum_{n=0}^k \frac{f^{(n)}(t)}{n!} (x-t)^n$, where t is the offset.

It follows that $F_k(x) = f(x)$ for all k (justified as every term becomes 0, other than the first term, $n=0$, where it's $\frac{f(x)}{0!}$). Therefore the error term is derived as follows, where $R_k(x)$ is the remainder after k terms;

$$F_k(x) - F_k(a) = f(x) - \sum_{n=0}^k \frac{f^{(n)}(a)}{n!} (x-a)^n = R_k(x)$$

Power Series Solutions of ODE

When we have a differential equation in the form $\frac{dy}{dx} = ky$, for a constant k , we can consider the power series for y . By differentiating term by term, we can compare coefficients;

$$\frac{dy}{dx} \equiv \frac{d}{dx} \sum_{i=0}^{\infty} a_i x^i \equiv \sum_{i=1}^{\infty} a_i i x^{i-1} \equiv ky \equiv k \sum_{i=0}^{\infty} a_i x^i \equiv k \sum_{i=1}^{\infty} a_{i-1} x^{i-1} \equiv \sum_{i=1}^{\infty} k a_{i-1} x^{i-1}$$

By matching coefficients of x^{i-1} , where $i \geq 1$, we have $k a_{i-1} = a_i i$, therefore we end up with the relation $a_i = \frac{k}{i} a_{i-1}$. By "expanding" this, we end up with $a_i = \frac{k}{i} a_{i-1} = \frac{k}{i} \cdot \frac{k}{i-1} a_{i-2} = \dots = \frac{k^i}{i!} a_0$.

If we were to have some boundary condition, such as when $x=0$, $y=1$, we know that $y(0) = a_0$, hence $a_0 = 1$. Therefore, this can be solved to the following;

$$y = \sum_{i=0}^{\infty} \frac{(kx)^i}{i!} = e^{kx}$$

Linear Algebra

Linear Equation Systems

For brevity, instead of writing;

$$ax_1 + bx_2 + cx_3 = j \quad (1)$$

$$dx_1 + ex_2 + fx_3 = k \quad (2)$$

$$gx_1 + hx_2 + ix_3 = l \quad (3)$$

We will write $\mathbf{A}\vec{x} = \vec{y}$, where $\mathbf{A} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$, and $\vec{y} = \begin{bmatrix} j \\ k \\ l \end{bmatrix}$

In general, for a real linear equation system, we have the following possible outcomes;

- no solution

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ 2 & 0 & 3 \end{bmatrix}, \text{ and } \vec{y} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

By adding equations (1), and (2), we get $2x_1 + 3x_3 = 5$, which contradicts equation (3), hence it is **inconsistent**, and therefore has no solutions.

- unique solution

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ 0 & 1 & 1 \end{bmatrix}, \text{ and } \vec{y} = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$$

By doing equation (1) - equation (3), we get $x_1 = 1$. Adding equations (1), and (2), we end up with $2x_1 + 3x_3 = 5$, however, we know the value of x_1 , therefore it follows that $x_3 = 1$. Finally, by using this value in equation (3), we have $x_2 = 1$, therefore the only solution is;

$$\vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- infinite solutions

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ 2 & 0 & 3 \end{bmatrix}, \text{ and } \vec{y} = \begin{bmatrix} 3 \\ 2 \\ 5 \end{bmatrix}$$

Note that this is the same \mathbf{A} from the first example. However, this time adding equations (1), and (2), is consistent with equation (3). If we let $x_3 = \lambda$, such that x_3 is a free variable, with $\lambda \in \mathbb{R}$, we can get both x_1 , and x_2 in terms of λ . By doing equations (1) - (2), we get $2x_2 - x_3 = 1$, hence $x_2 = \frac{1}{2} + \frac{1}{2}\lambda$. By doing equations (1) + (2), we get $2x_1 + 3x_3 = 5$, which leads to $x_1 = \frac{5}{2} - \frac{3}{2}\lambda$. As such, we have an infinite set of solutions, given $\lambda \in \mathbb{R}$;

$$\vec{x} = \begin{bmatrix} \frac{5}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} -\frac{3}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

Groups

With a set G , and some operation $\otimes : G \mapsto G$, the criteria for (G, \otimes) being a **group** are (1 - 4), and (1 - 5) for an **Abelian group**;

1. **closure**

$$\forall x, y \in G [x \otimes y \in G]$$

2. **associativity**

$$\forall x, y, z \in G [(x \otimes y) \otimes z = x \otimes (y \otimes z)]$$

3. **neutral element**

$$\exists e \in G [\forall x \in G [x \otimes e = x \wedge e \otimes x = x]]$$

4. inverse element

$$\forall x \in G[\exists y \in G[x \otimes y = e \wedge y \otimes x = e]]$$

often written as $y = x^{-1}$, note that this does **not** mean $y = \frac{1}{x}$, defined with respect to \otimes

5. commutative

$$\forall x, y \in G[x \otimes y = y \otimes x]$$

For example, we can say $(\mathbb{Z}, +)$ is a group, but $(\mathbb{N}, +)$ is not, since the latter doesn't have an inverse element; for example $-5 \notin \mathbb{N}$, so 5 doesn't have an inverse. While $(\mathbb{R} \setminus \{0\}, \cdot)$ is an Abelian group, (\mathbb{R}, \cdot) isn't a group as 0 doesn't have an inverse element ($\neg \exists x \in \mathbb{R}[0 \cdot x = 1]$). $(\mathbb{R}^n, +)$ is an Abelian groups under addition, as long as addition is defined component-wise.

Matrices

We can defined the set of all real-valued $m \times n$ matrices ($m, n \in \mathbb{N}$), where m is the number of rows, and n is the number of columns, as $\mathbb{R}^{m \times n}$. We can refer to each element as $a_{i,j}$, where $i \in [1, m]$, and $j \in [1, n]$, and $a_{i,j} \in \mathbb{R}$; thus giving us the following rectangular matrix;

$$\mathbf{A} = \begin{bmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,n} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \cdots & a_{m,n} \end{bmatrix}$$

A $\mathbb{R}^{1 \times n}$ matrix is referred to as a **row (vector)**, and a $\mathbb{R}^{m \times 1}$ matrix is referred to as a **column (vector)**, with component-wise addition, we are able to say $(\mathbb{R}^{m \times n}, +)$ is a group.

Multiplication on matrices, $\mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{B} \in \mathbb{R}^{n \times k}, \mathbf{C} \in \mathbb{R}^{m \times k}$, where $\mathbf{C} = \mathbf{AB}$, is defined as;

$$c_{i,j} = \sum_{l=1}^n a_{i,l}b_{l,j}, \text{ with } i \in [1, m], \text{ and } j \in [1, k]$$

In general, matrix multiplication is not commutative, hence $\mathbf{AB} \neq \mathbf{BA}$. However, it is associative, and distributive (given that the dimensions match).

The identity matrix, written as $\mathbf{I}_n \in \mathbb{R}^{n \times n}$, is all 0s, other than $a_{i,i} = 1$, where $i \in [1, n]$. The identity matrix has the following property; $\mathbf{I}_m \mathbf{A} = \mathbf{A} = \mathbf{AI}_n$, where $\mathbf{A} \in \mathbb{R}^{m \times n}$.

If a square matrix ($\mathbf{A} \in \mathbb{R}^{n \times n}$) is non-singular, we can find an inverse for it, $\mathbf{A}^{-1} \in \mathbb{R}^{n \times n}$, such that $\mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n$. However, not every matrix has an inverse. We also have the result $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$.

We can define the transpose of $\mathbf{A} \in \mathbb{R}^{n \times m}$ as $\mathbf{B} = \mathbf{A}^\top \in \mathbb{R}^{m \times n}$, where $b_{j,i} = a_{i,j}$. On the rectangular matrix representation, we are essentially mirroring it on the leading diagonal. A **square** matrix is symmetric $\Leftrightarrow \mathbf{A} = \mathbf{A}^\top$. The general results are $(\mathbf{A}^\top)^\top = \mathbf{A}$, $(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top$, and $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$. Also, if \mathbf{A} is non-singular, then $(\mathbf{A}^{-1})^\top = (\mathbf{A}^\top)^{-1}$.

A multiplication by scalar $\lambda \in \mathbb{R}$, means that for $\mathbf{A} \in \mathbb{R}^{m \times n}$, and $\mathbf{B} = \lambda \mathbf{A}$, it follows that $b_{i,j} = \lambda a_{i,j}$. This operation is distributive, and also associative.

Solving Linear Equation Systems

Given the general form of an equation system as;

$$\begin{array}{cccccc} a_{1,1}x_1 & + & a_{1,2}x_1 & + & \cdots & + & a_{1,n}x_n & = & b_1 \\ a_{2,1}x_1 & + & a_{2,2}x_2 & + & \cdots & + & a_{2,n}x_n & = & b_2 \\ \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m,1}x_1 & + & a_{m,2}x_2 & + & \cdots & + & a_{m,n}x_n & = & b_m \end{array}$$

We can easily represent it as $\mathbf{A}\vec{x} = \vec{b}$, where $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\vec{x} \in \mathbb{R}^{n \times 1}$, and $\vec{b} \in \mathbb{R}^{m \times 1}$.

For example, consider a simple linear equation system as follows;

$$\mathbf{A}\vec{x} = \begin{bmatrix} 1 & 0 & 8 & -4 \\ 0 & 1 & 2 & 12 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 42 \\ 8 \end{bmatrix}$$

Let us also say $\mathbf{A} = [\vec{c}_1 \mid \vec{c}_2 \mid \vec{c}_3 \mid \vec{c}_4]$, where \vec{c}_i represents the i^{th} column of \mathbf{A} , hence $\vec{b} = \sum_{i=1}^4 x_i \vec{c}_i$.

In order to find a particular solution, we can easily create \vec{b} as a linear combination of \vec{c}_1 , and \vec{c}_2 , such that $\vec{b} = 42\vec{c}_1 + 8\vec{c}_2 + 0\vec{c}_3 + 0\vec{c}_4$, therefore we can get a **particular** solution; $\vec{x} = [42 \ 8 \ 0 \ 0]^T$. This doesn't capture all the solutions of this equation system - if there are more unknowns than equations, then there are an infinite number of solutions. Recall that $\vec{b} + \vec{0} = \vec{b}$, hence we can attempt to find non-trivial $\vec{0}$ s to generate the set of all solutions.

Using the fact that \mathbf{A} is already in **row echelon form**, we can try say that $\vec{c}_3 = 8\vec{c}_1 + 2\vec{c}_2$, therefore, it follows that $\vec{0} = 8\vec{c}_3 + 2\vec{c}_2 - \vec{c}_3 + 0\vec{c}_4$, as such we've managed to create a zero solution when $\vec{x} = [8 \ 2 \ -1 \ 0]^T$. We can do the same with \vec{c}_4 , such that $\vec{c}_4 = -4\vec{c}_1 + 12\vec{c}_2 \Leftrightarrow \vec{0} = -4\vec{c}_1 + 12\vec{c}_2 + 0\vec{c}_3 - \vec{c}_4$, hence $\vec{x} = [-4 \ 12 \ 0 \ -1]^T$ is also a zero solution. With these two $\vec{0}$ s, we can say that the general solution is as follows (where $\lambda_1, \lambda_2 \in \mathbb{R}$);

$$\vec{x} = \begin{bmatrix} 42 \\ 8 \\ 0 \\ 0 \end{bmatrix} + \lambda_1 \begin{bmatrix} 8 \\ 2 \\ -1 \\ 0 \end{bmatrix} + \lambda_2 \begin{bmatrix} -4 \\ 12 \\ 0 \\ -1 \end{bmatrix}$$

In this solution, we first find a **particular** solution for $\mathbf{A}\vec{x} = \vec{b}$, and then combine this with **all** solutions to $\mathbf{A}\vec{x} = \vec{0}$. This was fairly trivial to solve, as the matrix was already in a special form. Generally, equation systems won't be given to us like this, however we can transform it into this form by **Gaussian elimination**.

Elementary Transformations

We are allowed to apply the following operations, which let us transform the equation system into a simpler state, but preserve the solution set;

- swap two equations (which means we can swap rows in the matrix)
- multiply an equation (matrix row) by a non-zero constant $\lambda \in \mathbb{R} \setminus \{0\}$
- add (or subtract) an equation (row) to / from another equation (row)

We can say that a matrix is in **row echelon form (REF)**, if all zero rows are below the non-zero rows, and the pivot of each row is **strictly** to the right of the pivot in the row above. The pivot of a row is defined as the first non-zero element in a given row, reading left-to-right.

Consider the following system of equations, and the result of it after Gaussian elimination;

$$\begin{array}{rrrrrrrrcl} -2x_1 & + & 4x_2 & - & 2x_3 & - & x_4 & + & 4x_5 & = & -3 \\ 4x_1 & - & 8x_2 & + & 3x_3 & - & 3x_4 & + & x_5 & = & 2 \\ x_1 & - & 2x_2 & + & x_3 & - & x_4 & + & x_5 & = & 0 \\ x_1 & - & 2x_2 & + & 0x_3 & - & 3x_4 & + & 4x_5 & = & a \\ & & & & & & & & & & \rightsquigarrow \\ x_1 & - & 2x_2 & + & x_3 & - & x_4 & + & x_5 & = & 0 \\ & & & & x_3 & - & x_4 & + & 3x_5 & = & -2 \\ & & & & & & x_4 & - & 2x_5 & = & 1 \\ & & & & & & & & 0 & = & a + 1 \end{array}$$

We can now work on the augmented matrix to transform it into **REF**;

$$\begin{aligned}
& \left[\begin{array}{ccccc|c} -2 & 4 & -2 & -1 & 4 & -3 \\ 4 & -8 & 3 & -3 & 1 & 2 \\ 1 & -2 & 1 & -1 & 1 & 0 \\ 1 & -2 & 0 & -3 & 4 & a \end{array} \right] \begin{array}{l} \leftrightarrow R_3 \\ \\ \leftrightarrow R_1 \end{array} \rightsquigarrow \left[\begin{array}{ccccc|c} 1 & -2 & 1 & -1 & 1 & 0 \\ 4 & -8 & 3 & -3 & 1 & 2 \\ -2 & 4 & -2 & -1 & 4 & -3 \\ 1 & -2 & 0 & -3 & 4 & a \end{array} \right] \begin{array}{l} \\ -4R_1 \\ +2R_1 \\ -R_1 \end{array} \rightsquigarrow \\
& \left[\begin{array}{ccccc|c} 1 & -2 & 1 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 & -3 & 2 \\ 0 & 0 & 0 & -3 & 6 & -3 \\ 0 & 0 & -1 & -2 & 3 & a \end{array} \right] -R_2 \rightsquigarrow \left[\begin{array}{ccccc|c} 1 & -2 & 1 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 & -3 & 2 \\ 0 & 0 & 0 & -3 & 6 & -3 \\ 0 & 0 & 0 & -3 & 6 & a-2 \end{array} \right] -R_3 \rightsquigarrow \\
& \left[\begin{array}{ccccc|c} 1 & -2 & 1 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 & -3 & 2 \\ 0 & 0 & 0 & -3 & 6 & -3 \\ 0 & 0 & 0 & 0 & 0 & a+1 \end{array} \right] \begin{array}{l} \\ \cdot -1 \\ \cdot -\frac{1}{3} \\ \end{array} \rightsquigarrow \left[\begin{array}{ccccc|c} 1 & -2 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 3 & -2 \\ 0 & 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 & 0 & a+1 \end{array} \right]
\end{aligned}$$

It follows that there is only a solution when $a = -1$. To clarify some definitions, we refer to a variable as being **basic**, if it corresponds to the pivot element of any given row, hence in this case, the basic variables are x_1, x_3, x_4 , and the rest of the variables are **free** (x_2 , and x_5 in our case). The pivot columns in our case (written as \vec{p}_i corresponding to x_i) are;

$$\vec{p}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \vec{p}_3 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \text{ and } \vec{p}_4 = \begin{bmatrix} -1 \\ -1 \\ 1 \\ 0 \end{bmatrix}$$

We implicitly set the **free** variables to 0 when we find our particular solution using this method.

$$x_1\vec{p}_1 + x_3\vec{p}_3 + x_4\vec{p}_4 = \vec{b} \Leftrightarrow x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -1 \\ -1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ -2 \\ 1 \\ 0 \end{bmatrix} \Leftrightarrow \vec{x} = \begin{bmatrix} 2 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix}$$

We can get $x_4 = 1$ straight away, and from that we can get $x_3 = -1$, and also $x_1 = 2$. By using the same method as before, where we try create the non-pivot columns as a linear combination of the pivot columns, we can get the following general solution ($\forall \lambda_1, \lambda_2 \in \mathbb{R}$) - note that we could just as easily negate all the values in the column vectors, since our λ_1, λ_2 can take any real value;

$$\vec{x} = \begin{bmatrix} 2 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix} + \lambda_1 \begin{bmatrix} -2 \\ -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \lambda_2 \begin{bmatrix} -2 \\ 0 \\ 1 \\ -2 \\ -1 \end{bmatrix}$$

Minus-1 Trick

We say that some equation system is in **reduced row echelon form (RREF)** (this may also be referred to as row-reduced echelon form, or row canonical form in other texts), if it is already in row echelon form, every pivot item is 1, and the pivot is the only non-zero entry in the column. Consider the following matrix, with the pivots coloured purple, and the matrix after adding rows of the form $[0 \ \cdots \ 0 \ \textcolor{red}{-1} \ 0 \ \cdots \ 0]$, to create a square matrix. By reading the columns containing the new **-1**s, we can get all solutions for $\mathbf{A}\vec{x} = \vec{0}$, once again $\forall \lambda_1, \lambda_2 \in \mathbb{R}$;

$$\mathbf{A} = \begin{bmatrix} \textcolor{violet}{1} & 3 & 0 & 0 & 3 \\ 0 & 0 & \textcolor{violet}{1} & 0 & 9 \\ 0 & 0 & 0 & \textcolor{violet}{1} & -4 \end{bmatrix} \quad \mathbf{A}' = \begin{bmatrix} \textcolor{violet}{1} & 3 & 0 & 0 & 3 \\ 0 & \textcolor{red}{-1} & 0 & 0 & 0 \\ 0 & 0 & \textcolor{violet}{1} & 0 & 9 \\ 0 & 0 & 0 & \textcolor{violet}{1} & -4 \\ 0 & 0 & 0 & 0 & \textcolor{red}{-1} \end{bmatrix} \quad \vec{x} = \lambda_1 \begin{bmatrix} 3 \\ -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \lambda_2 \begin{bmatrix} 3 \\ 0 \\ 9 \\ -4 \\ -1 \end{bmatrix}$$

Gaussian Elimination to Calculate the Inverse

This is a fairly simple technique for calculating the inverse matrix, where we essentially augment the original matrix with the identity, and apply Gaussian elimination to get the identity on the left hand side; $[\mathbf{A} \mid \mathbf{I}_n] \rightsquigarrow \dots \rightsquigarrow [\mathbf{I}_n \mid \mathbf{A}^{-1}]$.

This is one of the parts in this module that are fairly easy to understand, but just requires practice for the exams. This technique is applied to a simple $\mathbf{A} \in \mathbb{R}^{2 \times 2}$;

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right] \xrightarrow{-3R_1} \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & -2 & -3 & 1 \end{array} \right] \xrightarrow{-\frac{1}{2}} \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & 1 & \frac{3}{2} & -\frac{1}{2} \end{array} \right] \xrightarrow{-2R_2} \left[\begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & \frac{3}{2} & -\frac{1}{2} \end{array} \right]$$

Vector Spaces

We define a real-valued **vector space** (\mathbb{R} -vector space) is a set V with two operations;

- $+: V \times V \mapsto V$ inner operation
- $\cdot: \mathbb{R} \times V \mapsto V$ outer operation

The following properties hold;

- $(V, +)$ is an Abelian group
- the outer operation is distributive

$$\lambda \cdot (\vec{x} + \vec{y}) = \lambda \cdot \vec{x} + \lambda \cdot \vec{y} \quad \forall \lambda \in \mathbb{R}, \vec{x}, \vec{y} \in V$$

$$(\lambda + \psi) \cdot \vec{x} = \lambda \cdot \vec{x} + \psi \cdot \vec{x} \quad \forall \lambda, \psi \in \mathbb{R}, \vec{x} \in V$$
- the outer operation is associative

$$\lambda \cdot (\psi \cdot \vec{x}) = (\lambda \cdot \psi) \cdot \vec{x} \quad \forall \lambda, \psi \in \mathbb{R}, \vec{x} \in V$$
- the outer operation has a neutral element of 1

$$1 \cdot \vec{x} = \vec{x} \quad \forall \vec{x} \in V$$

Examples of vector spaces are $V = \mathbb{R}^n$, $V = \mathbb{R}^{m \times n}$, and $V = \mathbb{C}$ (the standard representation of the complex numbers).

Vector Subspaces

Suppose we have $U \subset V$, where V is a vector space, then U is a vector / linear subspace if it is a vector space with the same outer, and inner operations, but restricted to $U \times U$, and $\mathbb{R} \times U$, respectively.

For us to show that U is a vector subspace, we must show the following properties;

- $U \neq \emptyset$, especially have to show $\vec{0} \in U$
- closure of U
 - outer operation: $\lambda \vec{x} \in U$ $\forall \lambda \in \mathbb{R}, \vec{x} \in U$
 - inner operation: $\vec{x} + \vec{y} \in U$ $\forall \vec{x}, \vec{y} \in U$

We have trivial vector subspaces of the vector space V , namely V itself, and $\{\vec{0}\}$ (the subspace containing only the zero vector). The intersection of an arbitrary number of subspaces is also a subspace, and the intersection of all subspaces of V is referred to as the **linear hull** of V .

The solution space for a homogeneous linear equation system $\mathbf{A}\vec{x} = \vec{0}$, with n unknowns, is a subspace of \mathbb{R}^n . Note that a homogeneous equation means that it has the zero vector on the right hand side. The same also applies in reverse; such that for a vector subspace $U \subset V$, there is a homogeneous system which characterises U . In contrast, an inhomogeneous linear equation system $\mathbf{A}\vec{x} = \vec{b}$, where $\vec{b} \neq \vec{0}$ does not have a vector subspace for its solutions. Assume that it does have some subspace U , therefore, $\vec{0} \in U$ by definition. However, $\vec{0}$ doesn't satisfy the equation, since we know $\vec{b} \neq \vec{0}$.

Linear Independence

We can define a vector space V , with $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$, as being linearly dependent if, and only if, there exists a non-trivial solution (such that there exists at least one $\lambda_i \neq 0$) in the following equation;

$$\vec{0} = \sum_{i=1}^k \lambda_i \vec{x}_i$$

By the closure of vector spaces, a set of linearly dependent vectors contains some redundancy.

We can check if vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in V$ are linearly independent by constructing an augmented matrix; $\mathbf{A} = [\vec{x}_1 \mid \vec{x}_2 \mid \dots \mid \vec{x}_k]$. Once put into row echelon form, if all the columns are pivot columns, then it follows that it is linearly independent, otherwise the non-pivot columns can be made from the pivot columns to the left of it (therefore it is linearly dependent).

Consider the following vectors in \mathbb{R}^4 ;

$$\vec{x}_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \vec{x}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \text{ and } \vec{x}_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

We essentially want to solve $\lambda_1 \vec{x}_1 + \lambda_2 \vec{x}_2 + \lambda_3 \vec{x}_3 = \vec{0}$, which means we can set this up as follows;

$$\begin{array}{ccc} \left[\begin{array}{ccc|c} 1 & 1 & -1 & 0 \\ 2 & 1 & -2 & 0 \\ -3 & 0 & 1 & 0 \\ 4 & 2 & 1 & 0 \end{array} \right] \begin{array}{l} -2R_1 \\ +3R_1 \\ -4R_1 \end{array} & \rightsquigarrow & \left[\begin{array}{ccc|c} 1 & 1 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 3 & -2 & 0 \\ 0 & -2 & 5 & 0 \end{array} \right] \begin{array}{l} \\ +3R_2 \\ -2R_2 \end{array} & \rightsquigarrow & \left[\begin{array}{ccc|c} 1 & 1 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 5 & 0 \end{array} \right] \begin{array}{l} \\ \cdot -1 \\ \cdot -\frac{1}{2} \end{array} & \rightsquigarrow & \\ \left[\begin{array}{ccc|c} 1 & 1 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 5 & 0 \end{array} \right] -5R_3 & \rightsquigarrow & \left[\begin{array}{ccc|c} 1 & 1 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] & & \end{array}$$

As every column here is a pivot column, such that every column is linearly independent of the ones to its left, it follows that there isn't a non-trivial solution, as the only solution is $\lambda_1 = \lambda_2 = \lambda_3 = 0$, therefore the vectors are linearly independent.

Linear Combinations (or Something?)

Suppose there is some vector space V , with k linearly independent vectors $\vec{b}_1, \vec{b}_2, \dots, \vec{b}_k$, and m linear combinations, expressed as follows (with $j \in [1, m]$);

$$\vec{x}_j = \sum_{i=1}^k \lambda_{i,j} \vec{b}_i$$

We can then apply the same approach to test for linear independence;

$$\vec{0} = \sum_{j=1}^m \mu_j \vec{x}_j = \sum_{j=1}^m \mu_j \sum_{i=1}^k \lambda_{i,j} \vec{b}_i = \sum_{j=1}^m \sum_{i=1}^k \mu_j \lambda_{i,j} \vec{b}_i = \sum_{i=1}^k \left(\sum_{j=1}^m \mu_j \lambda_{i,j} \right) \vec{b}_i$$

This means that $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_m$ only has linear independence if, and only if the following column vectors are linearly independent;

$$\vec{\lambda}_1 = \begin{bmatrix} \lambda_{1,1} \\ \vdots \\ \lambda_{k,1} \end{bmatrix} \in \mathbb{R}^k, \dots, \vec{\lambda}_m = \begin{bmatrix} \lambda_{1,m} \\ \vdots \\ \lambda_{k,m} \end{bmatrix} \in \mathbb{R}^k$$

Consider the following example, with a set of linearly independent vectors $\vec{b}_1, \vec{b}_2, \vec{b}_3, \vec{b}_4 \in \mathbb{R}^3$, and;

$$\begin{array}{cccccc}
\vec{b}_1 & - & 2\vec{b}_2 & + & \vec{b}_3 & - & \vec{b}_4 & = & \vec{x}_1 \\
-4\vec{b}_1 & - & 2\vec{b}_2 & + & 0\vec{b}_3 & + & 4\vec{b}_4 & = & \vec{x}_2 \\
2\vec{b}_1 & + & 3\vec{b}_2 & - & \vec{b}_3 & - & 3\vec{b}_4 & = & \vec{x}_3 \\
17\vec{b}_1 & - & 10\vec{b}_2 & + & 11\vec{b}_3 & + & \vec{b}_4 & = & \vec{x}_4
\end{array}$$

Now, to check whether $\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4$ are linearly independent, we need to find the reduced row echelon form of the coefficient matrix;

$$\begin{array}{ccc}
\begin{bmatrix} 1 & -4 & 2 & 17 \\ -2 & -2 & 3 & -10 \\ 1 & 0 & -1 & 11 \\ -1 & 4 & -3 & 1 \end{bmatrix} \begin{array}{l} +2R_1 \\ -R_1 \\ +R_1 \end{array} & \rightsquigarrow & \begin{bmatrix} 1 & -4 & 2 & 17 \\ 0 & -10 & 7 & 24 \\ 0 & 4 & -3 & -6 \\ 0 & 0 & -1 & 18 \end{bmatrix} \begin{array}{l} \cdot -\frac{1}{10} \\ \\ \cdot -1 \end{array} \rightsquigarrow \begin{bmatrix} 1 & -4 & 2 & 17 \\ 0 & 1 & -\frac{7}{10} & -\frac{24}{10} \\ 0 & 4 & -3 & -6 \\ 0 & 0 & 1 & -18 \end{bmatrix} \begin{array}{l} \\ -4R_2 \\ \\ \end{array} \rightsquigarrow \\
\begin{bmatrix} 1 & -4 & 2 & 17 \\ 0 & 1 & -\frac{7}{10} & -\frac{24}{10} \\ 0 & 0 & -\frac{2}{10} & \frac{36}{10} \\ 0 & 0 & 1 & -18 \end{bmatrix} \begin{array}{l} \\ +\frac{2}{10}R_4 \end{array} & \rightsquigarrow & \begin{bmatrix} 1 & -4 & 2 & 17 \\ 0 & 1 & -\frac{7}{10} & -\frac{24}{10} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -18 \end{bmatrix} \begin{array}{l} \\ \\ \leftrightarrow R_4 \\ \leftrightarrow R_3 \end{array} \rightsquigarrow \begin{bmatrix} 1 & -4 & 2 & 17 \\ 0 & 1 & -\frac{7}{10} & -\frac{24}{10} \\ 0 & 0 & 1 & -18 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} -2R_3 \\ +\frac{7}{10}R_3 \\ \\ \end{array} \rightsquigarrow \\
\begin{bmatrix} 1 & -4 & 0 & 53 \\ 0 & 1 & 0 & -15 \\ 0 & 0 & 1 & -18 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} +4R_2 \\ \\ \\ \end{array} & \rightsquigarrow & \begin{bmatrix} 1 & 0 & 0 & -7 \\ 0 & 1 & 0 & -15 \\ 0 & 0 & 1 & -18 \\ 0 & 0 & 0 & 0 \end{bmatrix}
\end{array}$$

Since we have a non-pivot column, it follows that it's solvable non-trivially, and therefore it is linearly dependent. Therefore, the \vec{x} vectors are linear dependent since \vec{x}_4 lies in the span of $\vec{x}_1, \vec{x}_2, \vec{x}_3$.

Basis

Given a finite number of vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in V$, we can say that any vector of the following form, is a linear combination of the vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ (given $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$);

$$\vec{v} = \sum_{i=1}^k \lambda_i \vec{x}_i \in V$$

Suppose we have $A \subset V$, and $A = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$, if we can generate every vector $\vec{v} \in V$, as a linear combination of vectors in A , then A is a **generating set** / **span** of the vector space V . This is denoted as $V = [A] = [\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k]$.

We can define a generating set of V as a **basis** of V , if it is linearly independent, and minimal. We define a generating set of V minimal if there isn't a smaller set which spans V . With a real vector space V , and $B \subset V$, $B \neq \emptyset$, B is a basis of $V \Leftrightarrow B$ is a minimal generating set $\Leftrightarrow B$ is a maximal linearly independent subset of V .

It is also equivalent to say that every $\vec{v} \in V$ is a **unique** linear combination of vectors from B ;

$$\vec{v} = \sum_{i=1}^k \lambda_i \vec{b}_i = \sum_{i=1}^k \mu_i \vec{b}_i \Rightarrow \forall i \in [1, k] [\mu_i = \lambda_i]$$

While bases aren't unique, the number of elements (or $\dim(V)$) is constant for a vector space V . In \mathbb{R}^3 , we refer to the **standard** basis as the following set of vectors, thus $\dim(\mathbb{R}^3) = 3$, obviously;

$$B = \left[\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right]$$

With a vector subspace $U \subset V$, we can set an upper bound on the dimensions as $\dim(U) \leq \dim(V)$, and $\dim(U) = \dim(V) \Leftrightarrow U = V$.

Consider the following subspace $U \subset \mathbb{R}^5$, spanned by;

$$\vec{x}_1 = \begin{bmatrix} 1 \\ 2 \\ -1 \\ -1 \\ -1 \end{bmatrix}, \vec{x}_2 = \begin{bmatrix} 2 \\ -1 \\ 1 \\ 2 \\ -2 \end{bmatrix}, \vec{x}_3 = \begin{bmatrix} 3 \\ -4 \\ 3 \\ 5 \\ -3 \end{bmatrix}, \text{ and } \vec{x}_4 = \begin{bmatrix} -1 \\ 8 \\ -5 \\ -6 \\ 1 \end{bmatrix}$$

As we're once again trying to determine linear independence, we end up with another homogeneous equation system, that can be represented by taking the augmented matrix of the vectors, and then apply Gaussian elimination on it, down to reduced row echelon form (still doing all of it, for practice);

$$\begin{aligned} & \begin{bmatrix} 1 & 2 & 3 & -1 \\ 2 & -1 & -4 & 8 \\ -1 & 1 & 3 & -5 \\ -1 & 2 & 5 & -6 \\ -1 & -2 & -3 & 1 \end{bmatrix} \begin{matrix} -2R_1 \\ +R_1 \\ +R_1 \\ +R_1 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & 3 & -1 \\ 0 & -5 & -10 & 10 \\ 0 & 3 & 6 & -6 \\ 0 & 4 & 8 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \cdot \frac{1}{3} \rightsquigarrow \begin{bmatrix} 1 & 2 & 3 & -1 \\ 0 & -5 & -10 & 10 \\ 0 & 1 & 2 & -2 \\ 0 & 4 & 8 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} +5R_3 \\ -4R_3 \end{matrix} \rightsquigarrow \\ & \begin{bmatrix} 1 & 2 & 3 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} \rightarrow R_4 \\ \rightarrow R_2 \\ \rightarrow R_3 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & 3 & -1 \\ 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} +R_3 \\ +2R_3 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} -2R_2 \end{matrix} \rightsquigarrow \\ & \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

From this, we can see that the vectors corresponding to the pivot columns are \vec{x}_1 , \vec{x}_2 , and \vec{x}_4 , hence we can say that $[\vec{x}_1, \vec{x}_2, \vec{x}_4]$ forms a basis of U .

On the other hand, we can try to find a simple basis of U , which means we want as many coordinates equal to 0 as possible. We can do this by writing the vectors as **rows** in the matrix, and then apply the same Gaussian elimination;

$$\begin{aligned} & \begin{bmatrix} 1 & 2 & -1 & -1 & -1 \\ 2 & -1 & 1 & 2 & -2 \\ 3 & -4 & 3 & 5 & -3 \\ -1 & 8 & -5 & -6 & 1 \end{bmatrix} \begin{matrix} -2R_1 \\ -3R_1 \\ +R_1 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & -1 & -1 & -1 \\ 0 & -5 & 3 & 4 & 0 \\ 0 & -10 & 6 & 8 & 0 \\ 0 & 10 & -6 & -7 & 0 \end{bmatrix} \begin{matrix} +R_3 \end{matrix} \rightsquigarrow \\ & \begin{bmatrix} 1 & 2 & -1 & -1 & -1 \\ 0 & -5 & 3 & 4 & 0 \\ 0 & -10 & 6 & 8 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \begin{matrix} +2R_2 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & -1 & -1 & -1 \\ 0 & -5 & 3 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \begin{matrix} \cdot -\frac{1}{5} \\ \leftrightarrow R_4 \\ \leftrightarrow R_3 \end{matrix} \rightsquigarrow \\ & \begin{bmatrix} 1 & 2 & -1 & -1 & -1 \\ 0 & 1 & -\frac{3}{5} & -\frac{4}{5} & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} +R_3 \\ +\frac{4}{5}R_3 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 2 & -1 & 0 & -1 \\ 0 & 1 & -\frac{3}{5} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{matrix} -2R_2 \end{matrix} \rightsquigarrow \\ & \begin{bmatrix} 1 & 0 & \frac{1}{5} & 0 & -1 \\ 0 & 1 & -\frac{3}{5} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad U = \left[\begin{bmatrix} 1 \\ 0 \\ \frac{1}{5} \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -\frac{3}{5} \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right] \end{aligned}$$

By reading the rows with the leading 1s, we have a simple basis; $U = [\vec{b}_1, \vec{b}_2, \vec{b}_3]$

Rank

The rank of a matrix is the number of linearly independent columns, denoted $\text{rk}(\mathbf{A})$, where $\mathbf{A} \in \mathbb{R}^{m \times n}$. We are also given that the column rank is the same as the row rank, hence $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{A}^\top)$.

The columns of \mathbf{A} span a subspace $U \in \mathbb{R}^m$, with $\dim(U) = \text{rk}(\mathbf{A})$, and the rows span a subspace $W \in \mathbb{R}^n$, with $\dim(W) = \text{rk}(\mathbf{A})$. A basis of U , and W , can be found by applying Gaussian elimination to \mathbf{A} , and \mathbf{A}^\top respectively.

$\mathbf{A} \in \mathbb{R}^{n \times n}$ is invertible (regular) if, and only if, $\text{rk}(\mathbf{A}) = n$. For all $\mathbf{A} \in \mathbb{R}^{m \times n}$, and all $\vec{b} \in \mathbb{R}^m$, the equation system $\mathbf{A}\vec{x} = \vec{b}$ can only be solved if $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{A} \mid \vec{b})$. The subspace of solutions for $\mathbf{A}\vec{x} = \vec{0}$ (**kernel** / **nullspace**) has dimension $n - \text{rk}(\mathbf{A})$. A matrix is said to have **full rank** if $\text{rk}(\mathbf{A}) = \min(m, n)$, otherwise it is said to be **rank deficient**.

For example, to calculate the rank of \mathbf{A} , as below, we apply Gaussian elimination to row echelon form, and look at the number of pivot columns (which corresponds to the linearly independent columns);

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 \\ -2 & -3 & 1 \\ 3 & 5 & 0 \end{bmatrix} \xrightarrow{+2R_1, -3R_1} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 3 \\ 0 & -1 & -3 \end{bmatrix} \xrightarrow{+R_3} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

As we have two pivot columns, it follows that $\text{rk}(\mathbf{A}) = 2$.

Subspace Intersection

Suppose we have two subspaces $U_1 = [\vec{b}_1, \vec{b}_2, \dots, \vec{b}_k] \subset V$, and $U_2 = [\vec{c}_1, \vec{c}_2, \dots, \vec{c}_l] \subset V$. We're then interested in finding the basis of $U_1 \cap U_2$, which is all the $\vec{x} \in V$, such that $\vec{x} \in U_1 \wedge \vec{x} \in U_2$. For something to be in U_1 , or U_2 , it must be a linear combination of the basis vectors of the respective subspaces; hence we can get the following;

$$\sum_{i=1}^k \lambda_i \vec{b}_i = \vec{x} = \sum_{j=1}^l \mu_j \vec{c}_j \Leftrightarrow \sum_{i=1}^k \lambda_i \vec{b}_i - \sum_{j=1}^l \mu_j \vec{c}_j = \vec{0}$$

This gives a homogeneous equation system, where we want to solve;

$$\left[\vec{b}_1 \mid \vec{b}_2 \mid \dots \mid \vec{b}_k \mid -\vec{c}_1 \mid -\vec{c}_2 \mid \dots \mid -\vec{c}_l \right] \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_k \\ \mu_1 \\ \vdots \\ \mu_l \end{bmatrix} = \vec{0}$$

However, we only need to determine $\lambda_1, \lambda_2, \dots, \lambda_k$, or $\mu_1, \mu_2, \dots, \mu_l$, to find the intersection of U_1 , and U_2 .

Consider the example, with the following vector subspaces;

$$U_1 = \left[\begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right] \subset \mathbb{R}^4, \text{ and also } U_2 = \left[\begin{bmatrix} -1 \\ 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right] \subset \mathbb{R}^4$$

By generating the matrix shown above, we can now apply Gaussian elimination to reduced row echelon form;

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & -1 & -1 \\ 0 & 1 & 1 & -2 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \xrightarrow{-R_1} \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -2 & -1 \\ 0 & 1 & 1 & -2 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \xrightarrow{-R_2} \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -2 & -1 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \xrightarrow{-R_4} \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -2 & -1 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -2 & -1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \begin{matrix} \leftrightarrow R_4 \\ \leftrightarrow R_3 \end{matrix} \rightsquigarrow \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -2 & -1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

From this, we can determine that $\mu_2 = 0$, and that μ_1 is a free variable, since it doesn't have a corresponding pivot column. Therefore, it follows that $\vec{x} = \mu_1 \vec{c}_1$, hence we've determined that the basis of the intersection is as follows;

$$U_1 \cap U_2 = \left[\begin{bmatrix} -1 \\ 1 \\ 2 \\ 0 \end{bmatrix} \right]$$

Linear Mappings

A linear mapping preserves the structure of vector spaces. Therefore, for two \mathbb{R} -vector spaces V , W , and a mapping Φ (note that it can also be called a **vector space homomorphism**);

$$\Phi : V \mapsto W \text{ is } \mathbf{linear} \Leftrightarrow \forall \lambda, \mu \in \mathbb{R} \forall \vec{x}, \vec{y} \in V [\Phi(\lambda \vec{x} + \mu \vec{y}) = \lambda \Phi(\vec{x}) + \mu \Phi(\vec{y})]$$

Special cases of this are;

- **isomorphism**
- **endomorphism**
- **automorphism**
- **identity mapping**

$\Phi : V \mapsto W$ is linear, and bijective

$\Phi : V \mapsto V$ is linear

$\Phi : V \mapsto V$ is linear, and bijective

$\text{id}_V : V \mapsto V$ such that $\forall \vec{x} \in V [\text{id}_V(\vec{x}) = \vec{x}]$

For example, consider the mapping $\Phi : \mathbb{R}^2 \mapsto \mathbb{C}$, where $\Phi(\vec{x}) = x_1 + ix_2$. We can show that this is a linear mapping, by taking arbitrary $\vec{x}, \vec{y} \in \mathbb{R}^2$, and also arbitrary $\lambda \in \mathbb{R}$;

$$\begin{aligned} \Phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}\right) &= \Phi\left(\begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \end{bmatrix}\right) & \Phi\left(\lambda \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) &= \Phi\left(\begin{bmatrix} \lambda x_1 \\ \lambda x_2 \end{bmatrix}\right) \\ &= (x_1 + y_1) + i(x_2 + y_2) & &= (\lambda x_1) + i(\lambda x_2) \\ &= x_1 + ix_2 + y_1 + iy_2 & &= \lambda(x_1 + ix_2) \\ &= \Phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) + \Phi\left(\begin{bmatrix} y_1 \\ y_2 \end{bmatrix}\right) & &= \lambda \Phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) \end{aligned}$$

Image, and Kernel / Null Space