

# ML Programming assignment VIII

Name : 林欣妤

Student number : 314652023

October 26, 2025

## 1 Stochastic Differential Equations (SDE)

### 1.1 Definition

A **stochastic differential equation (SDE)** describes a process  $x_t \in \mathbb{R}^d$  whose infinitesimal change is influenced by both deterministic and random terms:

$$dx_t = \underbrace{f(x_t, t) dt}_{\text{drift}} + \underbrace{G(x_t, t) dW_t}_{\text{diffusion}}, \quad x(0) = x_0, \quad (1.1)$$

where

- $x_t$  is the stochastic process (unknown we want to solve for),
- $f \in \mathbb{R}^d$  is the drift vector,
- $G \in \mathbb{R}^{d \times d}$  is the diffusion matrix,
- $W_t$  is a standard Brownian motion (Wiener process).

The integral form of the SDE (Itô integral) is:

$$x_t = x_0 + \int_0^t f(x_s, s) ds + \int_0^t G(x_s, s) dW_s. \quad (1.2)$$

A stochastic process that solves an SDE is called an **Itô process**.

Drift and diffusion must satisfy certain regularity conditions such as Lipschitz continuity.

### 1.2 Special Cases

- Pure drift:  $dx_t = f(x_t, t) dt \rightarrow$  deterministic process
- Pure diffusion:  $dx_t = G(x_t, t) dW_t \rightarrow$  Brownian-like random motion

### 1.3 Stochastic Process

A stochastic process is a collection of random variables  $\{x_t\}_{t \in T}$  indexed by a set  $T$  (discrete or continuous), defined on a probability space, and taking values in  $\mathbb{R}^d$ .

- For each fixed  $t$ ,  $x_t(\omega)$  is a random variable
- For each  $\omega \in \Omega$ ,  $t \mapsto x_t(\omega)$  is called a **path** (realization)

## 1.4 Wiener Process (Brownian Motion)

$W_t \in \mathbb{R}^d$  is a continuous stochastic process satisfying:

- Initial condition:  $W_0 = 0$
- Stationary Gaussian increments:  $\Delta W_t = W(t + \Delta t) - W(t) \sim \mathcal{N}(0, \Delta t I)$
- Independent increments
- Continuous paths: with probability 1,  $t \mapsto W(t)$  is continuous

Properties:  $E[W(t)] = 0$ ,  $\text{Var}[W(t)] = t$ , and almost surely nowhere differentiable.

White noise can be interpreted as the (formal) derivative of Brownian motion:  $h(t) = \frac{dW}{dt}$ .

## 1.5 White Noise

A white noise process  $h(t) \in \mathbb{R}^d$  satisfies:

- $h(t)$  and  $h(t')$  are independent if  $t \neq t'$
- Gaussian process with zero mean and Dirac delta correlation:

$$E[h(t)] = 0, \quad E[h(t)h^T(s)] = \delta(t - s)I$$

Properties: almost everywhere discontinuous, unbounded, can take arbitrarily large values.

## 1.6 Interpretation of Stochastic Integral

$$\int_0^t G(x_s, s) dW_s = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} G(x_{t_k}, t_k) [W(t_{k+1}) - W(t_k)],$$

with  $0 = t_0 < t_1 < \dots < t_n = t$ , and  $W(t_{k+1}) - W(t_k) \sim \mathcal{N}(0, (t_{k+1} - t_k)I)$ .

Remark:  $\int_0^t dW_s = W_t - W_0 = W_t \sim \mathcal{N}(0, tI)$ .

## 1.7 Euler—Maruyama Method / Euler—Maruyama 方法

A forward-Euler-like scheme is used to numerically approximate the process:

Partition  $[0, T]$  into  $N$  equal subintervals of width  $\Delta t = T/N$ ,  $t_k = k\Delta t$ ,  $k = 0, \dots, N$ .

Set  $X_0 = x_0$ . Update rule:

$$X_{n+1} = X_n + f(X_n, t_n)\Delta t + G(X_n, t_n)\Delta W(t_n), \quad (1.3)$$

where  $\Delta W(t_n) = W(t_{n+1}) - W(t_n)$ , or equivalently

$$X_{n+1} = X_n + f(X_n, t_n)\Delta t + G(X_n, t_n)\Delta t Z(t_n), \quad (1.4)$$

with  $Z(t_i)$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

## References

- [1] Chat-GPT (Apply GPT to revise and correct the English content of the report)