

The Community Navigator programme for people with treatment resistant depression in community mental health teams in England: a randomised controlled trial

References

1. McIntyre RS, Alsuwaidan M, Baune BT, et al. Treatment-resistant depression: definition, prevalence, detection, management, and investigational interventions. *World Psychiatry* 2023; 22(3): 394-412.
2. Wang J, Mann F, Lloyd-Evans B, Ma R, Johnson S. Associations between loneliness and perceived social support and outcomes of mental health problems: a systematic review. *BMC psychiatry* 2018; 18(1): 156.
3. Stefanidou T, Evlat G, Gray H, et al. Associations between loneliness and outcomes of Common Mental Disorders (CMDs): A systematic review of longitudinal studies. *medRxiv* 2025: 2025.08.06.25333144.
4. Stefanidou T, Ambler G, Bartl G, et al. Randomised controlled trial of the Community Navigator programme to reduce loneliness and depression for adults with treatment-resistant depression in secondary community mental health services: trial protocol. *Trials* 2023; 24(1): 652.
5. Ma R, Mann F, Wang J, et al. The effectiveness of interventions for reducing subjective and objective social isolation among people with mental health problems: a systematic review. *Soc Psychiatry Psychiatr Epidemiol* 2020; 55(7): 839-76.
6. Kounali D, Button KS, Lewis G, et al. How much change is enough? Evidence from a longitudinal study on depression in UK primary care. *Psychol Med* 2022; 52(10): 1875-82.
7. Morriss R, Garland A, Nixon N, et al. Efficacy and cost-effectiveness of a specialist depression service versus usual specialist mental health care to manage persistent depression: a randomised controlled trial. *Lancet Psychiatry* 2016; 3(9): 821-31.