











Curve Fitting, Linear Regression, Regularization

Example

Handwritten Digit Recognition




Polynomial Curve Fitting




$$y(x, \mathbf{w}) = w_0 + w_1x + w_2x^2 + \dots + w_Mx^M = \sum_{j=0}^M w_jx^j$$

Sum-of-Squares Error Function




$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$


0th Order Polynomial




1st Order Polynomial




3rd Order Polynomial



9th Order Polynomial



Over-fitting




Root-Mean-Square (RMS) Error: $E_{\text{RMS}} = \sqrt{2E(\mathbf{w}^*)/N}$

Polynomial Coefficients

	$M = 0$	$M = 1$	$M = 3$	$M = 9$
w_0^*	0.19	0.82	0.31	0.35
w_1^*		-1.27	7.99	232.37
w_2^*			-25.43	-5321.83
w_3^*			17.37	48568.31
w_4^*				-231639.30
w_5^*				640042.26
w_6^*				-1061800.52
w_7^*				1042400.18
w_8^*				-557682.99
w_9^*				125201.43


Data Set Size: $N = 15$

9th Order Polynomial



Data Set Size: $N = 100$

9th Order Polynomial




Regularization

Penalize large coefficient values

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$

Linear Regression Result



Heat Flux	Skin Temperature	Heat Flux	Skin Temperature	Heat Flux	Skin Temperature
10.858	31.002	6.3221	31.581	4.3917	32.221
10.617	31.021	6.0325	31.618	4.2951	32.259
10.183	31.058	5.7429	31.674	4.2469	32.296
9.7003	31.095	5.5016	31.712	4.0056	32.334
9.652	31.133	5.2603	31.768	3.716	32.391
10.086	31.188	5.1638	31.825	3.523	32.448
9.459	31.226	5.0673	31.862	3.4265	32.505
8.3972	31.263	4.9708	31.919	3.3782	32.543
7.6251	31.319	4.8743	31.975	3.4265	32.6
7.1907	31.356	4.7777	32.013	3.3782	32.657
7.046	31.412	4.7295	32.07	3.3299	32.696
6.9494	31.468	4.633	32.126	3.3299	32.753
6.7081	31.524	4.4882	32.164	3.4265	32.791




Figure D.1. Measurements of heat flux and skin temperature of a person.

D.2.1 Least Square Method

Suppose we wish to fit the following linear model to the observed data:

$$f(x) = \omega_1 x + \omega_0, \quad (\text{D.3})$$

where ω_0 and ω_1 are parameters of the model and are called the **regression coefficients**. A standard approach for doing this is to apply the **method of least squares**, which attempts to find the parameters (ω_0, ω_1) that minimize the sum of the squared error

$$SSE = \sum_{i=1}^N [y_i - f(x_i)]^2 = \sum_{i=1}^N [y_i - \omega_1 x_i - \omega_0]^2, \quad (\text{D.4})$$

which is also known as the **residual sum of squares**.

This optimization problem can be solved by taking the partial derivative of E with respect to ω_0 and ω_1 , setting them to zero, and solving the corresponding system of linear equations.

$$\begin{aligned}\frac{\partial E}{\partial \omega_0} &= -2 \sum_{i=1}^N [y_i - \omega_1 x_i - \omega_0] = 0 \\ \frac{\partial E}{\partial \omega_1} &= -2 \sum_{i=1}^N [y_i - \omega_1 x_i - \omega_0] x_i = 0\end{aligned}\tag{D.5}$$

These equations can be summarized by the following matrix equation, which is also known as the **normal equation**:

$$\begin{pmatrix} N & \sum_i x_i \\ \sum_i x_i & \sum_i x_i^2 \end{pmatrix} \begin{pmatrix} \omega_0 \\ \omega_1 \end{pmatrix} = \begin{pmatrix} \sum_i y_i \\ \sum_i x_i y_i \end{pmatrix}.\tag{D.6}$$

Since $\sum_i x_i = 229.9$, $\sum_i x_i^2 = 1569.2$, $\sum_i y_i = 1242.9$, and $\sum_i x_i y_i = 7279.7$, the normal equations can be solved to obtain the following estimates for the parameters.

$$\begin{aligned} \begin{pmatrix} \hat{\omega}_0 \\ \hat{\omega}_1 \end{pmatrix} &= \begin{pmatrix} 39 & 229.9 \\ 229.9 & 1569.2 \end{pmatrix}^{-1} \begin{pmatrix} 1242.9 \\ 7279.7 \end{pmatrix} \\ &= \begin{pmatrix} 0.1881 & -0.0276 \\ -0.0276 & 0.0047 \end{pmatrix} \begin{pmatrix} 1242.9 \\ 7279.7 \end{pmatrix} \\ &= \begin{pmatrix} 33.1699 \\ -0.2208 \end{pmatrix} \end{aligned}$$

$$f(x) = 33.17 - 0.22x.$$

Figure D.2 shows the line corresponding to this model.






Figure D.2. A linear model that fits the data given in Figure D.1.

Regularization: $\ln \lambda = -18$



Regularization: $\ln \lambda = 0$



Polynomial Coefficients

	$\ln \lambda = -\infty$	$\ln \lambda = -18$	$\ln \lambda = 0$
w_0^*	0.35	0.35	0.13
w_1^*	232.37	4.74	-0.05
w_2^*	-5321.83	-0.77	-0.06
w_3^*	48568.31	-31.97	-0.05
w_4^*	-231639.30	-3.89	-0.03
w_5^*	640042.26	55.28	-0.02
w_6^*	-1061800.52	41.32	-0.01
w_7^*	1042400.18	-45.95	-0.00
w_8^*	-557682.99	-91.53	0.00
w_9^*	125201.43	72.68	0.01
